

**CHAPTER 2**  
**TABLE OF CONTENTS**

- 2.1.0 Target Population
- 2.2.0 General Guidelines
- 2.3.0 Weight Reduction
- 2.4.0 Macronutrient Composition
  - 2.4.1 Fat
  - 2.4.2 Protein
  - 2.4.3 Carbohydrate
- 2.5.0 Micronutrient Composition
- 2.6.0 Vitamin and Mineral Supplements
- 2.7.0 Nonnutritive Sweeteners
- 2.8.0 Alcohol
- 2.9.0 Healthy Dietary Pattern
- 2.10.0 Physical Activity
- Appendix A. Suggested Approximate Macronutrient Distribution According to Clinical Guideline

## CHAPTER 2.

# Clinical Nutrition Guideline for Overweight and Obese Adults With Type 2 Diabetes (T2D) or Prediabetes, or Those at High Risk for Developing T2D

*Osama Hamdy, MD, PhD; Om P. Ganda, MD, Chair; Clinical Oversight Committee; Melinda Maryniuk, MEd, RD, CDE; Robert A. Gabbay, MD, PhD, FACP; and the members of the Joslin Clinical Oversight Committee*

*From the Adult Diabetes and Clinical Research sections, Joslin Diabetes Center, Harvard Medical School, Boston, Massachusetts*

**Objective.** The Joslin Clinical Nutrition Guideline for Overweight and Obese Adults With Type 2 Diabetes (T2D) or Prediabetes, or Those at High Risk for Developing T2D is designed to assist primary care physicians, specialists, and other healthcare providers in individualizing the care of and setting goals for adult, nonpregnant patients with T2D or individuals at high risk for developing the disease. This guideline focuses on the unique needs of those individuals. Several components complement the *2015-2020 Dietary Guidelines for Americans*. The *Dietary Guidelines for Americans* are jointly developed every 5 years by the US Department of Health and Human Services and the US Department of Agriculture. This Guideline is not intended to replace sound medical judgment or clinical decision making and may need to be adapted for certain patient-care situations where more or less stringent interventions are necessary. This guideline was approved October 19, 2016; updated on January 28, 2018.

### 2.1.0 TARGET POPULATION

**TABLE 1.** Individuals Targeted for Intervention Meet 1 Criterion in Each of 2 Categories

Overweight or Obese <sup>a</sup>	Type 2 Diabetes or Prediabetes
BMI $\geq 25$ kg/m <sup>2</sup> or Waistline Men: $\geq 40$ in (102 cm) [1B] Women: $\geq 35$ in (88 cm) [1B]	Diagnosis of type 2 diabetes (T2D) or prediabetes, determined by IGT or IFG [1A] or High-risk of T2D, determined by: <ul style="list-style-type: none"> <li>• Metabolic syndrome, per AHA/NHLBI criteria [1B]</li> <li>• First-degree relative with T2D</li> <li>• Confirmed diagnosis of insulin resistance (eg, high basal insulin)</li> </ul>

AHA indicates American Heart Association; BMI, body mass index; IGT, impaired glucose tolerance; IFG, impaired fasting glucose; in, inches; NHLBI, National Heart, Lung, and Blood Institute, T2D, type 2 diabetes.

<sup>a</sup>For Asian populations (South Asian Indians, East Asians, and Malays), a BMI  $\geq 23$  kg/m<sup>2</sup> and a waistline  $\geq 35$  in/90 cm in men or  $\geq 31$  in/80 cm in women [1B].

### 2.2.0 GENERAL GUIDELINES

- There is strong evidence that weight reduction improves insulin sensitivity and glycemic control, lipid profile, and blood pressure in T2D, and decreases the

- risk of developing T2D in prediabetes and high-risk populations [1A].
- Refer individuals to a registered dietitian (RD) experienced in diabetes and weight management for individualized medical nutrition therapy (MNT); care should be coordinated with an interdisciplinary team including the patient's primary care physician (PCP) [1B].
  - To enhance effectiveness of MNT, a series of 3 to 4 encounters with an RD, each lasting 45 to 90 minutes, is recommended to begin at diagnosis
- Priorities for this population include:
  - Weight reduction
  - Glycemic control as well as achieving blood pressure and low-density lipoprotein cholesterol goals
  - Meal-to-meal consistency in carbohydrate distribution for those with fixed medication/insulin programs
  - Individualization for cultural and food preferences (eg, vegetarian)
  - Adoption of a healthy eating pattern that is sustainable over time. The Mediterranean diet, the DASH [Dietary Approaches to Stop Hypertension] diet, and a plant-based or vegetarian diet are examples of healthy dietary patterns
  - Integration of behavior-change therapies to adopt healthy eating behaviors and sustainable weight loss
- The meal plan composition, described below, is for general guidance only and may be individualized by the RD or other healthcare provider according to clinical judgment, individual (patient) preferences and needs, and metabolic response.
- Physical activity is an integral component of a weight loss program for both initial weight loss and for weight maintenance.

### 2.3.0 WEIGHT REDUCTION

- A structured lifestyle plan that combines dietary modification, activity, and behavioral modification, along with ongoing support, is necessary for weight reduction [1B]. To maintain long-term weight loss, ongoing weight-maintenance counseling and support is recommended.
- A modest and gradual weight reduction of 1 to 2 pounds every 1 to 2 weeks should be the optimal target [2A]. Reduction of daily caloric intake should be between 250-750 calories [1C]. Total daily intake should not be less than 1000 to 1200 calories for women and 1200 to 1600 calories for men, or based on an RD assessment of usual intake [1C].

- A 5% to 10% weight loss may result in significant improvement in blood glucose control among patients with diabetes and may help prevent the onset of diabetes among individuals with prediabetes [1B]. Weight reduction should be individualized and continued until an agreed-upon BMI and/or other metabolic goals are reached.
- Target individuals should meet with an RD to discuss a structured MNT plan for weight management that includes menus and snacks as well as education and practice in portion control, all effective components of weight-management plans [1B].
- Diabetes-Specific Meal Replacements (DSMRs) in the form of shakes, bars, ready-to-mix powders, and prepackaged meals that match these nutrition guidelines may be effective in initiating and maintaining weight loss.
  - Meal replacements should be used under the supervision of a RD
  - When meal replacements are initiated, glucose levels should be carefully monitored and, if needed, antihyperglycemic medications should be adjusted
  - Meal replacements should be used with caution by those with hyperkalemia
- Bariatric surgeries, although not without medical and nutrition risks, are effective options and may be discussed when indicated (consider in individuals with BMI  $\geq 40$  kg/m<sup>2</sup> and those with BMI  $\geq 35$  kg/m<sup>2</sup> with other comorbidities. Reduce calculations by 2.5 kg/m<sup>2</sup> for Asians) [2B]. To date, there is limited evidence to support the recommendation of bariatric surgeries for patients with BMI  $< 35$  kg/m<sup>2</sup> even if a person has diabetes or other comorbid conditions.
- Anti-obesity medications may be considered for patients who were not able to lose weight through lifestyle modifications, but the long-term risks and benefits of these medications are unclear [2C].
- The effect of diabetes medications should be evaluated throughout the weight loss program and adjusted as necessary to avoid hypoglycemia.

### 2.4.0 MACRONUTRIENT COMPOSITION

#### 2.4.1 Fat:

- Amount.** There is general agreement that the type of fat consumed is more important than the quantity (generally 30% to 40% of total calories). Trans fats from partially hydrogenated oil should be eliminated [1B].
- Monounsaturated and polyunsaturated fats should comprise the majority of fat intake [2B].
  - Limit saturated fat intake to  $< 10\%$  of total calories.
    - Recent evidence demonstrates that saturated fat from dairy foods (milk, yogurt, cheese) may be acceptable within the total daily caloric intake [2B]

- Despite recent evidence suggesting that saturated fat poses a weak or neutral effect on health, further research in this area is warranted

- Low-fat diets are generally less effective than low-carbohydrate diets for weight reduction [2C]

#### **Recommended.**

- Plant fats rich in mono- and polyunsaturated fats (eg, olive oil, canola oil, soybean oil, nuts/seeds, and avocado) [2A]
- Oily fish rich in omega-3 fatty acids (eg, salmon, herring, trout, sardines, fresh tuna) 2 times/week, as a source of these fatty acids [2B]

#### **Not recommended.**

- Foods high in saturated animal fat, including nonlean pork, lamb, and beef; processed meat; butter and cream
- Foods high in trans fats (eg, most fast foods; most commercially baked goods; margarines from partially hydrogenated oil)

### **2.4.2 Protein**

**Amount.** Protein intake should range between 1.0-1.5 grams/kg of adjusted body weight. To calculate adjusted body weight, first calculate excess weight: Excess weight = current weight – ideal body weight (IBW). Adjusted body weight = IBW + 0.25 of excess body weight. This amount generally accounts for 20% to 30% of total caloric intake.

- A modest increase in protein reduces appetite and helps achieve and maintain weight reduction [2B]. Protein also helps minimize loss of lean body mass during weight reduction [2B].
- No reliable scientific data support a protein intake that exceeds 2 grams/kg of adjusted body weight. Conversely, reduction of protein intake to less than 0.8 grams/kg day may result in protein malnutrition.

**Recommended.** Fish, skinless poultry, lean meat, dairy, egg whites, nuts, seeds, soy, and other legumes [2B].

**Not recommended.** Sources of protein that are high in saturated fat (eg nonlean pork, lamb, beef; processed meats) as they may be associated with increased cardiovascular risk [1B]. Heme iron in meat is also associated with an increased risk of T2D [2B].

**Patients with renal issues.** Although reducing total calories may result in a reduction of the total amount of protein intake, any patient with signs of kidney disease (both of the following: proteinuria; estimated glomerular filtration rate <60 ml/min) should consult a nephrologist before increasing the total or percentage of protein in their diet [1B]. Protein intake for these patients should be modified, but not lowered to a level that may jeopardize their overall health or increase their risk for malnutrition or hypoalbuminemia.

### **2.4.3 Carbohydrate**

**Amount.** The total daily intake of carbohydrate should be at least 130 grams/day and preferably 40% to 45% of the total caloric intake. Intake should be adjusted to meet the cultural and food preferences of the individual.

**Consideration of glycemic index/glycemic load.** The glycemic index/glycemic load is an important factor that patients should apply in their daily selection of carbohydrate foods. Foods with a lower glycemic index content should be selected [2B] (eg, whole grains, legumes, fruits, green leafy and nonstarchy vegetables, milk, yogurt).

**Recommended.** Green leafy and nonstarchy vegetables, whole fruits, legumes, whole and minimally processed grains, oats, milk, yogurt [2B].

#### **Not recommended.**

- Sugar, or added sugar, especially sugar-sweetened beverages, ice cream, candies, and grain-based desserts. Milk chocolate should be avoided.
- Refined grain products including white bread, white pasta, white rice, low-fiber wheat cereal, cakes, muffins, pizza. White bagels should be limited.
- High glycemic-index carbohydrates, including white potatoes and white rice.

#### **Fiber.**

- Approximately 14 grams of fiber/1000 cal (20-35 grams) per day is recommended [1C]. If tolerated, approximately 50 grams/day is effective in improving postprandial hyperglycemia; that quantity should be encouraged [2B].
- Fiber from unprocessed plant-based food, such as vegetables, fruits, seeds, nuts, and legumes, is preferable. However, if needed, fiber supplements such as psyllium, resistant starch, and  $\beta$ -glucan can be added [2B].

## **2.5.0 MICRONUTRIENT COMPOSITION**

**Sodium.** Daily consumption should be <2300 mg (about 1 tsp of salt) per day [1A]. Further reduction to 1500 mg is recommended in people aged >50 years, especially those including those with hypertension or chronic kidney disease [2B].

#### **Potassium.**

- Daily consumption should be a minimum of 4700 mg unless potassium excretion is impaired (eg, patients with chronic kidney disease; patients on certain drugs who retain potassium).
- Potassium helps offset high sodium intake by triggering more sodium excretion by the kidneys.
- Potassium-rich foods include fruits and vegetables like bananas, mushrooms, spinach, and almonds.

## **2.6.0 VITAMIN AND MINERAL SUPPLEMENTS**

- In individuals who are not deficient, there are no significant data supporting the routine use of vitamins or minerals to improve glucose control. However,

some individuals may benefit from multivitamin supplementation, as calorie-restricted diets may be inadequate in some nutrients, such as calcium.

- No significant data support the use of herbal supplements or spices to improve glucose control.

### 2.7.0 NONNUTRITIVE SWEETENERS

All FDA-approved nonnutritive sweeteners are permissible in moderate quantities.

### 2.8.0 ALCOHOL

- If alcohol is consumed, consumption must remain moderate: no more than 1 drink per day for women and no more than 2 drinks per day for men (1 drink is equal to 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of 80-proof distilled alcohol).
- Alcoholic beverages contain calories and are low in nutritional value. They may contribute to hypoglycemia or, in the case of high-carbohydrate alcoholic beverages, hyperglycemia.
- It is not advisable to increase alcohol consumption for the purpose of deriving purported health benefits.

### 2.9.0 HEALTHY DIETARY PATTERN

The following dietary patterns have been shown to be effective in the prevention and management of diabetes:

- Mediterranean diet
- DASH diet
- Plant-based, vegetarian, and vegan diets
- Moderately low carbohydrate consumption; high consumption of plant-based protein; fats from plants

The following specific foods have been shown in some study results to be associated with a reduced risk of developing T2D:

- Oat cereal
- Yogurt
- Dairy products
- Tea, coffee, and decaffeinated coffee
- Green leafy vegetables
- Fish and seafood (only in Asia)
- Red grapes, apples, blueberries
- Nuts (especially walnuts)

### 2.10.0 PHYSICAL ACTIVITY

- Physical activity should be an integral component of the weight loss and diabetes care plan to optimize glucose control, decrease cardiovascular risk factors, and achieve or maintain optimal body weight.
- All adults should consult their healthcare provider and/or see an exercise physiologist to discuss a safe exercise program appropriate to their abilities [1C].
- To increase lean body mass, full-body resistance training should be incorporated into the activity plan 3 to 4 days per week. The training should include upper-body,

core, and lower-body strengthening exercises using free weights, resistance machines, or resistance bands [1B].

### Guidelines for healthy adults with diabetes or prediabetes:

- Moderate-intensity aerobic (endurance) physical activity performed a minimum of 30 minutes 5 days per week, or vigorous-intensity aerobic physical activity performed a minimum of 20 minutes 3 days per week, should be achieved unless contraindicated. Activity can be accumulated toward the 30-minute minimum by performing bouts, each lasting 10 or more minutes [1A].
- A target of 60 to 90 minutes of moderate-intensity aerobic activity per day, 6 to 7 days per week, is encouraged for weight loss if overweight or obese [1B].
- Stretching exercises should be done when muscles are warm or at the end of the activity plan to loosen muscles and prevent soreness [1B].

### Additional guidelines for adults with medical or physical limitations:

- Incorporate balance exercises to prevent falls and injuries.
- Functional Fitness Testing is useful to assess patients' functionality and to track their progress. Testing such as 6-Minute Walk Test, 2-Minute Step Test, Balance Assessment and Hand Strength should be included at baseline and post intervention [1C].
- For those with proliferative diabetic retinopathy, retinal traction, or severe nonproliferative diabetic retinopathy, activity programs that involve strenuous lifting; harsh, high-impact components; or components that place the head in an inverted position for extended periods of time may need to be revised, depending on the level of retinopathy and other retinal disease. Consultation with an eye specialist in diabetes eye care is advised.

## Appendix A.

**TABLE 2.** Suggested Approximate Macronutrient Distribution According to Clinical Guideline

Daily Calorie Level	Carbohydrate		Protein		Fat	
	Grams	%	Grams	%	Grams	%
1000	130	~50*	75	30	22	20
1200	135	45	75-90	25-30	40	30
1500	150-170	40-45	75-110	20-30	50	30
1800	180-200	40-45	90-135	20-30	60	30
2000	200-225	40-45	100-150	20-30	70	~30

\*A minimum of 130 grams of carbohydrate per day, in a 1000-calorie meal plan, calculates to ~50% of the total daily calories.

**NOTE:** The diets within the rectangle represent most common diet plans for weight loss.

Source: American Diabetes Association

### CORRESPONDENCE TO:

Om P. Ganda, MD  
1 Joslin Place  
Boston, MA 02215  
Email: om.ganda@joslin.harvard.edu

## REFERENCES

### Recent Studies, Reviews and Meta-Analyses

1. Look AHEAD Research Group; Wing RR, Bolin P, Brancati FL, et al. Cardiovascular effects of intensive lifestyle intervention in type 2 diabetes [published correction appears in *N Engl J Med*. 2014;370(19):1866]. *N Engl J Med*. 2013;369(2):145-154. doi: 10.1056/NEJMoa1212914.
2. Mozaffarian D. Dietary and policy priorities for cardiovascular disease, diabetes and obesity: a comprehensive review. *Circulation*. 2016;133(2):187-225. doi: 10.1161/CIRCULATIONAHA.018585.
3. Ludwig D. Lowering the bar on the low-fat diet. *JAMA*. 2016;316(20):2087-2088. doi: 10.1001/jama.2016.15473.
4. Wang DD, Li Y, Chiuve SE, et al. Association of specific dietary fats with total and cause-specific mortality. *JAMA Intern Med*. 2016;176(8):1134-1145. doi: 10.1001/jamainternmed.2016.2417.
5. de Souza RJ, Mente A, Maroleanu A, et al. Intake of saturated and trans unsaturated fatty acids and risk of all cause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis of observational studies. *BMJ*. 2015;351:h3978. doi: 10.1136/bmj.h3978.
6. Chowdhury R, Warnakula S, Kunutsor S, et al. Association of dietary, circulating, and supplement fatty acids with coronary risk: a systematic review and meta-analysis. *Ann Intern Med*. 2014;160(6):398-406. doi: 10.7326/M13-1788.
7. Franz MJ, Boucher JL, Rutten-Ramos S, VanWormer JJ. Lifestyle weight-loss intervention outcomes in overweight and obese adults with type 2 diabetes: a systematic review and meta-analysis of randomized clinical trials. *J Acad Nutr Diet*. 2015;115(9):1447-1463. doi: 10.1016/j.jand.2015.02.031.
8. Emadian A, Andrews RC, England CY, Wallace C, Thompson JL. The effect of macronutrients on glycaemic control: a systematic review of dietary randomised controlled trials in overweight and obese adults with type 2 diabetes in which there was no difference in weight loss between treatment groups. *Br J Nutr*. 2015;114(10):1656-1666. doi: 10.1017/S0007114515003475.
9. Estruch R, Ros E, Salas-Salvadó J, et al; PREDIMED Study Investigators. Primary prevention of cardiovascular disease with a Mediterranean diet [published correction appears in *N Engl J Med*. 2013;370(9):886]. *N Engl J Med*. 2013;368(14):1279-1290. doi: 10.1056/NEJMoa1200303.
10. Ley SH, Hamdy O, Mohan V, Hu FB. Prevention and management of type 2 diabetes: dietary components and nutritional strategies. *Lancet*. 2014;383(9933):1999-2007. doi: 10.1016/S0140-6736(14)60613-9.
11. Mottalib A, Hamdy O. Effects of 3 different models of medical nutrition therapy on A1C and body weight in overweight and obese patients with type 2 diabetes: a randomized clinical study. *Diabetes*. 2016;65(suppl 1) abstr 759-P.
12. Balk EM, Earley A, Raman G, Avendano EA, Pittas AG, Remington PL. Combined diet and physical activity promotion programs to prevent type 2 diabetes among persons at increased risk: a systemic review for the Community Preventive Services Task Force. *Ann Intern Med*. 2015;163(6):437-451. doi: 10.7326/M15-0452.
13. Batterham RL, Cummings DE. Mechanisms of diabetes improvement following bariatric/metabolic surgery. *Diabetes Care*. 2016;39(6):893-901. doi: 10.2337/dc16-0145.
14. Rubino F, Nathan DM, Eckel RH, et al; Delegates of the 2nd Diabetes Surgery Summit. Metabolic surgery in the treatment algorithm for type 2 diabetes: a joint statement by international diabetes organizations. *Diabetes Care*. 2016;39(6):861-877. doi: 10.2337/dc16-0236.
15. Imamura F, Micha R, Wu JH, et al. Effects of saturated fat, polyunsaturated fat, monounsaturated fat, and carbohydrate on glucose-insulin homeostasis: a systematic review and meta-analysis of randomised controlled feeding trials. *PLoS Med*. 2016;13(7):e1002087. doi: 10.1371/journal.pmed.1002087.
16. Qian F, Korat AA, Malik V, Hu FB. Metabolic effects of monounsaturated fatty acid-enriched diets compared with carbohydrate or polyunsaturated fatty acid-enriched diets in patients with type 2 diabetes: a systematic review and meta-analysis of randomized controlled trials. *Diabetes Care*. 2016;39(8):1448-1457. doi: 10.2337/dc16-0513.
17. Eckel RH, Jakicic JH, Ard JD, et al; American College of Cardiology/American Heart Association Task Force on Practice Guidelines. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines [published correction appears in *Circulation*. 2014;129(25 Suppl 2):S100-S101]. *Circulation*. 2014;129(25 Suppl 2):S76-S99. doi: 10.1161/01.cir.0000437740.48606.d1.
18. Hamdy O, Mottalib A, Morsi A, et al. Long-term effect of intensive lifestyle intervention on cardiovascular risk factors in patients with diabetes in real-world clinical practice: a 5-year longitudinal study. *BMJ Open Diabetes Res Care* 2017; 5(1): e000259. doi:10.1136/bmjdn-2016.000259.

### Other Pertinent References

19. Anderson JW, Gustafson NJ, Bryant CA, Tietyan-Clark J. Dietary fiber and diabetes: a comprehensive review and practical application. *J Am Diet Assoc*. 1987;87(9):1189-1197.
20. Andrews RC, Cooper AR, Montgomery AA, et al. Diet or diet plus physical activity versus usual care in patients with newly diagnosed type 2 diabetes: the Early ACTID randomised controlled trial. *Lancet*. 2011;378(9786):129-139. doi: 10.1016/S0140-6736(11)60442-X.
21. Brinkworth GD, Noakes M, Parker B, Foster P, Clifton PM. Long-term effects of advice to consume a high-protein, low-fat diet, rather than a conventional weight-loss diet, in obese adults with type 2 diabetes: one-year follow-up of a randomised trial. *Diabetologia*. 2004;47(10):1677-1686.
22. Chandalia M, Garg A, Lutjohann D, von Bergmann K, Grundy SM, Brinkley LJ. Beneficial effects of high dietary fiber intake in patients with type 2 diabetes mellitus. *N Engl J Med*. 2000;342(19):1392-1398. doi: 10.1056/NEJM200005113421903.
23. Dansinger ML, Gleason JA, Griffith JL, Selker HP, Schaefer EJ. Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA*. 2005;293(1):43-53.
24. De Caterina R. n-3 fatty acids in cardiovascular disease. *N Engl J Med*. 2011;364(25):2439-2450. doi: 10.1056/NEJMra1008153.
25. Knowler WC, Barrett-Connor E, Fowler SE, et al; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med*. 2002;346(6):393-403. doi: 10.1056/NEJMoa012512.
26. Dumesnil JG, Turgeon J, Tremblay A, et al. Effect of a low-glycaemic index-low-fat-high protein diet on the atherogenic metabolic risk profile of abdominally obese men. *Br J Nutr*. 2001;86(5):557-568.
27. Esposito K, Maiorino MI, Ciotola M, et al. Effects of a Mediterranean-style diet on the need for antihyperglycemic drug therapy in patients with newly diagnosed type 2 diabetes [published correction appears in *Ann Intern Med*. 2009;151(8):591]. *Ann Intern Med*. 2009;151(5):306-314.
28. Foster GD, Wyatt HR, Hill JO, et al. Weight and metabolic outcomes after 2 years on a low-carbohydrate versus low-fat diet: a randomized trial. *Ann Intern Med*. 2010;153(3):147-157. doi: 10.1059/0003-4819-153-3-201008030-00005.
29. Foster GD, Wyatt HR, Hill JO, et al. A randomized trial of a low-carbohydrate diet for obesity. *N Engl J Med*. 2003;348(21):2082-2090. doi: 10.1056/NEJMoa022207.
30. Fukagawa NK, Anderson JW, Hageman G, Young VR, Minaker KL. High-carbohydrate, high-fiber diets increase peripheral insulin sensitivity in healthy young and old adults. *Am J Clin Nutr*. 1990;52(3):524-528. doi: 10.1093/ajcn/52.3.524.
31. Gannon MC, Nuttall FQ. Effect of a high-protein, low-carbohydrate diet on blood



- glucose control in people with type 2 diabetes. *Diabetes*. 2004;53(9):2375-2382.
32. Hamdy O, Horton ES. Protein content in diabetes nutrition plan. *Curr Diab Rep*. 2011;11(2):111-119. doi: 10.1007/s11892-010-0171-x.
  33. Hamdy O, Ledbury S, Mullooly C, et al. Lifestyle modification improves endothelial function in obese subjects with the insulin resistance syndrome. *Diabetes Care*. 2003;26(7):2119-2125. doi: 10.2337/diacare.26.7.2119.
  34. Hu FB, Stampfer MJ, Manson JE, et al. Dietary fat intake and the risk of coronary heart disease in women. *N Engl J Med*. 1997;337(21):1491-1499. doi: 10.1056/NEJM199711203372102.
  35. Hu FB. Plant-based foods and prevention of cardiovascular disease: an overview. *Am J Clin Nutr*. 2003;78(suppl 3):544S-551S.
  36. Johnston CS, Tjonn SL, Swan PD. High-protein, low-fat diets are effective for weight loss and favorably alter biomarkers in healthy adults. *J Nutr*. 2004;134(3):586-591.
  37. Keno Y, Matsuzawa Y, Tokunaga K, et al. High sucrose diet increases visceral fat accumulation in VMH-lesioned obese rats. *Int J Obes*. 1991;15(3):205-211.
  38. Kim JY, Nolte LA, Hansen PA, et al. High-fat diet-induced muscle insulin resistance: relationship to visceral fat mass. *Am J Physiol Regul Integr Comp Physiol*. 2000;279(6):R2057-R2065.
  39. Lara-Castro C, Garvey WT. Diet, insulin resistance, and obesity: zoning in on data for Atkins dieters living in South Beach. *J Clin Endocrinol Metab*. 2004;89(9):4197-4205.
  40. Larsen TM, Dalskov SM, van Baak M, et al; Diets with high or low protein content and glycemic index for weight-loss maintenance. *N Engl J Med*. 2010;363(22):2102-2113. doi: 10.1056/NEJMoa1007137.
  41. Look AHEAD Research Group; Wing RR. Long-term effects of a lifestyle intervention on weight and cardiovascular risk factors in individuals with type 2 diabetes mellitus: four-year results of the Look AHEAD trial. *Arch Intern Med*. 2010;170(17):1566-1575. doi: 10.1001/archinternmed.2010.334.
  42. Ludwig DS, Pereira MA, Kroenke CH, et al. Dietary fiber, weight gain, and cardiovascular disease risk factors in young adults. *JAMA*. 1999;282(16):1539-1546.
  43. Ludwig, D. S. and M. I. Friedman "Increasing adiposity: consequence or cause of overeating?" *JAMA*. 2014; 311(21): 2167-2168.
  44. Luscombe ND, Clifton PM, Noakes M, Parker B, Wittert G. Effects of energy-restricted diets containing increased protein on weight loss, resting energy expenditure, and the thermic effect of feeding in type 2 diabetes. *Diabetes Care*. 2002;25(4):652-657. doi: 10.2337/diacare.25.4.652.
  45. Mann JI. Diet and risk of coronary heart disease and type 2 diabetes. *Lancet*. 2002;360(9335):783-789.
  46. Meckling KA, O'Sullivan C, Saari D. Comparison of a low-fat diet to a low-carbohydrate diet on weight loss, body composition, and risk factors for diabetes and cardiovascular disease in free-living, overweight men and women. *J Clin Endocrinol Metab*. 2004;89(6):2717-2723.
  47. Meneghini LE, Orozco-Beltran D, Khunti K, et al. Weight beneficial treatments for type 2 diabetes. *J Clin Endocrinol Metab*. 2011;96(11):3337-3353. doi: 10.1210/jc.2011-1074.
  48. McAuley KA, Hopkins CM, Smith KJ, et al. Comparison of high-fat and high-protein diets with a high-carbohydrate diet in insulin-resistant obese women [published correction appears in *Diabetologia*. 2005;48(5):1033]. *Diabetologia*. 2005;48(1):8-16.
  49. Park Y, Subar AF, Hollenbeck A, Schatzkin A. Dietary fiber intake and mortality in the NIH-AARP Diet and Health Study. *Arch Intern Med*. 2011;171(12):1061-1068. doi: 10.1001/archinternmed.2011.18.
  50. Parker B, Noakes M, Luscombe N, Clifton P. Effect of a high-protein, high-monounsaturated fat weight loss diet on glycemic control and lipid levels in type 2 diabetes. *Diabetes Care*. 2002;25(3):425-430.
  51. Pereira MA, Swain J, Goldfine AB, Rifai N, Ludwig DS. Effects of a low-glycemic load diet on resting energy expenditure and heart disease risk factors during weight loss. *JAMA*. 2004;292(20):2482-2490.
  52. Raynor HA, Anderson AM, Miller GD, et al; Look AHEAD Research Group. Partial meal replacement plan and quality of the diet at 1 year: Action for Health in Diabetes (Look AHEAD) trial. *J Acad Nutr Dietetics*. 2015;115(5):731-742. doi: 10.1016/j.jand.2014.11.003.
  53. Rock CL, Flatt SW, Sherwood NE, Karanja N, Pakiz B, Thomson CA. Effect of a free prepared meal and incentivized weight loss program on weight loss and weight loss maintenance in obese and overweight women: a randomized controlled trial. *JAMA*. 2010;304(16):1803-1810. doi: 10.1001/jama.2010.1503.
  54. Skov AR, Toubro S, Rønn B, Holm L, Astrup A. Randomized trial on protein vs carbohydrate in ad libitum fat reduced diet for the treatment of obesity. *Int J Obes Relat Metab Disord*. 1999;23(5):528-536.
  55. Sofi F, Cesari F, Abbate R, Gensini GF, Casini A. Adherence to Mediterranean diet and health status: a meta-analysis. *BMJ*. 2008;337:a1344. doi: 10.1136/bmj.a1344.
  56. Stern L, Iqbal N, Seshadri P, et al. The effects of low-carbohydrate versus conventional weight loss diets in severely obese adults: one-year follow-up of a randomized trial. *Ann Intern Med*. 2004;140(10):778-785. doi: 10.7326/0003-4819-140-10-200405180-00007.
  57. Story L, Anderson JW, Chen WJ, Karounos D, Jefferson B. Adherence to high-carbohydrate, high-fiber diets: long-term studies of non-obese diabetic men. *J Am Diet Assoc*. 1985;85(9):1105-1110.
  58. Uusitupa M, Lindi V, Louheranta A, Salopuro T, Lindström J, Tuomilehto J; Finnish Diabetes Prevention Study Group. Long-term improvement in insulin sensitivity by changing lifestyles of people with impaired glucose tolerance: 4-year results from the Finnish Diabetes Prevention Study. *Diabetes*. 2003;52(10):2532-2538. doi: 10.2337/diabetes.52.10.2532.
  59. Yang Q, Liu T, Kuklina EV, et al. Sodium and potassium intake and mortality among US adults: prospective data from the Third National Health and Nutrition Examination Survey. *Arch Intern Med*. 2011;171(13):1183-1191. doi: 10.1001/archinternmed.2011.257.
  60. Gijsbers L, Ding EL, Malik VS, de Goede J, Geleijnse JM, Soedamah-Muthu SS. Consumption of dairy foods and diabetes incidence: a dose-response meta-analysis of observational studies. *Am J Clin Nutr*. 2016;103(4):1111-1124. doi: 10.3945/ajcn.115.123216.
  61. Ericson U, Hellstrand S, Brunkwall L, et al. Food sources of fat may clarify the inconsistent role of dietary fat intake for incidence of type 2 diabetes. *Am J Clin Nutr*. 2015;101(5):1065-1080. doi: 10.3945/ajcn.114.103010.
  62. Tobias DK, Chen M, Manson JE, Ludwig DS, Willett W, Hu FB. Effect of low-fat diet interventions versus other diet interventions on long-term weight change in adults: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol*. 2015;3(12):968-975. doi: 10.1016/S2213-8587(15)00367-8.