
...BIBLIOGRAPHY...

- Boers M, Verhoeven AC, Markusse HM, et al. Randomised comparison of combined step-down prednisolone, methotrexate and sulphasalazine with sulphasalazine alone in early rheumatoid arthritis. *Lancet* 1998;351:220.
- Brennan FM, Maini RN, Feldmann M. TNF α —A pivotal role in rheumatoid arthritis? *Br J Rheumatol* 1992;31:293-298.
- Elliott MJ, Maini RN, Feldmann M, et al. Randomised double-blind comparison of chimeric monoclonal antibody to tumour necrosis factor alpha (cA2) versus placebo in rheumatoid arthritis. *Lancet* 1994;344:1105-1110.
- Feldmann M, Brennan FM, Maini RN. *Rheumatoid arthritis*. *Cell* 1996;85:307-310.
- Felson DT, Anderson JJ, Boers M, et al. American College of Rheumatology preliminary definition of improvement in rheumatoid arthritis. *Arthritis Rheum* 1995;38:727-735.
- Gardiner PV, Sykes HR, Hassey GA, Walker DJ. An evaluation of the Health Assessment Questionnaire in long-term longitudinal follow-up of disability in rheumatoid arthritis. *Br J Rheumatol* 1993;32:724-728.
- Gran JT, Husby G, Thorsby E. The association between rheumatoid arthritis and the HLA antigen DR4. *Ann Rheum Dis* 1983;42:292-296.
- Harris ED Jr. Clinical features of rheumatoid arthritis. In: Kelley WN, Harris ED Jr, Duddy S, Sledge CB. *Textbook of Rheumatology*, 4th Ed. Philadelphia, PA: WB Saunders Company; 1983.
- Hawley DJ, Wolfe F. Sensitivity to change of the health assessment questionnaire (HAQ) and other clinical and health status measures in rheumatoid arthritis: Results of short-term clinical trials and observational studies versus long-term observational studies. *Arthritis Care Res* 1992;5:130-136.
- Lawrence RC, Helmick CG, Arnett FC, et al. Estimates of the prevalence of arthritis and selected musculoskeletal disorders in the United States. *Arthritis Rheum* 1998;41:778-799.
- Lipsky P, St. Clair W, Kavanaugh A, et al. Long-term control of signs and symptoms of rheumatoid arthritis with chimeric monoclonal anti-TNF α antibody (infliximab) in patients with active disease on methotrexate. *Arthritis Rheum* 1998;41(Suppl 9):S364.
- Mandell BF. COX 2-selective NSAIDs: biology, promises, and concerns. *Cleve Clin J Med* 1999;66:285-292.
- Matsuda Y, Yamanaka H, Higami K, Kashiwazaki S. Time lag between active joint inflammation and radiological progression in patients with early rheumatoid arthritis. *J Rheumatol* 1998;25:427-432.
- Moreland LW, Baumgartner SW, Schiff MH, et al. Treatment of rheumatoid arthritis with a recombinant human tumor necrosis factor receptor (p75)-Fc fusion protein. *New Engl J Med* 1997;337:141-147.
- Mottonen T, Hannonen P, Leirisalo-Repo M, et al. Comparison of combination therapy with single-drug therapy in early rheumatoid arthritis: A randomized trial. *Lancet* 1999;353:1568-1573.
- O'Dell J. Combination DMARD therapy for rheumatoid arthritis: Apparent universal acceptance. [Abstract] *Arthritis Rheum* 1997;40(Suppl 9):S50.
- O'Dell JR, Haire CE, Erikson N, et al. Treatment of rheumatoid arthritis with methotrexate alone, sulfasalazine and hydroxychloroquine, or a combination of all three medications. *N Engl J Med* 1996;334:1287-1291.
- O'Dell JR, Nepom BS, Haire C, et al. HLA-DRB1 typing in rheumatoid arthritis: Predicting response to specific treatments. *Ann Rheum Dis* 1998;57:209-213.
- Pincus T, Brooks RH, Callahan LF. Prediction of long-term mortality in patients with rheumatoid arthritis according to simple questionnaire and joint count measures. *Ann Intern Med* 1994;120:26-34.
- Prashker MJ, Meenan RF. The total costs of drug therapy for rheumatoid arthritis: A model based on costs of drug, monitoring, and toxicity. *Arthritis Rheum* 1995;38:318-325.
- Semble EL, Loeser RF, Wise CM. Therapeutic exercise for rheumatoid arthritis and osteoarthritis. *Semin Arthritis Rheum* 1990;20:32-40.
- Singh G, Terry R, Ramey D, Wolfe F, Fries J. Cost of medical care for patients with osteoarthritis and rheumatoid arthritis: A 13 year study. [Abstract]. *Arthritis Rheum* 1996;39:S71.
- Van der Heide A, Jacobs JWG, Bijlsma WJ, et al. The effectiveness of early treatment with "second-line" antirheumatic drugs: A randomized, controlled trial. *Ann Int Med* 1996;124:699-707.

...BIBLIOGRAPHY...

- Ward MM, Leigh JP, Fries JF. Progression of functional disability in patients with rheumatoid arthritis: Associations with rheumatology subspecialty care. *Arch Intern Med* 1993;153:2229-2237.
- Ware JE Jr, Bayliss MS, Rogers WH, Kosinski M, Tarlov AR. Differences in 4-year health outcomes for elderly and poor chronically ill patients treated in HMO and fee-for-service systems. Results from the Medical Outcomes Study. *JAMA* 1996;276:1039-1047.
- Weinblatt ME, Kremer JM, Bankhurst AD, et al. A trial of etanercept, a recombinant tumor necrosis factor receptor: Fc fusion protein, in patients with rheumatoid arthritis receiving methotrexate. *N Engl J Med* 1999; 340:253-259.
- Weinblatt ME, Kremer JM, Coblyn JS, et al. Leflunomide plus methotrexate in refractory RA: A pilot study. [Abstract] *Arthritis Rheum* 1997;40(Suppl 9): S193.
- Wolfe F, Hawley DJ. The long-term outcomes of rheumatoid arthritis: Work disability—A prospective 18 year study of 823 patients. *J Rheumatol* 1998;25:2108-2117.
- Wolfe F, Sharp JT. Radiographic outcome of recent-onset rheumatoid arthritis: a 19-year study of radiographic progression. *Arthritis Rheum* 1998;41:1571-1582.
- Wolfe F, Zwillich SH. The long-term outcomes of rheumatoid arthritis: a 23-year prospective, longitudinal study of total joint replacement and its predictors in 1,600 patients with rheumatoid arthritis. *Arthritis Rheum* 1998;41:1072-1082.
- Wolfe F. Psychological distress and rheumatic disease. *Scand J Rheumatol* 1999;28:131-136.
- Yelin E, Henke C, Epstein W. The work dynamics of the persons with rheumatoid arthritis. *Arthritis Rheum* 1987;30:507-512.
- Yelin E. The costs of rheumatoid arthritis: Absolute, incremental, and marginal estimates. *J Rheumatol* 1996;23:47-2351.
- Yelin EH, Criswell LA, Feigenbaum PG. Health care utilization and outcomes among persons with rheumatoid arthritis in fee-for-service and prepaid group practice settings. *JAMA* 1996;276:1048-1053.
- Yelin EH, Such CL, Criswell LA, Epstein WV. Outcomes for persons with rheumatoid arthritis with a rheumatologist versus a non-rheumatologist as the main physician for this condition. *Med Care* 1998;36:513-522.
- Ytterberg SR, Mahowald ML, Krug HE. Exercise for arthritis. *Balliere's Clin Rheumatol* 1994;8:161-187.