

**2020
Official Abstracts**



May 2020 • Volume 22, Supplement 2

International Journal of **MS**CARE

The Official Publication of the Consortium of Multiple Sclerosis Centers



***Abstracts from the
34th Annual Meeting of the
Consortium of Multiple Sclerosis Centers***

May – June 2020

Visit www.ms-care.org to view archived meeting content.

ijmsc.org

EDITORIAL BOARD

Editor in Chief

Francois Bethoux, MD
Cleveland Clinic
Cleveland, Ohio, USA

Associate Editor

Mary Alissa Willis, MD
University of Mississippi Medical Center
Jackson, Mississippi, USA

Ex Officio

June Halper, MSN, ANP, FAAN
Hackensack, New Jersey, USA

Project Manager

Charlene Belsole, CCRP
Cleveland Clinic
Cleveland, Ohio, USA

Board Members

Ted Brown, MD, MPH
Evergreen Neuroscience Institute and
Medical Center

Kirkland, Washington, USA

Susan Coote, PT, PhD
University of Limerick
Limerick, Ireland

Mary Filipi, PhD, APRN-C
Wahoo, Nebraska, USA

Marcia Finlayson, PhD, OT Reg (Ont), OTR
Queen's University
Kingston, Ontario, Canada

Frederick W. Foley, PhD
Yeshiva University
Bronx, New York, USA

Holy Name Medical Center
Teaneck, New Jersey, USA

Kathleen Fuchs, PhD
University of Virginia Health System
Charlottesville, Virginia, USA

Eduard Gappmaier, PT, PhD
University of Utah
Salt Lake City, Utah, USA

Christoph Heesen, MD
Institute of Neuroimmunology and
Clinical MS Research
University Medical Center
Hamburg, Germany

James Marriot, MD, FRCPC
University of Manitoba
Winnipeg, Manitoba, Canada

Lori Mayer, MSN, DNP(s), MSCN, CCRP
MS Clinic of Central Texas
Round Rock, Texas, USA

Sarah Morrow, MD, MSc, FRCPC
University of Western Ontario
London Health Sciences Centre
London, Ontario, Canada

Ahmed Z. Obeidat, MD, PhD
Medical College of Wisconsin
Milwaukee, WI, USA

Jacob J. Sosnoff, PhD
University of Illinois at Urbana-Champaign
Urbana, Illinois, USA

Megan Weigel, DNP, ARNP-c
Baptist Medical Center
Jacksonville Beach, Florida, USA

Publishers

Joseph J. D'Onofrio
Frank M. Marino
Delaware Media Group
PO Box 937
Glen Rock, NJ 07452, USA
jdonofrio@delmedgroup.com

Managing Editor

Anna Gillette

Art Director

James Ticchio

2020 Virtual Annual Meeting of the CMSC

The editorial team is pleased to present this supplement to the *International Journal of MS Care* (IJMSC) containing the abstracts from the 2020 Virtual Annual Meeting of the Consortium of Multiple Sclerosis Centers (CMSC). These abstracts include platform, poster, and Whitaker Research Track presentations, as well as a few late-breaking abstracts. In these unusual times, we all have had to modify many of our activities due to the COVID-19 pandemic, including the introduction of virtual scientific meetings. While the CMSC generously offers free registration to this meeting, many health care providers may not have the time to attend the live online sessions. The print version of this supplement is being distributed to members of the CMSC. The electronic version will be available to all on the IJMSC website at ijmsc.org.

We would like to thank Sanofi Genzyme for their support, which made this publication possible.

While reading abstracts does not completely replace in-person networking, our team hopes that this supplement will attest to the fact that the desire to share innovation and research has not been extinguished by the pandemic. We hope that you and your loved ones are keeping safe in these challenging times.

—Francois Bethoux, MD
Editor in Chief



CMSC Abstract Review Committee

CO-CHAIRS

Amy Perrin Ross, APN, MSN, CNRN, MSCN

Loyola University of Chicago MS Clinic
Chicago, IL
aperrinros@aol.com

Carrie Hersh, DO, MSc

Mellen Program for Multiple Sclerosis
Lou Ruvo Center for Brain Health
Cleveland Clinic Nevada
Las Vegas, NV
hershc@ccf.org

REVIEWERS

Farren Briggs, PhD, ScM

Case Western Reserve University
Cleveland, OH
farren.briggs@case.edu

Dorothea Cassidy-Pfohl, RN, BS, MSCN

Bala Cynwyd, PA
dcpfohl@verizon.net

Kathleen Czuba, SLP

Loyola University Medical Center
Oak Brook Terrace, IL
kateczuba@gmail.com

Robert Fallis, MD, MSCS

Ohio State University
Columbus, OH
robert.fallis@osumc.edu

Mary Kay Fink, RN, MSN, ACNS-BC, MSCN

John L. Trotter Multiple Sclerosis Center
St. Louis, MO
marykayfink@wustl.edu

Cindy Gackle, OTR/L, MSCS

Minneapolis, MN
cgackle@cptelecom.us

Ajay Gupta, MD, MSCS

Fort Wayne Neurological Center
Fort Wayne, IN
agupta@fwnc.com

Colleen Harris, MN, NP, MSCN, MSCS

University of Calgary MS Clinic
Calgary, AB, Canada
colleen.harris@albertahealthservices.ca

Rock Heyman, MD

University of Pittsburgh Medical Center
Pittsburgh, PA
heyhra@upmc.edu

Gloria Hou, MD

University of Washington
Seattle, WA
ghou1@u.washington.edu

Sue Kushner, MS, PT

Slippery Rock University
Valencia, VA
susan.kushner@sru.edu

Beverly Layton, RN, BSN, CCRC, MSCN

Birmingham VA Medical Center
Birmingham, AL
blayton55@bellsouth.net

Deborah Miller, PhD, LISW-S

Mellen Center for MS Treatment and Research
Cleveland, OH
millerd@ccf.org

Kelly Morton, PharmD

Loyola University of Chicago
MS Clinic
Chicago, IL
Kelly.Morton@lubs.org

Nancy Navarro, APN

Suburban Physicians
Aurora, IL
N.Navarro1@comcast.net

Cynthia Sullivan, PhD

Neuropsychology Associates of Fairfax, LLC
Fairfax, VA
csullivan@neuropsychologyfairfax.com

Gloria von Geldern, MD

University of Washington Medical Center
Seattle, WA
vgeldern@uw.edu

Jeff Wilken, PhD

Neuropsychology Associates of Fairfax, LLC
Fairfax, VA
jwilken@neuropsychologyfairfax.com

Annette Wundes, MD

Western MS Center at UWMC / MSRRTC
Seattle, WA
awundes@u.washington.edu

STAFF

Rachelle U. Ramirez

Multimedia Manager
Consortium of MS Centers
Hackensack, NJ
rramirez@mscare.org

Jennifer Pichardo

Project Coordinator
Consortium of MS Centers
Hackensack, NJ
jpichardo@mscare.org

PLATFORMS**DISEASE ASSESSMENT AND MANAGEMENT**

- (DAM01) International Registry Tracking Pregnancy Outcomes in Women Treated with Dimethyl Fumarate
- (DAM02) Menarche and Relapses in Girls with Pediatric Multiple Sclerosis
- (DAM03) Complexity of Aging with Multiple Sclerosis: Graceful Concessions or Kicking and Screaming?
- (DAM04) Serum Glial Fibrillary Acidic Protein Is Elevated in a Subset of Neuromyelitis Optica Patients and Associated with Increased Risk of Attacks
- (DAM05) Machine Learning Algorithms Applied to Visual Metrics to Classify Demyelinating Disease Diagnosis in Children
- (DAM06) Updated Recommendations for a Standardized Magnetic Resonance Imaging Protocol for Multiple Sclerosis

DISEASE-MODIFYING THERAPY

- (DMT01) Comparative Effectiveness of Switching from Natalizumab to a Moderate- Versus High-Efficacy Disease-Modifying Therapy in Clinical Practice
- (DMT02) Yearly Efficacy and Safety Outcomes Over 4 Years After Last Alemtuzumab Course in Pooled CARE-MS I and II Patients by Number of Additional Courses Received Through Year 9
- (DMT03) Efficacy and Safety of Ofatumumab Versus Teriflunomide in Patients with Relapsing Multiple Sclerosis: Phase 3 ASCLEPIOS I and II Trials
- (DMT04) Treatment-Emergent Adverse Events Occurring Early in the Treatment Course of Cladribine Tablets in Two Phase 3 Trials in Multiple Sclerosis

PSYCHOSOCIAL: COGNITION

- (PSY01) A Mindfulness Group Intervention in Newly Diagnosed Persons with Multiple Sclerosis: A Pilot Study
- (PSY02) Effects of Weekly Participation in a Wellness Program on Self-Reported Measures for People Living with Multiple Sclerosis: A 3-Year Analysis
- (PSY03) Examining Multilevel Environmental Correlates of Physical Activity Among Older Adults with Multiple Sclerosis
- (PSY04) Effect of Nabiximols Cannabinoid Oromucosal Spray on Depressive Symptoms, Suicidality, and Cognition in Patients with Multiple Sclerosis
- (PSY06) Multiple Sclerosis Management: Predicting Disease Trajectory of Multiple Sclerosis on Multidimensional Data Including Digital Cognitive Assessments and Patient-Reported Outcomes Using Machine Learning Techniques

REHABILITATION

- (RHI01) Significant Structural Neuroplasticity Changes Can Follow Physical Behavioral Change Therapy for Multiple Sclerosis
- (RHI02) A New Look at the Symbol Digit Modalities Test in Multiple Sclerosis and Disabilities
- (RHI03) Creating a Yoga Program as Part of a Comprehensive Multiple Sclerosis Care Model
- (RHI04) Feasibility of "Sit Less, Move More": An Intervention for Reducing Sedentary Behavior Among African Americans with Multiple Sclerosis
- (RHI05) The Effect of Aerobic Fitness on Physical and Cognitive Function and Brain Volume in Older Adults with Multiple Sclerosis
- (RHI06) Functional Electrical Stimulation Cycling Exercise Reduces Lower Limb Strength Asymmetry in Persons with Multiple Sclerosis

POSTERS**COMPLEMENTARY AND ALTERNATIVE THERAPIES**

- (CAM01) Multiple Sclerosis Imbalance: Visual Rehabilitation
- (CAM02) Acupuncture and Electromagnetotherapy for Chronic Pain Relief in Multiple Sclerosis
- (CAM03) The Effects of Reflexology in People with Multiple Sclerosis
- (CAM04) The Effects of CBD:THC Tincture Oil in Reducing Symptoms and Overall Symptom Management Medication Dosages, in Persons with Multiple Sclerosis
- (CAM05) Challenges and Opportunities in Progressive Multiple Sclerosis Trials: Lessons from Lipoic Acid
- (CAM06) Exercise in Medicine: A Complementary Exercise Promotion Approach Within Comprehensive Multiple Sclerosis Care
- (CAM07) Changes in Dietary Habits of Individuals Living with Multiple Sclerosis Enrolled in a Day Wellness Program

CASE REPORTS/CASE SERIES

- (CRS01) Seasonal Variation and Other Observations in Myelin Oligodendrocyte Glycoprotein (MOG) Antibody-Associated Disease

- (CRS02) Multiple Surgeries and Misdiagnosis Before Multiple Sclerosis Diagnosis: A Case Report
- (CRS03) Head Trauma as Onset for Multiple Sclerosis Diagnosis: A Case Report
- (CRS04) Team Approach Yields Surprising Functional Progress and Quality-of-Life Changes in a Challenging Case of Neuromyelitis Optica
- (CRS05) Differential Diagnosis and Treatment of Tumefactive Demyelination in a Teenaged Girl
- (CRS06) A Long-Standing Case of Recurrent Transverse Myelitis Due to Myelin Oligodendrocyte Glycoprotein (MOG)-IgG Antibody Mimicking Multiple Sclerosis
- (CRS07) Case Report of Severe Multiple Sclerosis Relapse Due to B-Cell Reconstitution Post Alemtuzumab
- (CRS08) Demographics, Clinical Characteristics, and Outcomes of Myelin Oligodendrocyte Glycoprotein (MOG) Antibody Disease Followed Up at Washington University in St. Louis
- (CRS09) A Fatal Case of Alemtuzumab-Induced Immune Thrombocytopenic Purpura in a Patient with Relapsing Multiple Sclerosis
- (CRS10) Colitis Associated with Teriflunomide
- (CRS11) Remarkable Recovery of Fulminant Multiple Sclerosis After Treatment Induction with Cyclophosphamide
- (CRS12) Neurofibromatosis Type 1 and Multiple Sclerosis in the Same Patient
- (CRS13) Successful Use of Immunotherapy for Osmotic Demyelination Syndrome

DISEASE-MODIFYING THERAPY

- (DXT01) Maintenance of Working Status and Work Productivity in Persons with Multiple Sclerosis Treated with Dimethyl Fumarate: A 5-Year Analysis of the North American Research Committee on Multiple Sclerosis (NARCOMS) Registry
- (DXT02) Early Effect of Ofatumumab on B-Cell Counts and Magnetic Resonance Imaging Activity in Relapsing Multiple Sclerosis Patients: Results from the APLIOS Study
- (DXT03) Analyses of the Effect of Disease Duration on the Efficacy and Safety of Siponimod in Patients with Active Secondary Progressive Multiple Sclerosis from the EXPAND Study
- (DXT04) Siponimod First-Dose Effects in Patients with Secondary Progressive Multiple Sclerosis Receiving Concomitant Selective Serotonin Reuptake Inhibitor Therapy
- (DXT05) Efficacy of Diroximel Fumarate in Highly Active Relapsing-Remitting Multiple Sclerosis: Interim Results from the Phase 3 EVOLVE-MS-1 Study
- (DXT06) Real-World Effectiveness of Peginterferon Beta-1a Versus Interferon Beta-1a and Glatiramer Acetate in US Multiple Sclerosis Patients
- (DXT07) Injection Site Reactions and Risk of Discontinuation Among New and Experienced Peginterferon Beta-1a Users in the Plegridy Observational Program
- (DXT08) Post Hoc Analysis of Efficacy of Cladribine Tablets in Patients with Relapsing-Remitting Multiple Sclerosis Diagnosed Within 3 or 4 Years Prior to the CLARITY Study
- (DXT09) Exploration of Factors Which Influence Treatment Decisions of Patients with Multiple Sclerosis
- (DXT10) Siponimod Affects Disability Progression in Patients with Secondary Progressive Multiple Sclerosis Independent of Relapse Activity: Results from the Phase 3 EXPAND Study
- (DXT11) The Implications of Suboptimal Treatment Outcomes with Disease-Modifying Drugs in Employees with Multiple Sclerosis
- (DXT12) Real-World Effectiveness of Peginterferon Beta-1a Versus Teriflunomide in US Multiple Sclerosis Patients
- (DXT13) Disease-Modifying Therapies: How Confident Are We That We Understand Their Risk?
- (DXT14) Long-Term Safety and Efficacy of Eculizumab in Neuromyelitis Optica Spectrum Disorder
- (DXT15) Inebilizumab Reduces Neuromyelitis Optica Spectrum Disorder Disability Worsening: Outcomes and Long-Term Follow-up Data from the N-MOMentum Trial
- (DXT16) Effectiveness of Delayed-Release Dimethyl Fumarate Relative to Duration of Prior Glatiramer Acetate in Patients Enrolled in the RESPOND Study
- (DXT17) Long-term Follow-up Results from the Phase 2 Multicenter Study of Ublituximab (UTX), a Novel Glycoengineered Anti-CD20 Monoclonal Antibody, in Patients with Relapsing Multiple Sclerosis
- (DXT18) Adherence and Compliance with Subcutaneous Administration of Ofatumumab in Relapsing Multiple Sclerosis

- (DXT20) Glatiramer Acetate (GA) Produced by Mapi Pharma Is Equivalent to Commercially Available GA Preparations
- (DXT22) Characterization of Incidence and Time-to-Recovery from Grade 3/4 Lymphopenia Lasting ≥ 6 Months in Patients with Multiple Sclerosis Treated with Cladribine Tablets
- (DXT23) Disease-Modifying Therapy Landscape: An Evaluation of Cost and Care
- (DXT24) Two Expanded Disability Status Scale Subscales Evaluated in Patients with Relapsing-Remitting or Secondary Progressive Multiple Sclerosis
- (DXT26) Long-Term Disease Stability Assessed by the Expanded Disability Status Scale in Patients Treated with Cladribine Tablets in the CLARITY and CLARITY Extension Studies
- (DXT27) Integrated Lymphopenia Analysis in Younger and Older Patients with Multiple Sclerosis Treated with Cladribine Tablets
- (DXT28) Effectiveness of Cladribine Tablets in Patients with Relapsing-Remitting Multiple Sclerosis with Baseline Expanded Disability Status Scale Score ≥ 3.5 or ≤ 3.0 in CLARITY
- (DXT29) ACAPELLA: Real-World Experience with Ocrelizumab: An Observational Study Evaluating Safety in Patients with Relapsing and Progressive Multiple Sclerosis, Year 3 Data
- (DXT30) ACAPELLA: Hypogammaglobulinemia and JC Virus Status in Ocrelizumab-Treated Patients, Year 2 Data
- (DXT31) Impact of Eculizumab on Hospitalization Rates and Relapse Treatment in Patients with Neuromyelitis Optica Spectrum Disorder: Phase 3 PREVENT Study
- (DXT33) ACAPELLA: B-Cell Reconstitution in Ocrelizumab-Treated Patients
- (DXT34) Revealing the Immune Cell Subtype Reconstitution Profile in Cladribine-Treated Patients at the 96-Week Timepoint (CLARITY) Using Deconvolution Algorithms
- (DXT35) Real-World Experience with Ocrelizumab: A Safety Analysis
- (DXT36) Effect of Evobrutinib, a Bruton's Tyrosine Kinase Inhibitor, on Immune Cell and Immunoglobulin Levels over 48 Weeks in a Phase 2 Study in Relapsing Multiple Sclerosis
- (DXT37) Effect of Teriflunomide on Brain Volume Loss in Patients with Relapsing Multiple Sclerosis of Differing Ages in TEMSO
- (DXT38) Effects of Ozanimod on Information Processing Speed: Findings from the Phase 3 SUNBEAM and DAYBREAK Extension Trials
- (DXT40) Effect of the S1P1/5 Receptor Modulator Ozanimod on Cognitive Processing Speed in Subjects with Relapsing Multiple Sclerosis: Design of the ENLIGHTEN Study
- (DXT41) Eculizumab Benefits a Broad Range of Patients with Aquaporin-4 Antibody-Positive Neuromyelitis Optica Spectrum Disorder: The Phase 3 PREVENT Study
- (DXT42) Rationale and Design of CLASSIC-MS Study Evaluating Long-Term Efficacy for Patients with Multiple Sclerosis Treated with Cladribine Tablets
- (DXT43) Analyses of the Effect of Baseline Age on the Efficacy and Safety of Siponimod in Patients with Active Secondary Progressive Multiple Sclerosis from the EXPAND Study
- (DXT44) Real-World Patterns of Disease Progression in Patients with Multiple Sclerosis Who Are Adherent Versus Nonadherent to Disease-Modifying Treatments over 6 Years
- (DXT45) Pharmacist-Based Intervention for Improving Baseline Laboratory Monitoring for Patients on Multiple Sclerosis Disease-Modifying Therapies
- (DXT46) Cognitive Performance and Disability Across Age Groups in Teriflunomide-Treated Patients in the Teri-PRO Study
- (DXT48) Efficacy of Subcutaneous Interferon Beta-1a in Patients with a First Clinical Demyelinating Event: REFLEX Study – Outcomes in Patients Stratified by 2017 McDonald Criteria
- (DXT49) Post Hoc Analysis of Efficacy of Cladribine Tablets in Patients with Relapsing-Remitting Multiple Sclerosis Aged Over and Under 30 Years in the CLARITY Study
- (DXT50) Prevalence of Serious Adverse Pregnancy Outcomes After Exposure to Interferon Beta Before or During Pregnancy: Stratification by Characteristics of Pregnant Women with Multiple Sclerosis in a Register-Based Cohort Study in Finland and Sweden
- (DXT51) High Rates of Adherence to Oral Diroximel Fumarate and Dimethyl Fumarate Are Observed and Sustained in Relapsing Multiple Sclerosis Patients
- (DXT52) Efficacy and Safety of Eculizumab in Patients with Neuromyelitis Optica Spectrum Disorder Previously Treated with Rituximab: The Phase 3 PREVENT Study
- (DXT53) Multiple Sclerosis Clinical Phenotypes: Using Technology to Educate Patients and Optimize Treatment
- (DXT54) Assessment of the Discontinuation Rates of Disease-Modifying Therapy in Veterans with Multiple Sclerosis
- (DXT55) Herpes Zoster Virus (HZV) Infections Among Multiple Sclerosis Patients Treated with Various Disease-Modifying Therapies
- (DXT56) Potential Weight Changes Among Patients with Multiple Sclerosis Undergoing Treatment with Ocrevus (Ocrelizumab)
- (DXT57) FAST: Faster and Safe Administration of Tysabri
- (DXT58) Reduction of Risk of Secondary Progressive Multiple Sclerosis within 2 Years of Treatment with Cladribine Tablets: An Analysis of the CLARITY Study
- (DXT59) The CLARITY Study: Efficacy Outcomes Among Patients Who Received Disease-Modifying Drugs Prior to Treatment with Cladribine Tablets
- (DXT60) Correlations Between Four Common Measures of Cognition in Patients with Secondary Progressive Multiple Sclerosis
- (DXT61) Injection-Related Reactions with Subcutaneous Administration of Ofatumumab in Relapsing Multiple Sclerosis: Pooled Analysis of the Phase 3 ASCLEPIOS I and II Trials
- (DXT62) Real-World Treatment Patterns in Patients with Multiple Sclerosis Using Disease-Modifying Therapies
- (DXT63) Associations Between Treatment Satisfaction, Medication Beliefs, and Adherence to Disease-Modifying Therapies in Patients with Multiple Sclerosis Among Adult Saudis: A Tertiary Care Center Experience
- (DXT64) Evaluation of Rituximab Regimens and Outcomes in Neuromyelitis Optica Patients from a Single Academic Medical Center: A Retrospective Chart Review
- (DXT65) Longitudinal Disability Follow-up in Patients with 6-Month Confirmed Disability Improvement or Worsening in the CARE-MS and Extension Studies
- (DXT66) Clinical Benefits of Eculizumab Monotherapy in Neuromyelitis Optica Spectrum Disorder: Findings from the Phase 3 PREVENT Study
- (DXT67) Cognitive Functions over the Course of 5 Years in Multiple Sclerosis Patients Treated with Disease-Modifying Therapies
- (DXT69) One-Year Interim Analysis of Real-World Patient-Reported Outcomes in Relapsing-Remitting Multiple Sclerosis Patients Transitioning to Alemtuzumab (PRO-ACT Study)
- (DXT70) Clinical Characteristics and Outcomes of Peginterferon Beta-1a Treatment by Age: A Subgroup Analysis of the Plegridy Observational Program
- (DXT71) Efficacy and Safety of Teriflunomide in Patients with Relapsing-Remitting Multiple Sclerosis of Varying Disease Duration: Analysis of Pooled Clinical Trials
- (DXT73) Updated Safety of Cladribine Tablets in the Treatment of Patients with Multiple Sclerosis: Integrated Safety Analysis and Postapproval Data
- (DXT74) An Analysis of the Relationship Between Cladribine Dose and Risk of Malignancies in Patients with Multiple Sclerosis
- (DXT75) Switches to Established and Recently Approved Oral Disease-Modifying Therapies: Comparison of Patient Clinical Profiles and Therapy Selection Drivers
- (DXT76) First-Line Ocrelizumab Use for Relapsing-Remitting Multiple Sclerosis in the United States: Trend and Comparison to Glatiramer Acetate and Dimethyl Fumarate
- (DXT77) Alemtuzumab Maintains Efficacy on Clinical and Magnetic Resonance Imaging Lesion Outcomes, Including Slowing of Brain Volume Loss, Over 9 Years in Relapsing-Remitting Multiple Sclerosis Patients: CARE-MS II Follow-up (TOPAZ Study)
- (DXT78) The FLUENT Study: Changes in Immune Cell Profile, and in Clinical and Safety Outcomes, in Fingolimod-Treated Patients with Relapsing Multiple Sclerosis
- (DXT79) Efficacy of Ocrelizumab Treatment on Cognitive Functions in Persons with Multiple Sclerosis

EPIDEMIOLOGY AND GENETICS

- (EPI01) Determining the Effect of Early Versus Later Diagnosis of Multiple Sclerosis on Long-Term Prognosis in a Real-World Setting
- (EPI02) Motor Impairment in Multiple Sclerosis: Analysis from the North American Registry for Care and Research in Multiple Sclerosis (NARCRMS)
- (EPI03) Increase in Family Recurrence in Patients Diagnosed with Multiple Sclerosis in the Years 2017-2019 in Hispanic Population of Puerto Rico
- (EPI04) Diet Quality and Nutritional Adequacy of Micronutrients Among People with Relapsing-Remitting Multiple Sclerosis: An Analysis of Weighed Food Records

FAMILY AND CAREGIVERS

- (FAM01) Characterizing Predictors of Resilience Among Family Caregivers of People with Advanced Multiple Sclerosis Disability: Work in Progress
- (FAM02) Understanding Units of Energy: Key to Communicating Multiple Sclerosis Fatigue

INTERNET AND INFORMATION SERVICES

- (IIS01) North American Registry for Care and Research in Multiple Sclerosis (NARCRMS) Model for Implementing OpenClinica Insight for Data Sharing and Visualization

IMAGING

- (IMG01) Conformance to CMSC Magnetic Resonance Imaging (MRI) Guidelines in a Real-World Multicenter MRI Dataset
- (IMG02) Optic Nerve Head Volume Is Significantly Decreased in Pediatric Multiple Sclerosis and Not Pediatric Myelin Oligodendrocyte Glycoprotein (MOG)-Related Disorders
- (IMG03) Cerebellar Connectivity Is Associated with Verbal Memory Impairment in Multiple Sclerosis
- (IMG04) The Association Between Magnetic Resonance Imaging Brain Volumes and Computerized Cognitive Scores of People with Multiple Sclerosis
- (IMG05) Analysis of Demyelinating Injuries in Magnetic Resonance Imaging in People with Multiple Sclerosis

MULTIDISCIPLINARY CARE

- (MDC01) Social Assistance Intervention in Multiple Sclerosis
- (MDC02) Is a 2-Week Intensive Day Program an Effective Approach to Provide Outpatient Services for People with Multiple Sclerosis?
- (MDC03) The Waiting Room: A Successful Experience in the Multiple Sclerosis Care and Treatment Center (CATEM)
- (MDC05) Implementation of a Pharmacist-Led Immunization Program in a Center for Comprehensive Multiple Sclerosis (MS) Care
- (MDC06) The African American Experience and Multiple Sclerosis
- (MDC07) The Impact of the Nurse Practitioner Model of Care within Multiple Sclerosis
- (MDC08) Late-Onset Multiple Sclerosis: Comorbidity and Disease Progression
- (MDC09) Comparing Patient Perceptions on Multiple Sclerosis Management and Care: A Subanalysis of Geographic Differences

METHODS OF CARE

- (MOC01) Understanding the Health Care Provider–Patient Relationship in Treating Multiple Sclerosis
- (MOC02) Cancelling Clinic Appointments: What Factors Are Associated with Higher Rates of Cancellations in Patients with Multiple Sclerosis?
- (MOC03) From Therapy Enrollment to First Dose: A Quality Improvement Initiative for Multiple Sclerosis Care
- (MOC04) Nurse Telephone Encounters in a Multiple Sclerosis Clinic in 2020
- (MOC05) Pioneering Multiple Sclerosis Center Program with MSHA Certification to Improve Patient Care and Experience
- (MOC06) Conceptualizing Access Through the Perspectives of Canadians with Multiple Sclerosis
- (MOC07) Access to Health Care for Canadians with Multiple Sclerosis: Prioritizing Concerns
- (MOC08) Use of a Clinical Decision Support Tool to Support Monitoring and Care of Patients with Multiple Sclerosis Receiving Disease-Modifying Therapy
- (MOC09) Multiple Sclerosis Disease Impact Monitoring: Longitudinal Exploration of the Relationship of Ocular Coherence Tomography to Computerized Cognitive Testing
- (MOC11) Improving Understanding of Clinical Phenotype for Patients with Multiple Sclerosis: Design and Implementation of Smarttools in Electronic Health Record Systems
- (MOC12) Changing Language to Acknowledging Patients' Perceptions of Treatment in Multiple Sclerosis Care

NEUROIMMUNOLOGY AND DISEASE MODELS

- (NDM01) Efficacy of the Influenza Vaccine in Multiple Sclerosis Patients: A Systematic Review and Meta-analysis

NONIMAGING BIOMARKERS

- (NIB01) Higher Sensitivity of Quantitative Reverse Transcriptase–Polymerase Chain Reaction Compared with Flow Cytometry for Quantification of B Cells After Anti-CD20 Monoclonal Antibody Therapy
- (NIB02) Quantification of Smooth Pursuit Dysfunction in Multiple Sclerosis

NEUROPHYSIOLOGY, NEUROPSYCHOLOGY, AND NEUROPSYCHIATRY

- (NNN02) Education as a Moderating Variable in the Relationship Between Patient Self-Perception of Cognitive Impairment and Symbol Digit Modalities Test Performance in Multiple Sclerosis

- (NNN03) Preliminary Cognitive Outcomes Following Mesenchymal Stem Cell Therapy in Multiple Sclerosis
- (NNN04) Relationship Between Expanded Disability Status Scale Scoring and Attention Performance in People with Multiple Sclerosis
- (NNN05) Objective Measurement of Cognitive Impairment in Multiple Sclerosis Patients Using Novel Computerized Testing

PROGRAMS

- (PGM01) The Use of a Multiple Sclerosis Documentary Film Screening Program as an Educational Intervention to Increase Knowledge and Awareness About MS and Support Resources
- (PGM03) Dance for MS: A Structured Dance Program Targeted for Multiple Sclerosis Patients
- (PGM04) Development of an Effective Age-Span Program for Women with Multiple Sclerosis: A Patient Perspective
- (PGM05) National MS Society Pathways to a Cure: An In-Person Educational Program for People Affected by Multiple Sclerosis
- (PGM06) How Well Do Junior Neurology Residents Recognize Multiple Sclerosis? Analysis of the "Close the Loop" Clinical Acumen Project
- (PGM07) Time to Adult: Transitioning from Pediatric to Adult Health Care in Demyelinating Disorders
- (PGM08) Successful Pilot of MS VA-ECHO Tele-education Program for Rural Providers
- (PGM09) Current Topics in MS Webinar Series: A Professional Education Collaboration Between the National MS Society, Consortium of Multiple Sclerosis Centers, and the VA MS Centers of Excellence
- (PGM10) Multiple Sclerosis Nurse Fellowship Pilot: A 6-Month Immersion
- (PGM11) Collaborative Working Between Multiple Sclerosis (MS) Nurses and a Pharmaceutical Company: An Educational Project from the Consortium of Multiple Sclerosis Centers (CMSC) Conference, Seattle, 2019

PSYCHOSOCIAL FACTORS

- (PSF01) Differences in Depressive Symptomology Between Females and Males with Relapsing-Remitting Multiple Sclerosis
- (PSF03) Me Too MS: Physical, Sexual, and Other Forms of Violence Experience in Women with Multiple Sclerosis
- (PSF04) Predictors for Self-Efficacy for People Living with Multiple Sclerosis
- (PSF05) The Association Between Health Literacy, Health Outcomes, and Medication Adherence in Patients with Multiple Sclerosis
- (PSF06) It Takes a Village: The Veterans Health Administration (VHA) MS Centers of Excellence and National Multiple Sclerosis Society Partnership for Facilitating Communication, Collaboration, and Coordination of Services for Veterans with Multiple Sclerosis
- (PSF07) Discussing Multiple Sclerosis (MS) Progression with Patients: Experiences of UK Health Care Professionals from the Spectrum Project
- (PSF08) Development and Implementation of a Patient Education and Cognitive Wellness Program for Veterans with Multiple Sclerosis
- (PSF09) The Effects of Customized Psychoeducation-Based Neurocounseling Interventions on the Coping Flexibility of African American Women with Multiple Sclerosis
- (PSF10) The Conformity of Masculine Norms and the Effects on Coping, Health Behaviors, and Quality of Life in Men with Multiple Sclerosis
- (PSF11) Understanding the Lived Experience of Health Through the Exploration of Well-Being of Women with Multiple Sclerosis: Preliminary Findings

QUALITY OF LIFE AND OUTCOMES

- (QOL01) Alemtuzumab Effects on Urogenital Function: Results Pooled From the CARE-MS 9-Year Functional Assessment of Multiple Sclerosis Quality-of-Life Survey
- (QOL02) The Impact of Relapses on Quality of Life in Patients with Neuromyelitis Optica Spectrum Disorder: Data from the Phase 3 PREVENT Study
- (QOL04) Understand Common Multiple Sclerosis Symptoms Experienced Among MS Patients Participating in an Online MS Community
- (QOL05) Uncovering the Needs and Gaps in Care Among Multiple Sclerosis Patients Participating in an Online MS Community
- (QOL06) System-Level Variation in All-Cause Hospitalizations in Multiple Sclerosis: Year-1 Results of the Multiple Sclerosis Continuous Quality Improvement (MS-CQI) Research Collaborative
- (QOL07) Relapse Rate Is Influenced by System-Level Variation: Year-2 Results of the Multiple Sclerosis Continuous Quality Improvement (MS-CQI) Research Collaborative
- (QOL09) Living with Secondary Progressive Multiple Sclerosis: Results from an MS Coalition Survey
- (QOL10) Corticosteroid Pulse Therapy–Hyperglycemia: Protocol for Capillary Blood Glucose Control

- (QOL11) Natalizumab Is Associated with Improvement in Cognitive Processing Speed and Health-Related Quality of Life: STRIVE 4-Year Results
- (QOL12) Multiple Sclerosis Patient Perspectives: Disease Education and Communication Needs
- (QOL13) Multiple Sclerosis Patient Perspectives: Impact on Employment
- (QOL14) Determining the Relationship of Demographic and Clinical Variables with Fatigue in Multiple Sclerosis, Using the 5-Item Modified Fatigue Impact Scale (MFIS-5)
- (QOL15) Multiple Sclerosis Management and Expanded Disability Status Scale: A Great Start, but a Reason for Change Was Never So Apparent and Needed
- (QOL16) A Coach-Supported Standardized Approach for Quality Improvement (QI) in the Multiple Sclerosis Continuous Quality Improvement (MS-CQI) Research Collaborative
- (QOL18) Does Participation in a 6-Week Mindfulness Course Improve Mood and Overall Emotional Wellness for People Living with Multiple Sclerosis?
- (QOL19) The Mediterranean Diet and Fatigue, Depression, and Emotional Well-being in Multiple Sclerosis: A Study in Patient-Reported Outcomes

REHABILITATION

- (REH01) Assistive Technology for Progressive Deficits in Communication and Access in People with Advanced Multiple Sclerosis: Case Studies in Iterative Design
- (REH02) Predictors of Improvement in Respiratory Function Following Resistive Inspiratory Muscle Training in Advanced Multiple Sclerosis
- (REH03) Rehabilitation of Dysarthria in Multiple Sclerosis
- (REH04) Daily Occupational Performance in Multiple Sclerosis
- (REH05) Assessing the Benefits of Using Telehealth in Conjunction with a Fitbit to Improve Walking in Veterans with Multiple Sclerosis
- (REH06) Feasibility of Telehealth Rehabilitation for Veterans with Progressive Neuromuscular Disease
- (REH07) Systematic Review on Exercise Training as a Neuroplasticity-Inducing Behavior in Multiple Sclerosis
- (REH08) Aerobic Reserve in People with Multiple Sclerosis: Measurement and Correlates
- (REH09) The Impact of Vascular Comorbidities on Perceived Functional Impact in Persons with Multiple Sclerosis
- (REH10) Is Treadmill Walking Analogous to Overground Walking in Persons with Multiple Sclerosis?
- (REH11) Telerehabilitation Compared to Outpatient Rehabilitation for Patients with Multiple Sclerosis and Mobility Disorders
- (REH12) Management of Low Back Pain for Individuals with Multiple Sclerosis: A Case Series
- (REH13) Predicting Fall Risk in Persons with Multiple Sclerosis Utilizing the 12-Item Multiple Sclerosis Walking Scale
- (REH14) The Impact of Lower Limb Strength on Walking in Persons with Multiple Sclerosis: A Preliminary Analysis
- (REH15) The Effects of Intermittent Versus Continuous Walking on Distance to Fatigue in Persons with Multiple Sclerosis
- (REH16) The Effects of Cooling Vests on Gait Fatigability in Persons with Multiple Sclerosis
- (REH17) A Combination of Core Exercise and Balance-Based Torso Weighting for Women with Multiple Sclerosis
- (REH21) Impact of Restless Legs Syndrome Severity on Cognitive Function in Adults with Multiple Sclerosis and Restless Legs Syndrome
- (REH22) Utilization of Therapy Services Among Patients with Multiple Sclerosis
- (REH23) Proximal Movement Compensations Are Related to Muscle Function and Walking Capacity in People with Multiple Sclerosis
- (REH24) Effects of a Weight-Based Training Program on Bone Density, Cognition, and Quality of Life of Multiple Sclerosis Patients
- (REH25) Orchestrating a New Path for Multiple Sclerosis Rehabilitation: Empowering Patients Through Both Physical and Music Therapies
- (REH26) Improving Quality of Life Using an End-Effector Robotic Rehabilitation Approach in Progressive Multiple Sclerosis
- (REH27) Strategies to Foster Buy-in for Physical and Occupational Therapists: Engagement Across Multiple Sites and States for a Study on Tele-Exercise and Multiple Sclerosis (TEAMS)
- (REH28) Correlates of Change in First Trial Exposures Across 2 Days of Protective Steps Among Those with Multiple Sclerosis
- (REH29) Lifestyle Redesign for Multiple Sclerosis: A Case Series of Female Hispanic Patients
- (REH30) Cognitive Processing Speed as a Predictor of Motor Skill Learning in Healthy Adults and Persons with Multiple Sclerosis

RELAPSE THERAPY

- (RTH01) Safety of Satralizumab Based on Pooled Data from Phase 3 Studies in Patients with Neuromyelitis Optica Spectrum Disorder (NMOSD)
- (RTH02) Adolescents with Neuromyelitis Optica Spectrum Disorder Achieved Similar Exposures and Favorable Safety Profile When Treated with the Adult Satralizumab Dosing Regimen
- (RTH03) Efficacy and Safety Outcomes from a Prospective Observational Registry of Repository Corticotropin Injection for Relapse of Multiple Sclerosis
- (RTH04) A Prospective Clinical Trial Utilizing an iOS Mobile Application and Apple Watch to Manage Multiple Sclerosis and Predict Disease Relapses
- (RTH05) Characterization of the Pharmacokinetics and Pharmacodynamics of Satralizumab, a Recycling Antibody, to Support Once-Every-4-Weeks Dosing in Patients with Neuromyelitis Optica Spectrum Disorder

SELF-CARE

- (SEL01) Understanding Gaps in Knowledge Surrounding Flu Shots and Immunizations as They Relate to Multiple Sclerosis
- (SEL02) Experimental Project Fashion Design and Social Inclusion: Multiple Women
- (SEL03) An Assessment of the Feasibility of a Dyadic Physical Activity Intervention for Persons with Advanced Multiple Sclerosis and Their Family Caregivers: Work in Progress

SYMPTOM MANAGEMENT

- (SXM01) Effect of Nabiximols on Spasticity and Muscle Strength in Patients with Multiple Sclerosis Across 3 Randomized Controlled Trials
- (SXM02) MS Action Plan May Be Effective Tool Helping Patients with Acute Change in Multiple Sclerosis Symptoms
- (SXM03) Intrathecal Baclofen Therapy in Ambulatory and Nonambulatory Multiple Sclerosis Patients: A Single-Center Experience
- (SXM05) Virtual Delivery of Mindfulness-Based Art Therapy (MBAT) to Improve Symptoms Among Adults with Multiple Sclerosis (MS)
- (SXM06) A Pilot Study of Mirabegron (Myrbetriq) and Behavioral Modification Including Pelvic Floor Exercise for Overactive Bladder in Multiple Sclerosis
- (SXM07) The Clinical Spectrum of Myelin Oligodendrocyte Glycoprotein (MOG) Antibody-Associated Demyelinating Disorders: Three Case Reports

WHITAKER RESEARCH TRACK

- (WHI01) Characterizing the Acute Exercise Response in Nonambulatory People with Progressive Multiple Sclerosis
- (WHI03) Youth with Multiple Sclerosis Have Low Fitness Levels
- (WHI04) Dietary Patterns and Health-Related Quality of Life of Individuals with Multiple Sclerosis

LATE BREAKING

- Real-World Evidence Assessment of Betaseron (interferon beta-1b) Adherence Following the Introduction of the BETACONNECT Autoinjector
- A Unique Case of a Patient with Tuberous Sclerosis and Recent Diagnosis of Neuromyelitis Optica
- Shorter Infusion Time of Ocrelizumab: Primary Results from the ENSEMBLE PLUS Study in Patients with Relapsing-Remitting Multiple Sclerosis
- Effect of Ofatumumab on Serum Immunoglobulin Levels and Infection Risk in Relapsing Multiple Sclerosis Patients from the Phase 3 ASCLEPIOS I and II Trials
- Ofatumumab Versus Teriflunomide in Relapsing Multiple Sclerosis: Analysis of No Evidence of Disease Activity (NEDA-3) from the ASCLEPIOS I and II Trials
- Real-World Findings of Usability and Usefulness of Multiple Sclerosis Progression Discussion Tool: A Physician-Completed Digital Tool to Evaluate Early Signs of Disease Progression
- Efficacy and Safety of the Bruton's Tyrosine Kinase Inhibitor (BTKI) Evobrutinib in Relapsing Multiple Sclerosis over 108 Weeks: Open-Label Extension to a Phase 2 Study
- Natalizumab-Treated Patients with Relapsing-Remitting Multiple Sclerosis Report Better "Feel-Good" Outcomes on Key Physical, Emotional, and Cognitive Domains Compared to Other Disease-Modifying Therapies

PLATFORMS

DISEASE ASSESSMENT AND MANAGEMENT

(DAM01) International Registry Tracking Pregnancy Outcomes in Women Treated with Dimethyl Fumarate

Maria Houtchens,¹ Kerstin Hellwig,² David Rog,³ Christopher McGuigan,⁴ Denise R. Bruen,⁵ Kun Chen,⁶ Xiaomei Peng,⁶ Cynthia C. Jones⁶

¹Brigham and Women's Hospital, Boston, MA; ²University of Bochum, Neurological Clinic, Bochum, Germany; ³Manchester Centre for Clinical Neurosciences, Salford, United Kingdom; ⁴Department of Neurology, St. Vincent's University Hospital, Dublin, Ireland; ⁵Department of Neuroscience, University of Virginia Health, Charlottesville, VA; ⁶Biogen, Cambridge, MA

Background: To date, delayed-release dimethyl fumarate (DMF) exposure during pregnancy has not shown any safety signals in clinical trial and postmarketing data, however the DMF label recommends use during pregnancy only if potential benefit justifies the potential risk to the fetus. In the general population, 62% of pregnancies end in live birth, 22% end in induced abortion, and 16% end in fetal loss. Similar rates have been observed in patients with multiple sclerosis (MS). **Objectives:** An international registry (trial registration: NCT01911767) was started to prospectively evaluate pregnancy outcomes in women with MS exposed to DMF since 1 day before the first day of their last menstrual period before conception or during pregnancy; results are reported for the United States as well as the overall population. **Methods:** In this ongoing registry, data were collected at enrollment, 6-7 months of gestation, 4 weeks after estimated delivery date, and at 4, 12, and 52 weeks after birth. Infant and maternal outcomes included ectopic and molar pregnancies, birth defects, congenital anomalies or infant death occurring at ≤ 52 weeks of age, and maternal death at ≤ 12 weeks postdelivery. Potential birth defects were adjudicated by an external expert. Gestational size was classified as small ($<10^{\text{th}}$ percentile), appropriate ($10^{\text{th}}-90^{\text{th}}$), or large ($>90^{\text{th}}$) based on standardized growth charts. **Results:** As of April 2019, 263 patients were enrolled; 57 in the United States. Median gestational week at first DMF exposure was 1 (range, 1-13); median fetal DMF exposure duration was 5 (range, 0.1-40) weeks. Of the 214 pregnancy outcomes reported to date, 197 (92%) were live births and 17 (8%) fetal losses. In the United States, 38 pregnancy outcomes have been reported to date, 34 (89%) live births and 4 (11%) fetal losses. Of infants with known gestational age ($n = 194$), 176 (89%) births were full-term and 18 (9%) premature (<37 weeks). In the United States, 30 (97%) births were full-term and 1 (3%) premature. There were 16 spontaneous abortions (4 in US; 1 ectopic pregnancy outside of US), and 1 fetal death at ≥ 28 weeks' gestation. No perinatal, infant, or maternal deaths were reported. Infants (163 with gestational size data) were classified as small 18 (11%), appropriate 134 (82%), and large 11 (7%). Seven (4%) infants had confirmed birth defects. **Conclusions:** The adverse pregnancy outcome frequencies from the interim analysis did not exceed those observed in the MS and general populations. No safety signal has been observed to date.

Supported by: Biogen

Disclosure: *Maria Houtchens:* Nothing to disclose. *Kerstin Hellwig:* Bayer, Biogen, Genzyme, Sanofi, Teva, Roche, Novartis, Merck (consulting fee); Bayer, Biogen, Merck, Novartis, Sanofi Genzyme, Teva (contracted research, speakers' bureau). *David Rog:* Actelion, Biogen, Merck Serono, Novartis, Sanofi, Teva Neuroscience, TG Therapeutics (contracted research); Biogen, MedDay, Merck Serono, Novartis, Roche, Sanofi, Teva Neuroscience (consulting fee); Biogen, Merck Serono, Novartis, Roche, Sanofi, Teva Neuroscience (speakers' bureau). *Christopher McGuigan:* Actelion, Biogen, Merck, Novartis, Roche, Sandoz, Sanofi Genzyme (consulting fee, contracted research, speakers' bureau). *Denise R. Bruen:* Biogen, Genentech, Sanofi Genzyme, EMD Serono, Novartis, Celgene (consulting fee). *Kun Chen, Xiaomei Peng, Cynthia C. Jones:* Biogen (ownership interest, salary).

Keywords: Disease-modifying treatments in MS, Pregnancy

(DAM02) Menarche and Relapses in Girls with Pediatric Multiple Sclerosis

Kristen M. Krysko,¹ Michael Waltz,² Tanuja Chinnis,³ Bianca Weinstock-Guttman,⁴ Greg Aaen,⁵ Leslie Benson,⁶ Mark Gorman,⁶ Yolanda Harris,⁷ Lauren B. Krupp,⁸ Timothy Lotze,⁹ Soe Mar,¹⁰ Manikum Moodley,¹¹ Jayne Ness,¹² Mary Rensel,¹³ Moses Rodriguez,¹⁴ John Rose,¹⁵ Alice Rutatangwa,¹ Teri Schreiner,¹⁶ Emmanuelle Waubant,¹ T. Charles Casper,² Jennifer S. Graves^{17,18}

¹Neurology, University of California San Francisco, San Francisco, CA; ²Pediatrics, University of Utah, Salt Lake City, UT; ³Pediatric Neurology, Massachusetts General Hospital, Boston, MA; ⁴Neurology, State University of New York at Buffalo, Buffalo, NY; ⁵Pediatrics, Loma Linda University, San Bernardino, CA; ⁶Neurology, Boston Children's

Hospital, Boston, MA; ⁷Nursing, University of Alabama at Birmingham, Birmingham, AL; ⁸Neurology, NYU Langone Medical Center, New York, NY; ⁹Neurology, Texas Children's Hospital, Houston, TX; ¹⁰Neurology, Washington University in Saint Louis, St. Louis, MO; ¹¹Pediatrics and Neurology, Dell Children's Hospital, University of Texas, Austin, TX; ¹²Pediatrics, University of Alabama at Birmingham, Birmingham, AL; ¹³Neurology, Cleveland Clinic, Cleveland, OH; ¹⁴Neurology, Mayo Clinic, Rochester, MN; ¹⁵Neurology, University of Utah, Salt Lake City, UT; ¹⁶Neurology and Pediatrics, University of Colorado, Aurora, CO; ¹⁷Neurosciences, University of California San Diego, San Diego, CA; ¹⁸Neurology, University of California San Diego, La Jolla, CA

Background: Sex steroid hormones have a clinical impact on the immune system. Puberty may trigger multiple sclerosis (MS) disease activity, with mean age of pediatric MS onset occurring near age 13 years. **Objectives:** To evaluate the association between menarche and disease course in pediatric MS through comparison of relapse rates across the premenarche, perimenarche, and postmenarche periods. **Methods:** This is a retrospective analysis of a prospectively followed cohort of girls meeting MS criteria within the US Network of Pediatric MS Centers database. Only individuals with known menarche dates were included in the analysis. Relapses were collected prospectively. Both negative binomial and repeated Cox regression models were used to assess the association of pubertal development stage with relapse rate, adjusted for tier of disease-modifying therapy and body mass index. **Results:** Of the 503 girls included, onset was during premenarche in 53, perimenarche in 84 (within ± 1 year of menarche), and postmenarche in 366. The median time of MS onset was 2.5 years after menarche. In adjusted negative binomial analysis, annual relapse rate during the premenarche period was 0.43, perimenarche period was 0.50, and postmenarche period was 0.36 ($P = .16$). In adjusted repeated-events Cox regression analysis, there was increased hazard to relapse with stage of development from premenarche through menarche [premenarche HR 0.60 [95% CI, 0.45-0.79] and perimenarche HR 0.79 [95% CI, 0.62-1.02] compared to the reference of postmenarche, $P = .0006$]. **Conclusions:** Before menarche girls have lower relapse rates. Onset of puberty may be a time of increase in disease activity and may require consideration of a change in therapeutic approach.

Supported by: None

Disclosure: *Kristen M. Krysko:* Biogen (fellowship funding). *Michael Waltz, Soe Mar, Manikum Moodley, Moses Rodriguez, T. Charles Casper:* Nothing to disclose. *Tanuja Chinnis:* Biogen, Novartis, Octave, Serono, Mallinckrodt, Verily (research support); Biogen, Novartis, Sanofi Genzyme, Bayer, Celgene, Genentech (consulting fee); Sanofi Genzyme, Novartis (contracted research). *Bianca Weinstock-Guttman:* Biogen, Novartis, Genentech, EMD Serono (research support); Biogen, Novartis, Genentech, Celgene, Mallinckrodt, EMD Serono (consulting fee). *Greg Aaen, Leslie Benson:* Biogen (contracted research). *Mark Gorman:* Novartis, Biogen (contracted research); Pfizer (research support). *Yolanda Harris:* Celgene (consulting fee). *Lauren B. Krupp:* Biogen, Novartis (research support); Biogen, Novartis, Red Hill Pharmaceutical, Genzyme, Gerson Lehman, Sanofi, Celgene (consulting fee); biotechnology and pharmaceutical companies for the fatigue severity scale (royalty). *Timothy Lotze:* Biogen (consulting fee). *Jayne Ness:* Novartis, Chugai, Roche (contracted research). *Mary Rensel:* EMD Serono, Biogen, Teva, Genzyme, Novartis (consulting fee); Genzyme (educational grants); MedImmune, Novartis, Biogen (MS Paths), Genentech (research support); Novartis, Genzyme, Biogen, Multiple Sclerosis Association of America (speakers' bureau). *John Rose:* Biogen, Teva Neuroscience, Viela Bio, Accredo, Genentech, Novartis (unrestricted educational grants); Teva Neuroscience, Biogen (research support). *Alice Rutatangwa:* Biogen (fellowship grant). *Teri Schreiner:* Biogen, Adamas, MSDx (contracted research). *Emmanuelle Waubant:* Genentech, Biogen (contracted research). *Jennifer S. Graves:* Biogen, Octave, Genentech (research support); Novartis, Genentech, Alexion, Celgene (consulting fee).

Keywords: Epidemiology of MS, Hormonal factors in MS, Natural history of MS

(DAM03) Complexity of Aging with Multiple Sclerosis: Graceful Concessions or Kicking and Screaming?

Emma V. Richardson, Robert W. Moll

Department of Physical Therapy, University of Alabama at Birmingham, Birmingham, AL

Background: Over the past 3 decades there have been significant advances in the development of pharmaceutical and rehabilitative treatments for persons with multiple sclerosis (MS), such that life expectancy is continuing to increase. As a result of these advancements, there is a "graying" phenomenon within the global MS population and a demographic shift of the aging landscape. While these advancements are exciting, there also concerns and unknowns regarding what it is like to age with MS. **Objectives:** The objectives of this research were to explore the experiences of aging in conjunction with having MS, and the different ways persons older than 60 with MS interpreted this phenomenon. **Methods:** Semistructured interviews with 40 persons with MS older

than 60 years were conducted. Thereafter data were subject to a pluralistic analysis approach utilizing phenomenological and narrative traditions. **Results:** This research highlighted the complexity of aging with MS and the various ways persons older than 60 with MS experience and interpret this phenomenon. Most participants experienced a continued progression of physical and cognitive deficits, however aging narratives and what is culturally expected as a person older than 60 allowed for positive interpretations of this stage of life. For example, participants diagnosed in early adulthood stated having health problems was "normal" over 60, resulting in a sense of belonging that they had not experienced since being diagnosed. Some participants also perceived they were aging more successfully than peers without MS as they had engaged in health-enhancing behaviors from an earlier age, while others believed they had "aged out" of MS and were experiencing a peak of health and wellness. Participants who were diagnosed in middle age, however, noted a sharp progression of age-related and MS symptoms, but stated "everyone has something" and perceived the diagnosis of MS was less impactful in older age with regards to what is expected at this life stage. Concerns remained, however, regarding whether a new physical or cognitive experience was aging or MS, what the future holds regarding losing independence, losing spouses and caregivers, and growing "too old" such that quality of life is completely diminished. **Conclusions:** This qualitative research has highlighted the complexity of aging with MS. All participants noted a continued progression of health, physical function, and cognitive ability but stated that they felt a sense of normality and were living a life that culturally aligned with aging narratives. Fears about the future do remain as participants were concerned about living too long for being able to thrive. More research must be done that focuses on maintaining quality of life among older person with MS as quantity of life continues to increase.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Aging and MS, Management of activities of daily living in MS

(DAM04) Serum Glial Fibrillary Acidic Protein Is Elevated in a Subset of Neuromyelitis Optica Patients and Associated with Increased Risk of Attacks

Hans-Peter Hartung,¹ Orhan Aktas,² Michael A. Smith,³ William Rees,³ Kazuo Fujihara,⁴ Friedemann Paul,⁵ Romain Marignier,⁶ Jeffrey L. Bennett,⁷ Ho Jin Kim,⁸ Brian Weinschenker,⁹ Sean J. Pittock,¹⁰ Dean Wingerchuk,¹¹ Gary Cutter,¹² Ari Green,¹³ Maureen A. Mealy,³ Jorn Drappa,³ Gerard Barron,³ Soraya Madani,³ Liangwei Wang,³ Dewei She,³ Daniel Cimbora,³ John N. Ratchford,³ Eliezer Katz,³ Bruce A.C. Cree¹⁴

¹Department of Neurology, Medical Faculty, Heinrich Heine University, Dusseldorf, Germany; ²Medical Faculty, Heinrich-Heine-University, Dusseldorf, Germany; ³Viola Bio, Gaithersburg, MD; ⁴Department of Multiple Sclerosis Therapeutics, Southern Tohoku Research Institute for Neuroscience (STRINS), Koriyama, Japan; ⁵Experimental and Clinical Research Center, Max Delbrueck Center for Molecular Medicine and Charite - Universitätsmedizin Berlin, Berlin, Germany; ⁶Lyon University Hospital, Lyon, France; ⁷School of Medicine, University of Colorado, Aurora, CO; ⁸Research Institute and Hospital, National Cancer Center, Goyang, Korea, Republic of (South); ⁹Neurology, Mayo Clinic, Rochester, MN; ¹⁰Mayo Clinic, Rochester, MN; ¹¹Neurology, Mayo Clinic Arizona, Scottsdale, AZ; ¹²Biostatistics, University of Alabama at Birmingham, Birmingham, AL; ¹³Department of Neurology and Department of Ophthalmology, UCSF Weill Institute for Neurosciences, San Francisco, CA; ¹⁴Department of Neurology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA

Background: In patients with neuromyelitis optica spectrum disorder (NMOSD), pathogenic anti-aquaporin-4 antibodies cause astroglial injury resulting in increased levels of cerebrospinal fluid glial fibrillary acidic protein (GFAP). N-MOMentum is a randomized, placebo-controlled, double-masked trial of inebilizumab, an anti-CD19 monoclonal B-cell depleting antibody, in patients with NMOSD. **Objectives:** To investigate the relationship between prospectively sampled serum GFAP (sGFAP) levels and disease activity in N-MOMentum trial participants. **Methods:** sGFAP (Quanterix Simoa GFAP assay) was measured in 1260 serial and attack-related samples collected from N-MOMentum participants (n = 220) and in healthy controls (n = 25); the relationship between sGFAP levels and NMOSD attacks was assessed in N-MOMentum participants. **Results:** Median (interquartile range [IQR]) sGFAP levels were elevated in patients with NMOSD compared with controls (128.3 [92.5, 182.1] vs 73.3 [52.1, 108.7] pg/mL). Elevated sGFAP, defined as ≥ 3 SDs above the control mean (≥ 171 pg/mL), was observed at baseline in 29% (61/215) of NMOSD study participants. Study participants with elevated baseline sGFAP levels were 2.9 times more likely to experience an adjudicated NMOSD attack than those with lower baseline sGFAP during the 28-week randomized controlled period (RCP: $P = .002$). During the RCP, sGFAP levels increased significantly within 1 week of an NMOSD attack in placebo-treated participants (median fold change [IQR]: 20.2 [4.4, 98.3]; n = 17 attacks; $P = .001$) but did not increase significantly

during attacks in inebilizumab-treated patients (1.1 [0.75, 24.6]; n = 20 attacks; $P > .05$). Of 143 participants who did not have an adjudicated NMOSD attack during the RCP, there were fewer inebilizumab- than placebo-treated participants with elevated sGFAP levels at the end of the RCP (16% [19/117] vs 35% [9/26]). Of 514 samples drawn from inebilizumab-treated patients during the RCP who did not have adjudicated attacks, 14 samples (2.7%) from 12 patients displayed ≥ 2 -fold increase in sGFAP from baseline vs 9 samples (of 116; 8%) from 6 placebo-treated patients (odds ratio: 3.0, $P = .023$). **Conclusions:** Study participants with NMOSD had increased sGFAP levels compared with controls. sGFAP may prove to be a useful biomarker of attack risk and disease activity and severity.

Supported by: None

Disclosure: Hans-Peter Hartung: Bayer HealthCare, Biogen, CSL Behring, GeNeuro, Genzyme, MedImmune, Merck Serono, Novartis, Opexa, Receptos, Roche, Sanofi, Teva, Viola Bio (consulting, speaking, serving on steering committees). Orhan Aktas: Almirall, MedImmune, Merck Serono, Roche (personal fees); Bayer HealthCare, Biogen, Genzyme, Novartis, Teva, Viola Bio (grants, personal fees); German Research Foundation (DFG), German Ministry of Education and Research (BMBF) (grants). Michael A. Smith, William Rees, Maureen A. Mealy, Jorn Drappa, Gerard Barron, Soraya Madani, Liangwei Wang, Dewei She, Daniel Cimbora, John N. Ratchford, Eliezer Katz: Viola Bio (salary). Kazuo Fujihara: Alexion, Bayer Schering, Biogen, Chugai, MedImmune, Merck Serono, Mitsubishi Tanabe Pharma, Nihon Pharmaceutical, Novartis, Ono, Viola Bio (scientific advisory boards); Asahi Kasei Medical, Astellas, Bayer Schering, Biogen, Daiichi Sankyo, Dainippon Sumitomo, Eisai, Mitsubishi Tanabe Pharma, Nihon Pharmaceutical, Novartis, Takeda (funding for travel, speaking honoraria); Asahi Kasei Medical, Bayer Schering, Biogen, Chemo-Sero-Therapeutic Research Institute, Chugai, Genzyme Japan, Mitsubishi Tanabe Pharma, Nihon Pharmaceutical, Ono, Teijin, Teva, Ministry of Education, Culture, Sports, Science and Technology of Japan, Ministry of Health, Welfare and Labor of Japan (contracted research). Friedemann Paul: Bayer, Biogen, Merck Serono, Novartis, Sanofi Genzyme, Teva (research support, speaking honoraria, travel reimbursement); Guthy Jackson Charitable Foundation (travel funding); Novartis (steering committee); German Research Council (DFG Exc 257), German Competence Network for Multiple Sclerosis (financial support). Romain Marignier: Biogen, Merck Serono, Novartis, Roche, Sanofi Genzyme, Teva, Viola Bio (funding for travel, honoraria); MedImmune, Viola Bio (scientific advisory board). Jeffrey L. Bennett: AbbVie, Chugai, Clene Nanomedicine, Equillium, Frequency Therapeutics, Genentech, Genzyme, Alexion (personal fees); Aquaporinab (patent issued); EMD Serono, Novartis (grants, personal fees). MedImmune/Viola Bio (consulting fee); Guthy-Jackson Charitable Foundation, Mallinckrodt, National Institutes of Health (grants). Ho Jin Kim: Eisai, HanAll BioPharma, MedImmune/Viola Bio, Merck Serono, Novartis (personal fees); Journal of Clinical Neurology and Multiple Sclerosis Journal (associate editor/co-editor); National Research Foundation of Korea (grants); Sanofi Genzyme, Teva-Handok (grants, personal fees). Brian Weinschenker: Alexion, MedImmune, Viola Bio (royalties, payments); Mitsubishi Tanabe Pharma (consulting fee); RSR Ltd, Oxford University, Hospices Civils de Lyon, MVZ Labor PD Dr Volkemann und Kollegen GbR (royalty). Sean J. Pittock: Alexion, Autoimmune Encephalitis Alliance, Grifols, Euroimmun, NIH RO1 NS065829-01, MedImmune/Viola Bio, Guthy-Jackson Charitable Foundation (consulting fees, research support); named inventor on filed patents that relate to functional AQP4/NMO-IgG assays and NMO-IgG as a cancer marker (receipt of intellectual property rights/patent holder). Dean Wingerchuk: Alexion Pharmaceuticals, Terumo BCT, Guthy-Jackson Charitable Foundation (contracted research); BrainStorm Therapeutics, Caladrius Biosciences, Celgene, MedImmune, Novartis, Ono Pharmaceutical (personal fees); Chugai Pharmaceutical (consulting fee); MedImmune, Viola Bio (clinical trial adjudication committee). Gary Cutter: AMO Pharma, BioLineRx, Horizon Pharmaceuticals, Merck, Merck/Pfizer, National Heart, Lung, and Blood Institute (protocol review committee); National Institute of Child Health and Human Development (OPRU oversight committee), Neurim, OPKO Biologics, Orphazyme, Reata Pharmaceuticals, Receptos/Celgene, Sanofi-Aventis, Teva Pharmaceuticals (personal fees for participation on data and safety monitoring boards); Atara Biotherapeutics, Axon, Argenix, Biogen, Biotherapeutics, Brainstorm Cell Therapeutics, Charleston Laboratories Inc, Click Therapeutics, Genentech, Genzyme, GW Pharma, Klein Bunden Inc, MedDay, MedImmune, Novartis, Roche, SciFluor, Somablation, Teva Pharmaceuticals, TG Therapeutics, UTHealth Houston (consulting fee); Pythagoras Inc, AL (president of organization); Teva Neuroscience (grant); University of Alabama at Birmingham (salary). Ari Green: Biomure, Inception Sciences, JAMA Neurology, MedImmune/Viola Bio, Mylan, Synthron, Trims Pharma (other financial relationships [for activities as expert witness, associate editor, advisory board steering committee participation and end point adjudication]); Conrad N. Hilton Foundation, Tom Sberak MS Hope Foundation (grants); Pipeline Therapeutics (personal fees, other financial relationships). Bruce A.C. Cree: Akili, Alexion,

Biogen, GeNeuro, Novartis, Sanofi Genzyme, TG Therapeutics (consulting fee).

Keywords: Non-imaging biomarkers

(DAM05) Machine Learning Algorithms Applied to Visual Metrics to Classify Demyelinating Disease Diagnosis in Children

Beyza Ciftci,¹ Can Kavaklioglu,² Lauren Erdman,¹ Anna Goldenberg,^{3,4,5,6} Tara Berenbaum,⁷ Tara Feltham,⁷ Fiona Costello,⁸ Jean Mah,⁹ Arun Reginald,¹⁰ Brenda Banwell,¹¹ Giulia Longoni,¹² E. Ann Yeh¹²

¹Department of Pediatrics, The Hospital for Sick Children, Toronto, ON, Canada;

²Department of Mechanical and Industrial Engineering, Ryerson University, Toronto, ON, Canada;

³Genetics and Genome Biology Lab, The Hospital for Sick Children, Toronto, ON, Canada;

⁴Department of Computer Science, University of Toronto, Toronto, ON, Canada;

⁵Vector Institute, Toronto, ON, Canada;

⁶CFAR, Toronto, ON, Canada;

⁷Department of Neurosciences and Mental Health, The Hospital for Sick Children, Toronto, ON, Canada;

⁸Departments of Clinical Neurosciences and Surgery, University of Calgary, Calgary, AB, Canada;

⁹Department of Pediatrics, University of Calgary, Calgary, AB, Canada;

¹⁰Ophthalmology and Vision Sciences, The Hospital for Sick Children, Toronto, ON, Canada;

¹¹Neurology, The Children's Hospital of Philadelphia, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA;

¹²Department of Paediatrics, Division of Neurology, The Hospital for Sick Children, Toronto, ON, Canada

Background: Predicting disease classification in youth with a first episode of demyelination is feasible in some but not all cases. Machine learning algorithms can provide highly accurate predictions of outcomes directly from images, removing a major bottleneck in the speed with which predictions can be made. **Objectives:** To use visual metrics to predict disease class in youth with demyelination. **Methods:** Standardized, prospectively collected clinical and visual data at disease onset from 224 pediatric subjects, classified using consensus definitions of demyelinating disorders and serum antibody testing for myelin oligodendrocyte glycoprotein (MOG) and aquaporin 4 (healthy control = 72, multiple sclerosis = 69, anti-MOG = 18, neuromyelitis optica spectrum disorder [NMOSD] = 10, monophasic acquired demyelinating syndromes [MonoADSs] = 55) were recruited through the Demyelinating Disorders Program at The Hospital for Sick Children (Toronto, Ontario) and University of Calgary (2010-2019). We used 9 supervised machine learning algorithms (random forest classifier, CLS bagging classifier, AdaBoost classifier, XGBoost classifier, CLS logistic regression, MLP classifier, CLS SVM, decision tree, and Kneighbors classifier). Input variables were optical coherence tomography (OCT) parameters including retinal nerve fiber layer (RNFL) and ganglion cell inner plexiform layer thickness (GC IPL), low-contrast visual acuity (LCVA), and color vision for each eye. Separate sets of analyses were run with balanced (n set arbitrarily at 44, oversampling and under-sampling used appropriately) and unbalanced data. **Results:** In the analysis of the balanced data set, the random forest classifier (accuracy [a] = 80%, recall [r] = 50%, and precision [p] = 99%), XGBoost classifier (a = 80%, r = 50%, p = 80%), and decision tree (a = 90%, r = 99%, p = 90%) algorithms yielded the highest accuracy, recall, and precision levels for each disease class using the combination of RNFL, GC IPL, LCVA, and color vision. Analysis of the unbalanced data set showed lower overall levels of predictive accuracy (60%-90%) for each class using the same algorithms. Use of the OCT data alone yielded lower predictive accuracy in both balanced and unbalanced analyses. **Conclusions:** Machine learning algorithms used on combined structural and functional visual metrics may classify youth with demyelinating disorders. Implementation of artificial neural networks using raw OCT data and images as input are underway.

Supported by: None

Disclosure: Beyza Ciftci, Can Kavaklioglu, Lauren Erdman, Anna Goldenberg, Tara Berenbaum, Tara Feltham, Fiona Costello, Jean Mah, Arun Reginald, Giulia Longoni, E. Ann Yeh: Nothing to disclose. Brenda Banwell: Biogen, Merck Serono, Novartis, Teva Neuroscience (consulting fee).

Keywords: Machine learning and MS

(DAM06) Updated Recommendations for a Standardized Magnetic Resonance Imaging Protocol for Multiple Sclerosis

Anthony Traboulsee,¹ David Li,¹ Brenda Banwell,² Frederik Barkhof,³ Kathleen Costello,⁴ Peter Damiri,⁵ Scott D. Newsome,⁶ Jiwon Oh,⁷ Friedemann Paul,⁸ Daniel S. Reich,⁹ Mitchell T. Wallin,¹⁰ Jerry S. Wolinsky,¹¹ June Halper,¹² Sarah A. Morrow,¹³ Wim Van Hecke,¹⁴ Laura Barlow,¹⁵ Jason Shewchuk,¹⁶ Russell Shinohara,¹⁷ Amy Verrinder,¹⁸ Micki Maes,¹⁸ Patrick Quarterman,¹⁹ Kim van de Ven,²⁰ Shivraman Giri,²¹ Lori Saslow,²² CMSC MRI Guidelines Working Group

¹The University of British Columbia, Vancouver, BC, Canada; ²Neurology, The Children's Hospital of Philadelphia, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ³Institutes of Neurology and Healthcare Engineering, University College London, London, United Kingdom; ⁴National MS Society, New York, NY; ⁵Multiple Sclerosis

Association of America, Cherry Hill, NJ; ⁶Neurology, Johns Hopkins University School of Medicine, Baltimore, MD; ⁷University of Toronto, Toronto, ON, Canada; ⁸Experimental and Clinical Research Center, Max Delbrueck Center for Molecular Medicine and Charité – Universitätsmedizin Berlin, Berlin, Germany; ⁹Translational Neuroradiology Unit, National Institute of Neurological Disorders and Stroke, Bethesda, MD; ¹⁰Washington, DC, VA Medical Center, MS Center of Excellence, Washington, DC; ¹¹University of Texas Health Science Center at Houston (UTHealth), Houston, TX; ¹²CEO-CMSC, Hackensack, NJ; ¹³Western University, London, ON, Canada; ¹⁴Cometrix, Leuven, Belgium; ¹⁵UBC MRI Research Centre, University of British Columbia, Vancouver, BC, Canada; ¹⁶Vancouver General Hospital, Vancouver, BC, Canada; ¹⁷University of Pennsylvania, Philadelphia, PA; ¹⁸Cortechs Labs, Inc, San Diego, CA; ¹⁹GE Healthcare, New York, NY; ²⁰Philips Healthcare, Eindhoven, Netherlands; ²¹Siemens Healthineers, Boston, MA; ²²Consortium of Multiple Sclerosis Centers, Hackensack, NJ

Background: Standardize magnetic resonance imaging (MRI) protocols are important for the diagnosis and monitoring of patients with multiple sclerosis (MS). The Consortium of Multiple Sclerosis Centers (CMSC) convened an international panel of MRI experts to review and update the current guidelines. **Objectives:** To update the standardized MRI protocol and clinical guidelines for diagnosis and follow-up of MS and develop strategies for them to be universally accepted as standard of care in MRI for MS. **Methods:** The CMSC convened an expert panel in October 2019 to update the standardized MRI protocol. Conference attendees included neurologists, radiologists, magnetic resonance technologists, and imaging scientists with expertise in MS. Representatives from CMSC, Magnetic Resonance Imaging in MS (MAGNIMS), North American Imaging in Multiple Sclerosis Cooperative, National MS Society, Multiple Sclerosis Association of America, MRI manufacturers, and commercial image analysis companies were present. Before the meeting, CMSC members were surveyed about standardized MRI protocol, gadolinium, diffusion-weighted imaging, and the central vein sign. **Results:** The use of subcallosal plane for consistent repositioning and requirement for thin (≤ 3 mm), contiguous slices were re-emphasized. Updated Standardized Brain MRI Protocol includes core sequences optimized for MS lesion detection, and the judicious use of gadolinium when indicated. A new category of optimum plus sequences allows for brain atrophy monitoring and identifying lesions with a central vein sign. Other updates include progressive multifocal encephalopathy surveillance and spinal cord sequences. The group worked to make CMSC and MAGNIMS MRI protocols similar so the updated protocol could ultimately be accepted by international consensus. **Conclusions:** The international expert group developed revised clinical MRI protocols with the vision and action plans for them to be universally useful and useable and become the standard of care for patients with MS.

Supported by: None

Disclosure: Anthony Traboulsee: Biogen, Chugai, Roche, Sanofi, Teva (consulting and/or speaking fees, and grant/research support). David Li, Brenda Banwell, Frederik Barkhof, Kathleen Costello, Peter Damiri, Scott D. Newsome, Jiwon Oh, Friedemann Paul, Daniel S. Reich, Mitchell T. Wallin, Jerry S. Wolinsky, June Halper, Sarah A. Morrow, Wim Van Hecke, Laura Barlow, Jason Shewchuk, Russell Shinohara, Amy Verrinder, Micki Maes, Patrick Quarterman, Kim van de Ven, Shivraman Giri, Lori Saslow: Nothing to disclose.

Keywords: Disease-modifying treatments in MS, Imaging and MS

DISEASE-MODIFYING THERAPY

(DMT01) Comparative Effectiveness of Switching from Natalizumab to a Moderate- Versus High-Efficacy Disease-Modifying Therapy in Clinical Practice

Carrie M. Hersh,¹ Haleigh Harris,² Devon Conway,³ Le H. Hu¹

¹Lou Ruvo Center for Brain Health, Cleveland Clinic, Las Vegas, NV; ²Las Vegas, NV;

³Mellen Center for Multiple Sclerosis Treatment and Research, Cleveland, OH

Background: Natalizumab (NTZ) is a highly effective disease-modifying therapy (DMT) for relapsing multiple sclerosis (MS). Long-term use of NTZ is limited by potential safety risks that can be reduced by switching to an alternative therapy. However, NTZ discontinuation may trigger rebound disease, resulting in disability. Our previous study showed patients switching to moderate- (Mod) DMT vs high-efficacy therapy (HET) were at higher risk of early magnetic resonance imaging (MRI) disease activity by 6 months. **Objectives:** To assess the comparative effectiveness of switching from NTZ to a Mod DMT vs HET in patients with MS over 24 months. **Methods:** Patients discontinuing NTZ at 2 MS centers (n = 556) who switched to Mod DMT (n = 270) vs HET (n = 130) were assessed using propensity score (PS) weighting. PS model covariates included demographics and baseline clinical and radiographic disease characteristics. Outcomes included annualized relapse rate and proportions with new T2 and/or gadolinium-enhancing (GdE) lesions, absence of disease activity (a composite measure of no relapses and/or MRI activ-

ity), time-to-first relapse and GdE lesion, and 20% worsening of the Timed 25-Foot Walk (T25FW) and 9-Hole Peg Test (9-HPT). All outcomes were reported as Mod DMT vs HET. **Results:** In our cohort, 48.6% switched to Mod DMT (dimethyl fumarate, n = 130; fingolimod, n = 140) vs 23.4% who switched to HET (acrelizumab, n = 106; rituximab, n = 17; alemtuzumab, n = 7). Reasons for NTZ discontinuation included PML risk (54.9%), breakthrough disease (15.3%), and adverse effects (17.3%). PS weighting demonstrated excellent covariate balance. After PS adjustment, there were no differences in annualized relapse rate (rate ratio = 1.44 [95% CI, 0.69-1.59], P = .334) or time-to-first relapse (HR = 2.12 [95% CI, 0.87-5.17], P = .090) by 24-month follow-up. However, patients switching to Mod DMT had higher proportions with new T2 lesions (OR = 2.15 [95% CI, 1.18-3.01], P = .011), new GdE lesions (OR = 1.99 [95% CI, 1.12-2.73], P = .022), and 20% worsening of the T25FW (OR = 1.83 [95% CI, 1.06-3.02], P = .043) and 9-HPT (OR = 1.81 [95% CI, 1.05-3.56], P = .044), and lower proportion with absence of disease activity (OR = 0.41 [95% CI, 0.21-0.71], P = .004). Switchers to Mod DMT were also at higher risk of earlier time-to-first GdE lesion (HR = 6.67 [95% CI, 2.06-9.16], P = .002). **Conclusions:** By 24 months, NTZ switchers to Mod DMT vs HET had lower cumulative probability of no disease activity by 24 months and were at higher risk of disability accumulation.

Supported by: None

Disclosure: *Carrie M. Hersh:* Biogen, Genentech (consulting fee, contracted research); EMD Serono (consulting fee); Genzyme, Novartis (consulting fee, speakers' bureau); PCORI (contracted research). *Haleigh Harris:* Nothing to disclose. *Devon Conway:* National MS Society (contracted research); Novartis (consulting fee, contracted research); Tanabe Laboratories (consulting fee). *Le H. Hua:* Biogen, Celgene, EMD Serono, Genentech, Novartis (consulting fee); Genzyme (consulting fee, speakers' bureau).

Keywords: Comparative effectiveness, Disease-modifying treatments in MS

(DMT02) Yearly Efficacy and Safety Outcomes Over 4 Years After Last Alemtuzumab Course in Pooled CARE-MS I and II Patients by Number of Additional Courses Received Through Year 9

Regina Berkovich,^{1,2} Raed Alroughani,³ Ann D. Bass,⁴ Aaron L. Boster,⁵ Giancarlo Comi,⁶ Ho Jin Kim,⁷ Volker Limmroth,⁸ Jan Lycke,⁹ Richard A.L. Macdonell,¹⁰ Sven Schippling,¹¹ Basil Sharrack,¹² Mar Tintoré,¹³ Anthony Traboulsee,¹⁴ Patrick Vermersch,¹⁵ Heinz Wiendl,¹⁶ Tjalf Ziemssen,¹⁷ Nadia Daizadeh,¹⁸ Alan Jacobs,¹⁸ Elizabeth M. Poole,¹⁸ Barry A. Singer,¹⁹ on behalf of the CARE-MS I, CARE-MS II, CAMMS03409, and TOPAZ investigators

¹Keck School of Medicine, University of Southern California, Los Angeles, CA; ²Regina Berkovich, MD, PhD, Inc, West Hollywood, CA; ³Amiri Hospital, Sharq, Kuwait; ⁴Neurology Center of San Antonio, San Antonio, TX; ⁵Boster MS Center, Columbus, OH; ⁶University Vita-Salute San Raffaele, Milan, Italy; ⁷Research Institute and Hospital, National Cancer Center, Goyang, Korea, Republic of (South); ⁸Klinik für Neurologie und Palliativmedizin, Cologne, Germany; ⁹Institute of Neuroscience and Physiology, University of Gothenburg, Gothenburg, Sweden; ¹⁰Austin Health and Florey Institute of Neuroscience and Mental Health, Melbourne, VIC, Australia; ¹¹University Hospital Zürich and University of Zürich, Zürich, Switzerland; ¹²NIHR Sheffield Biomedical Research Centre, Sheffield Teaching Hospitals, University of Sheffield, Sheffield, United Kingdom; ¹³University Hospital Vall d'Hebron, Barcelona, Spain; ¹⁴The University of British Columbia, Vancouver, BC, Canada; ¹⁵University of Lille, INSERM U995, CHU Lille, Lille, France; ¹⁶Department of Neurology, University of Münster, Münster, Germany; ¹⁷Center for Clinical Neuroscience, Carl Gustav Carus University Hospital, Dresden, Germany; ¹⁸Sanofi, Cambridge, MA; ¹⁹The MS Center for Innovations in Care, Missouri Baptist Medical Center, St Louis, MO

Background: In CARE-MS I and II (trial registration: NCT00530348, NCT00548405), alemtuzumab treatment (12 mg/day; baseline: 5 days; 12 months later: 3 days) improved clinical and magnetic resonance imaging (MRI) outcomes vs subcutaneous interferon beta-1a over 2 years in patients with relapsing-remitting multiple sclerosis (MS). In 2 consecutive extensions (NCT00930553, NCT02255656 [TOPAZ]), patients could receive additional alemtuzumab (12 mg/day; 3 days; ≥12 months apart). **Objectives:** Evaluate yearly efficacy and safety of alemtuzumab in pooled CARE-MS patients who did or did not receive additional alemtuzumab through year 9. **Methods:** Pooled CARE-MS patients were stratified by the total number of courses received (exactly 2 courses, exactly 3 courses, ≥4 courses). Inclusion criteria: additional alemtuzumab (ie, courses 3 or 4) received by month 97 to allow ≥12 months of follow-up; no other disease-modifying therapy through year 9. Data were censored at course 5 (if received) in the ≥4 courses group. Outcomes data were rebaselined after the last alemtuzumab course. **Results:** 742/811 (91%) alemtuzumab-treated patients entered the extension and could receive additional courses; courses 3 and 4 were given most frequently in years 3 (19%) and 4 (6%), respectively. Of 742 extension patients, 359 (48%), 148 (20%), and 121 (16%) were included in the 2-, 3-, and ≥4-courses groups, with 303, 76, and 15 remaining on study in year 4 after last

course, respectively. Over 4 years after last course, annualized relapse rate was 0.07, 0.08, and 0.23 in the 2-, 3-, and ≥4-courses groups, respectively, and change in mean Expanded Disability Status Scale score at year 4 vs after last course was -0.06, +0.08, and +0.56, respectively. Over 4 years, 83%, 85%, and 94% were free of 6-month confirmed disability worsening, and 23%, 11%, and 15% had 6-month confirmed disability improvement in the 2-, 3-, and ≥4-courses groups, respectively. In year 4, 78%, 73%, and 69% were free of MRI disease activity. Serious adverse events were generally similar between cohorts during years 1-3 after last treatment (5.1%-11.4% per year), but low patient numbers in the ≥4-courses group confounded analysis of serious adverse events in year 4 after last course. **Conclusions:** Efficacy of additional alemtuzumab was maintained over 4 years after last course in CARE-MS patients, although the ≥4-courses group had higher disease activity and disability, as expected. Alemtuzumab safety was generally consistent between groups, except for the ≥4-courses cohort in year 4 after last course wherein interpretation was limited by low numbers of available patients.

Supported by: Sanofi, Bayer HealthCare Pharmaceuticals

Disclosure: *Regina Berkovich:* Acorda, Avanir, Bayer, Biogen, Novartis, Questcor, Sanofi, Teva (advisory boards, consulting fees). *Raed Alroughani:* Bayer, Biogen, GlaxoSmithKline, Lundbeck, Merck Serono, Novartis, Roche, Sanofi (speaking honoraria, research grants, scientific advisory boards). *Ann D. Bass:* Actelion, Biogen, EMD Serono, Mallinckrodt, Novartis, Roche-Genentech, Sanofi, TG Therapeutics (consulting fees/fees for non-CME services from commercial interests or their agents/grant and research support). *Aaron L. Boster:* Biogen, Mallinckrodt, Medtronic, Novartis, Sanofi, Teva (consulting fees and/or fees for non-CME services). *Giancarlo Comi:* Almirall SpA, Biogen, Biogen Italia Srl, Celgene, EXCEMED, F. Hoffman-La Roche, Forward Pharma, Genzyme, Genzyme Europe, MedDay, Merck KGaA, Merck Serono SpA, Roche SpA (consulting fee); Bayer, Biogen Dompé, Merck Serono, Novartis, Serono Symposia International Foundation, Teva (lecture fees); Sanofi (consulting fee, lecture fees). *Ho Jin Kim:* Alexion, Celltrion, Eisai, HanAll BioPharma, Merck Serono, Novartis, Sanofi, Teva-Handok, Viela Bio (consulting and speaking fees); Journal of Clinical Neurology, Multiple Sclerosis Journal (coeditor/associate editor); MedImmune/Viela Bio (steering committee); National Research Foundation of Korea (research support). *Volker Limmroth:* Bayer, Biogen, Merck Serono, Novartis, Roche, Sanofi, Teva (honoraria for consulting and speaking at symposia with approval by HR Department of Cologne General Hospital and University of Cologne). *Jan Lycke:* Acta Neurologica Scandinavica (editorial board); Almirall (scientific advisory boards); Biogen, Novartis, Teva (scientific advisory boards, travel support and/or lecture honoraria, unconditional research grants); Merck, Sanofi (scientific advisory boards, travel support and/or lecture honoraria). *Richard A.L. Macdonell:* Bayer, Merck, Roche (compensation for advisory boards and/or speaking fees); Biogen, Novartis, Sanofi, Teva (compensation for advisory boards and/or speaking fees, research support); Merck Serono (research support). *Sven Schippling:* Biogen, Merck Serono, Teva (consulting and/or speaking fees); Novartis, Sanofi (consulting and/or speaking fees, grant/research support). *Basil Sharrack:* Biogen, Merck, Novartis, Roche, Sanofi, Teva (research and travel grants, honoraria for expert advice on MS, and speaking fees). *Mar Tintoré:* Almirall, Bayer, Biogen, Genzyme, Merck Serono, Novartis, Roche, Sanofi-Aventis, Teva (speaking honoraria and travel expenses for scientific meetings). *Anthony Traboulsee:* Biogen, Chugai, Roche, Sanofi, Teva (consulting and/or speaking fees, grant/research support). *Patrick Vermersch:* Almirall, Biogen, Celgene, Merck, Novartis, Roche, Sanofi, Servier, Teva (consulting and/or speaking fees, research support). *Heinz Wiendl:* Bayer, Behring, Biogen, EMD Serono, Fresenius Medical Care, Merck Serono, Roche, Sanofi, Teva (consulting and/or speaking fees); Huber-Verlag (license fee payments); Neotope Biosciences, PML Consortium (grant/research support); Novartis (consulting and/or speaking fees, grant/research support). *Tjalf Ziemssen:* Almirall, Bayer, Merck, Roche (consulting and/or speaking fees); Biogen, Novartis, Sanofi, Teva (consulting and/or speaking fees, grant/research support). *Nadia Daizadeh, Alan Jacobs, Elizabeth M. Poole:* Sanofi (salary). *Barry A. Singer:* AbbVie, Biogen, Novartis, Roche, Sanofi (research support, speaking and/or consulting); Acorda, Alexion, Bayer, Celgene, EMD Serono, Genentech, Teva, TG Therapeutics (speaking and/or consulting); Alkermes, MedImmune (research support).

Keywords: Disease-modifying treatments in MS

(DMT03) Efficacy and Safety of Ofatumumab Versus Teriflunomide in Patients with Relapsing Multiple Sclerosis: Phase 3 ASCLEPIOS I and II Trials

Anne H. Cross,¹ Ludwig Kappos,² Amit Bar-Or,³ Jeffrey A. Cohen,⁴ Giancarlo Comi,⁵ Jorge Correale,⁶ Patricia K. Coyle,⁷ Jerome de Seze,⁸ David Leppert,⁹ Xavier Montalban,^{10,11} Krzysztof W. Selmaj,¹² Heinz Wiendl,¹³ Cecile Kerlough,¹⁴ Roman Willi,¹⁴ Bingbing Li,¹⁵ Algarda Kakariaka,¹⁴ Davorka Tomić,¹⁴ Alexandra Goodyear,¹⁵ Ratanakar Pingili,¹⁵ Dieter A. Haering,¹⁴ Krishnan Ramanathan,¹⁴ Martin Merschhemke,¹⁴ Stephen L. Hauser¹⁶

¹Department of Neurology, Division of Neuroimmunology, Washington University School

of Medicine, St Louis, MO; ²Neurologic Clinic and Polyclinic, Departments of Medicine, Clinical Research, Biomedicine and Biomedical Engineering, University Hospital of Basel, Basel, Switzerland; ³Center for Neuroinflammation and Experimental Therapeutics and Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ⁴Department of Neurology, Mellen MS Center, Neurological Institute, Cleveland Clinic, Cleveland, OH; ⁵University Vita-Salute San Raffaele, Milan, Italy; ⁶Institute for Neurological Research Dr. Raul Carrea, Buenos Aires, Argentina; ⁷Stony Brook University, Stony Brook, NY; ⁸University Hospital of Strasbourg, Strasbourg, France; ⁹Neurologic Clinic and Polyclinic, Departments of Medicine, Clinical Research, Biomedicine and Biomedical Engineering, University Hospital and University of Basel, Basel, Switzerland; ¹⁰Centre d'Esclerosi Múltiple de Catalunya (Cemcat), Hospital Universitario Vall d'Hebron, Barcelona, Spain; ¹¹St. Michael's Hospital, University of Toronto, Toronto, ON, Canada; ¹²Department of Neurology, Medical Academy of Lodz, Lodz, Poland; ¹³Department of Neurology, University of Münster, Münster, Germany; ¹⁴Novartis Pharma AG, Basel, Switzerland; ¹⁵Novartis Pharmaceuticals Corporation, East Hanover, NJ; ¹⁶University of California, San Francisco, San Francisco, CA

Background: Ofatumumab is the first fully human anti-CD20 monoclonal antibody, administered with a monthly 20 mg subcutaneous dosing regimen. **Objectives:** To investigate the efficacy and safety of ofatumumab vs teriflunomide in patients with relapsing multiple sclerosis (MS). **Methods:** ASCLEPIOS I and II were 2 identical phase 3, double-blind, double-dummy, active comparator-controlled, parallel-group, innovative, adaptive-design (with flexible duration), multicenter trials in patients aged 18-55 years with an Expanded Disability Status Scale score of 0-5.5 at screening. Patients were randomized (1:1) to receive subcutaneous ofatumumab 20 mg (loading dose: days 1, 7, and 14; maintenance dose: every 4 weeks from week 4) or oral teriflunomide 14 mg once daily, for up to 30 months. The primary end point was annualized relapse rate. Key secondary end points included 3- and 6-month confirmed disability worsening (3mCDW/6mCDW), 6-month confirmed disability improvement (6mCDI), magnetic resonance imaging-related outcomes, and serum neurofilament light chain (NFL) levels. Safety and tolerability were also assessed. **Results:** Of 1882 enrolled patients (ASCLEPIOS I/II: N = 927/955), 1578 completed the core study. Ofatumumab reduced annualized relapse rate (ASCLEPIOS I and II: 50.5% and 58.5%), gadolinium-enhancing T1 lesions (97.5% and 93.8%), and new/enlarging T2 lesions (82.0% and 84.5%) vs teriflunomide (all, $P < .001$). In the pre-specified ASCLEPIOS I/II pooled analysis, ofatumumab reduced the risk of 3mCDW by 34.4% ($P = .002$) and 6mCDW by 32.5% ($P = .012$), and numerically increased the probability to achieve 6mCDI by 35.2% ($P = .094$), vs teriflunomide. Ofatumumab reduced serum NFL levels vs teriflunomide in the first measurement at month 3 (ASCLEPIOS I, $P = .011$; ASCLEPIOS II, $P < .001$) and in all subsequent assessments (all, $P < .001$). No difference in the slope of brain volume change from baseline was observed between treatments ($P = .116$ [ASCLEPIOS I] and 0.129 [ASCLEPIOS II] vs teriflunomide). Adverse events occurred in 83.6% and 84.2% of patients receiving ofatumumab and teriflunomide, respectively. Systemic injection-related reactions occurred in 20.6% and 15.3% of ofatumumab and teriflunomide-treated patients, respectively. Rates of serious infections (ofatumumab, 2.5%; teriflunomide, 1.8%) and malignancies (0.5% and 0.3%, respectively) were low. **Conclusions:** Ofatumumab demonstrated superior efficacy vs teriflunomide, with an acceptable safety profile, in patients with relapsing MS.

Supported by: None

Disclosure: *Anne H. Cross:* Academic CME (CE provider) (speaking, slide preparation for CME); Biogen, Celgene, TG Therapeutics (consulting fee); EMD Serono (consulting fee, contracted research); Genentech/Roche (consulting fee, contracted research, speakers' bureau); Novartis (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent, scientific advisory board for ASCLEPIOS I and II); Projects in Knowledge (CE provider) (preparation of educational manuscripts, activities); Rockpointe (Potomac Center for Medical Education) (CE provider) (speakers' bureau for CME talks). *Ludwig Kappos:* Actelion, Mimoryx, Receptos, Santhera (consulting fee, steering committee; advisory board); Allergan, Allmirall, CSL Behring, Pfizer (support for educational activities); Bayer HealthCare (consulting fee, speakers' bureau, steering committee; advisory board; support for educational activities; license fees for Neurostatus products; grants); Biogen, Merck (consulting fee, steering committee; advisory board; support for educational activities; license fees for Neurostatus products; grants); Celgene, Genzyme (consulting fee, steering committee; advisory board; support for educational activities); INNO-Swiss, Roche research foundation (grants); Novartis, Roche (consulting fee, contracted research, speakers' bureau, steering committee; advisory board; support for educational activities; license fees for Neurostatus products; grants); Sanofi (consulting fee, speakers' bureau, steering committee; advisory board; support for educational activities); Teva (consulting fee, steering committee; advisory board; support for educational activities; license fees for Neurostatus products). *Amit Bar-Or:* Atara Biotherapeutics, Biogen, Celgene, EMD Serono, Genentech/Roche, GlaxoSmithKline, Mapi, MedImmune, Merck, Novartis, Receptos, Sanofi Genzyme (consulting fee, grant support, speakers'

bureau). *Jeffrey A. Cohen:* Adamas, Convelo, Mylan, Population Council (consulting fee); Multiple Sclerosis Journal (co-editor). *Giancarlo Comi:* Almirall SpA, Biogen, Biogen Italia Srl, Celgene Group, EXCEMED, F. Hoffman-La Roche, Forward Pharma, Genzyme Corp, MedDay, Merck KGaA, Merck Serono SpA, Novartis, Roche SpA, Sanofi Genzyme, Teva Italia Srl, Teva Pharmaceutical Industries Ltd (consulting fee); Genzyme Europe (consulting fee, speakers' bureau). *Jorge Correale:* Biogen, Genzyme, Merck-Serono, Novartis, Roche (board member, contracted research). *Patricia K. Coyle:* Accordant, Alexion, Bayer, Biogen, Celgene, Sanofi Genzyme, GlaxoSmithKline, Mylan, Serono, TG Therapeutics (consulting fee); Actelion, Alkermes, Corrona LLD, MedDay, National Institute of Neurological Disorders and Stroke, PCORI (contracted research); Genentech/Roche, Novartis (consulting fee, contracted research). *Jerome de Seze:* Alexion, Allergan, Almirall, Bayer, Biogen, Chugai, CSL Behring, F. Hoffmann-La Roche Ltd, Genzyme, LFB, Merck, Novartis, Teva (consulting fee, expert on advisory boards). *David Leppert:* Orion, Quanterix, Sanofi (consulting fee, speakers' bureau, travel reimbursement). *Xavier Montalban:* AbbVie, Teva Pharmaceutical (contracted research); Actelion, Celgene, EMD Serono, Genzyme, Immunic, Nervgen, TG Therapeutics (consulting fee); Biogen, MedDay, Merck, Novartis, Roche, Sanofi Genzyme (consulting fee, contracted research). *Krzysztof W. Selmaj:* Biogen, Celgene, F. Hoffmann-La Roche Ltd, Genzyme, Merck, Novartis, Synthon (honoraria for advisory boards). *Heinz Wiendl:* Bayer HealthCare, Biogen, Fresenius Medical Care, GlaxoSmithKline (consulting fee, honoraria). *Cecile Kerloeguen, Roman Willi, Bingbing Li, Algirdas Kakarietka, Daworka Tomic, Alexandra Goodyear, Ratnakar Pingili, Dieter A. Haering, Krishnan Ramanathan, Martin Merschhemke:* Novartis (salary). *Stephen L. Hauser:* Alector (scientific advisory board member [stock options received]); Annexon, Bionure, Molecular Stethoscope, Symbiotix (scientific advisory board member [stock options previously received]); F. Hoffmann-La Roche (travel support/reimbursement and writing support for anti-CD20 meetings and presentations); Neurona (board of trustees member [stock options previously received]); Novartis (travel support/reimbursement for anti-CD20 meetings and presentations).

Keywords: Disease-modifying treatments in MS, Efficacy and safety, Immunology and MS

(DMT04) Treatment-Emergent Adverse Events Occurring Early in the Treatment Course of Cladribine Tablets in Two Phase 3 Trials in Multiple Sclerosis

Jiwon Oh,¹ Bryan Walker,² Gavin Giovannoni,³ Dominic Jack,⁴ Fernando Dangond,⁵ Axel Nolting,⁴ Julie Aldridge,⁵ Lori Lebson,⁶ Thomas P. Leist⁷

¹Keenan Research Center for Biomedical Sciences, St. Michaels Hospital, Toronto, ON, Canada; ²Neurological Department, Duke University School of Medicine, Durham, NC; ³Bilzard Institute of Cell and Molecular Science, London, United Kingdom; ⁴Merck KGaA, Darmstadt, Germany; ⁵EMD Serono, Inc, Billerica, MA; ⁶EMD Serono, Inc, Rockland, MA; ⁷Jefferson University Hospital, Philadelphia, PA

Background: Tolerability and adherence to disease-modifying drugs (DMDs) can be influenced by treatment-emergent adverse events (TEAEs) that start soon after therapy initiation. One potential advantage of cladribine tablets is that patients only receive doses for two 4- to 5-day periods/treatment year. **Objectives:** To identify TEAEs early in the course of treatment in patients enrolled in the phase 3 CLARITY and ORACLE-MS clinical trials. **Methods:** This was a post hoc analysis of safety populations in CLARITY and ORACLE-MS studies. Patients received cladribine tablets 3.5 mg/kg (cumulative dose over 2 years; N = 636) or placebo (N = 641). The incidence of early adverse events, TEAEs, serious TEAEs, drug-related TEAEs, and TEAEs leading to discontinuation were summarized based on incidence within 2, 6, and 12 weeks (W) after commencement of therapy. **Results:** The incidence of TEAEs occurring within the first 2-12W of treatment across both trials in both treatment groups was generally low, and most events were mild (placebo: 68.4%-53.8%; cladribine tablets: 68.0%-54.4%). The most common TEAEs by time epoch after initiating placebo and cladribine tablets 3.5 mg/kg treatment, respectively, were nausea: 3.3% vs 4.9% (2W), 3.7% vs 6.4% (6W), and 4.5% vs 8.0% (12W); fatigue: 2.0% vs 1.4% (2W), 3.1% vs 2.5% (6W), and 4.4% vs 3.1% (12W); headache: 8.3% vs 9.0% (2W), 11.9% vs 14.8% (6W), and 15.1% vs 18.4% (12W); lymphopenia: 0.0% vs 2.5% (6W) and 0.5% vs 6.8% (12W); leukopenia: 0.0% vs 1.3% (12W). Other end points to be shown in the final presentation. **Conclusions:** Incidence of TEAEs experienced during the first 12 weeks of treatment with cladribine tablets in phase 3 clinical trials was low and mostly mild. Nausea, headache, and lymphopenia were seen more frequently in patients treated with cladribine tablets. These findings suggest that cladribine tablets are generally well tolerated, which may facilitate treatment adherence.

Supported by: None

Disclosure: *Jiwon Oh:* Biogen, Roche, Sanofi Genzyme (consulting fee, research funding); Brain Canada, MS Society of Canada (research funding); Celgene,

EMD Serono, Novartis (consulting fee). *Bryan Walker*: Biogen, Celgene, EMD Serono, Novartis, Sanofi Genzyme (consulting fee). *Gavin Giovannoni*: AbbVie, Actelion, Almirall, Atara Bio, Bayer Schering Pharma, Five Prime, GlaxoSmith-Kline, GW Pharma, Merck KGaA, Pfizer Inc, Protein Discovery Laboratories, Sanofi Genzyme, Teva Pharmaceuticals Industries Ltd, UCB, Vertex Pharmaceuticals (consulting fee); Biogen, Ironwood, Merck and Co, Novartis (consulting fee, unrelated research support). *Dominic Jack, Axel Nolting*: Merck KGaA (salary). *Fernando Dangond, Julie Aldridge, Lori Lebson*: EMD Serono Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). *Thomas P. Leist*: Alkermes, Bayer, Biogen, EMD Serono, Genentech, Genzyme, Novartis (consulting fee).

Keywords: Disease-modifying treatments in MS

PSYCHOSOCIAL: COGNITION

(PSY01) A Mindfulness Group Intervention in Newly Diagnosed Persons with Multiple Sclerosis: A Pilot Study

Sarah A. Morrow,¹ Nancy Vording,¹ Jordan Ward,¹ Courtney S. Casserly,² Heather Rosehart,³ Arlene Macdougall⁴

¹Clinical Neurological Sciences, London Health Sciences Centre, London, ON, Canada;

²Clinical Neurological Sciences, Western University, London, ON, Canada; ³Clinical Neurological Sciences, University of Western Ontario, London Health Sciences Center, London, ON, Canada; ⁴Psychiatry and Epidemiology & Biostatistics, St. Joseph's Health Care, Parkwood Institute, London, ON, Canada

Background: Relapsing multiple sclerosis (RMS) is a lifelong disease without a cure, usually diagnosed between age 20-40 years. Being newly diagnosed with RMS is a highly stressful event due to the unpredictable disease course after diagnosis. Thus, it is imperative that persons with multiple sclerosis have the skills and support to cope with the negative physical and emotional effects of the disease. **Objectives:** To assess whether a mindfulness-based intervention (MBI) will lessen the negative consequences of stress due to being newly diagnosed with RMS. **Methods:** A single-blind, randomized, prospective study of a 10-week MBI vs usual standard of care alone in persons newly diagnosed (within 1 year) with RMS. Primary outcomes included the Brief COPE measure and the Hospital Anxiety and Depression Scale (HADS). Secondary outcomes included measures of perceived stress, cognitive function, fatigue, and quality of life. Subjects were assessed at baseline, postintervention, and 6 months later. Analysis of covariance was used to compare longitudinal changes, with baseline scores used as covariates. **Results:** 25 subjects were recruited (16 MBI, 9 controls); most were women (21 [84%]), with a mean age of 38.4 ± 9.5 years. The groups were well matched on baseline characteristics. All controls completed the study, while 4 MBI participants did not. The MBI group improved significantly on the COPE measure when compared to the control group ($P = .024$) as well as on the HADS depression subscale ($P = .007$) pre- and postintervention; there was no significant difference over time on the HADS anxiety subscale ($P = .179$). On the secondary outcomes, there was a significant improvement on the Perceived Stress Scale ($P = .015$), and a trend towards improvement on the SF-36 ($P = .073$; quality of life) and the MSNQ ($P = .066$; perceived cognitive impairment) comparing pre- and postintervention assessments. Six-month data will be available at the time of this presentation. **Conclusions:** This pilot study demonstrates that an MBI improves coping, depression, and perceived stress in newly diagnosed persons with RMS.

Supported by: None

Disclosure: *Sarah A. Morrow*: Biogen, Novartis (contracted research); *Celgene* (consulting fee); *EMD Serono, Roche, Sanofi Genzyme* (speakers' bureau). *Nancy Vording, Jordan Ward, Courtney S. Casserly, Heather Rosehart, Arlene Macdougall*: Nothing to disclose.

Keywords: Comprehensive care and MS, Psychological issues and MS

(PSY02) Effects of Weekly Participation in a Wellness Program on Self-Reported Measures for People Living with Multiple Sclerosis: A 3-Year Analysis

Brian Hutchinson,¹ John Schafer,² Lacey Sayre,³ Tiffany Malone³

¹Multiple Sclerosis Achievement Center, Dignity Health, Sacramento, CA; ²Mercy MS Center, Carmichael, CA; ³Multiple Sclerosis Achievement Center, Dignity Health, Citrus Heights, CA

Background: The Multiple Sclerosis Achievement Center (MSAC) conducts day wellness programs to address physical, cognitive and social well-being. Program activities include individualized and group exercise, cognitive stimulation, education, socialization, and community outings. Baseline, 1-year, and 2-year follow-up data were collected and presented at previous Consortium of Multiple Sclerosis Centers annual meetings. **Objectives:** To determine, through the use of patient-reported

outcome (PRO) measures, if members of these wellness programs improve in areas of self-reported disease impact and quality of life over a 3-year period. **Methods:** Initial analysis, comparing data of baseline and 1-year participation in these wellness programs, was completed through paper/pencil outcome measures between December 2016 and August 2017 for 95 people with multiple sclerosis (PwMS). Of those 95, 2-year data for 70 PwMS were collected in January 2019 and 3-year data are anticipated for 66 people. Outcome measures used for these analyses include the Multiple Sclerosis Impact Scale (MSIS-29), Multiple Sclerosis Self-Efficacy Scale-10 item (MSSE), Godin Leisure Time Exercise Questionnaire (GLTEQ), and Neuro-QoL (questions from the Anxiety, Depression, Emotion & Behavior, Positive Affect, Cognition, Ability to Participate, and Social Roles sections were used). All outcomes were completed onsite at the MSAC as part of the members' weekly participation in the program. Analysis will be completed to compare data from the initial analysis to the 3-year results. **Results:** As previously reported, a correlation between reports of Self-Efficacy, Anxiety, Ability to Participate, and Positive Affect (per MSSE and Neuro-QoL) were seen with both 1- and 2-year analyses. Analysis of baseline to 2-year data demonstrated statistically significant changes in Neuro-QoL sections Ability to Participate ($P = .02$) and Social Roles ($P = .001$). Another notable change in the 2-year analysis was an increase in physical activity, as measured by GLTEQ (average change of 2.34 from baseline to 2 years). Data analysis will measure any changes in 3-year data compared to baseline, 1-year, and 2-year results. **Conclusions:** Complete collection and analysis of 3-year comparative data will be finalized in February 2020 for presentation at the meeting.

Supported by: None

Disclosure: *Brian Hutchinson*: Biogen (speakers' bureau). *John Schafer*: Biogen, EMD Serono, Genentech, Sanofi Genzyme, Teva Neuroscience (speakers' bureau). *Lacey Sayre, Tiffany Malone*: Nothing to disclose.

Keywords: Comprehensive care and MS, Psychological issues and MS, Wellness

(PSY03) Examining Multilevel Environmental Correlates of Physical Activity Among Older Adults with Multiple Sclerosis

Stephanie L. Silveira, Jessica F. Baird, Robert W. Motl

Physical Therapy, University of Alabama at Birmingham, Birmingham, AL

Background: With a growing population of older adults with multiple sclerosis (MS), appropriate strategies are needed to promote physical activity (PA) as a second-line approach for symptom management. **Objectives:** This cross-sectional study examined built environment, social environment, and individual social cognitive theory (SCT) variables as hierarchical correlates of PA in older adults with MS using a social ecological model (SEM) framework. **Methods:** 363 participants completed the online survey including demographics, the Abbreviated Neighborhood Walkability Scale (NEWS-A), Social Provisions Scale (SPS), Exercise Self-Efficacy Scale (EXSE), Multidimensional Outcome Expectations Scale for Exercise Scale (MOEES), and Godin Leisure-Time Exercise Questionnaire (GLTEQ) Total and Health Contribution score (HCS). Spearman rank-order correlation analyses were used to examine associations among NEWS-A subscales, SPS, EXSE, MOEES, and GLTEQ Total and HCS. We then conducted hierarchical, linear regression analysis whereby we regressed GLTEQ with NEWS-A subscales (built environment) in step 1, SPS (social environment) in step 2, then EXES and MOEES (individual determinants) in step 3. **Results:** Land-use mix diversity, land-use mix access, street connectivity, and aesthetics were significantly correlated with GLTEQ Total, whereas land-use mix diversity, land-use mix access, infrastructure and safety for walking, aesthetics, and crime were significantly correlated with GLTEQ HCS. Hierarchical linear regression analyses were then conducted whereby we regressed GLTEQ Total with NEWS-A subscales (step 1) with significant associations noted for land-use mix diversity and aesthetics ($R^2 = 0.09$), step 2 included SPS with significant associations noted for SPS, land-use mix diversity, and aesthetics ($R^2 = 0.15$), and finally EXSE and MOEES were included in step 3 and were the only significant correlates of GLTEQ total ($R^2 = 0.38$). Regarding GLTEQ HCS, land-use mix diversity, aesthetics, and crime were significant correlates in step 1 ($R^2 = 0.10$), SPS and land-use mix diversity were the only significant correlates in step 2 ($R^2 = 0.14$), and EXSE was the only significant correlate in step 3 ($R^2 = 0.36$). **Conclusions:** This study provides guidance for researchers and practitioners on relevant targets for tailoring PA interventions for older adults with MS and supports the continued emphasis on self-efficacy as a primary predictor of health behavior change and maintenance.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Management of activities of daily living in MS, Older adults with MS, Psychological issues and MS

(PSY04) Effect of Nabiximols Cannabinoid Oromucosal Spray on Depressive Symptoms, Suicidality, and Cognition in Patients with Multiple Sclerosis

John DeLuca,¹ Dawn Langdon,² Joris Berwaerts,³ Joanne Wagner³

¹Kessler Foundation, West Orange, NJ; ²Department of Psychology, Clinical, Health and Social Psychology, Royal Holloway, University of London, Egham, United Kingdom; ³Greenwich Biosciences, Inc, Carlsbad, CA

Background: Substantial evidence has shown nabiximols, a complex botanical mixture containing delta-9-tetrahydrocannabinol and cannabidiol as the principal cannabinoids, can reduce spasticity associated with multiple sclerosis (MS). This analysis assesses whether nabiximols affects other patient outcomes such as depressive symptoms, suicidality, and cognition. **Objectives:** Report the effect of nabiximols on depression, suicidality, and cognition using data from 2 placebo-controlled randomized controlled trials, GWSP0604 (12 weeks) and GWMS1137 (48 weeks), in patients with spasticity due to MS. **Methods:** Mood and suicidality were assessed using the Beck Depression Inventory-II (BDI-II) in both trials. In GWMS1137, suicidality was assessed using the Columbia-Suicide Severity Rating Scale (C-SSRS) and working memory/processing speed using the Paced Auditory Serial Addition Test (PASAT). The combined PASAT total score was calculated combining both PASAT-3 and -2 tests scores (total of 120 points). Outcome differences between nabiximols and placebo are summarized. **Results:** 241 patients from GWSP0604 and 121 from GWMS1137 were included. The baseline and end-of-treatment mean BDI-II total scores were 8.7 vs 9.5 for nabiximols and 9.7 vs 10.4 for placebo in GWSP0604, and 15.7 vs 12.5 for nabiximols and 13.5 vs 11.1 for placebo in GWMS1137. Differences between nabiximols and placebo of the BDI-II change from baseline adjusted means were -0.06 (-1.62, 1.49) in GWSP0604 (no significant difference) and -0.29 (-2.91, 2.33) in GWMS1137 (statistically noninferior). Question 9 of BDI-II (suicidal thoughts or wishes) showed no notable treatment differences in either trial, with only 1 patient treated with nabiximols reporting a score ≥ 2 . On the C-SSRS, 3 (5.1%) patients randomized to placebo and 1 (1.6%) to nabiximols had a “flag” (ie, “yes” as a response), but further questioning revealed no emergent suicidal ideations or behavior in any of these patients. For GWMS1137, the baseline and end-of-treatment PASAT total score means were 71.3 vs 78.6 for nabiximols and 74.5 vs 82.7 for placebo; increases may reflect practice effects. Treatment difference of the adjusted mean was -1.47 (-6.41, 3.48), indicating nabiximols does not adversely affect working memory/cognitive processing speed in patients with MS over a 48-week period compared with placebo. **Conclusions:** Nabiximols had no notable effects on depression, suicidality, or working memory/processing speed in patients with MS.

Supported by: None

Disclosure: *John DeLuca:* Biogen (consulting fee, contracted research, speakers' bureau); Celgene, Novartis (consulting fee); EMD Serono (contracted research); Sanofi Genzyme (speakers' bureau). *Professor Dawn Langdon:* Nothing to disclose. *Joris Berwaerts:* GW Pharmaceuticals (salary). *Joanne Wagner:* Greenwich Biosciences/GW Pharmaceuticals (salary); GW Pharmaceuticals (ownership interest).

Keywords: Complementary/alternative therapies in MS, MS symptom management, Psychological issues and MS

(PSY06) Multiple Sclerosis Management: Predicting Disease Trajectory of Multiple Sclerosis on Multidimensional Data Including Digital Cognitive Assessments and Patient-Reported Outcomes Using Machine Learning Techniques

Mark Gudesblatt,¹ Jared Srinivasan,¹ Olivia Kaczmarek,¹ Daniel Kantor,² Daniel Golan,³ Myassar Zarif,¹ Barbara Bumstead,¹ Marijean Buhse,¹ Lori Fafard,¹ Timothy Fratto,⁴ Jeffrey Wilken,⁴ Cynthia Sullivan,⁵ Glen Doniger⁶

¹South Shore Neurologic Associates, Patchogue, NY; ²Florida Atlantic University, Boca Raton, FL; ³Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel; ⁴Washington Neuropsychology Research Group, Washington, DC; ⁵Nursing, State University of New York at Stony Brook, Stony Brook, NY; ⁶NeuroTrax Corp, Modiin, Israel

Background: Multiple sclerosis (MS) disease impact is traditionally measured by magnetic resonance imaging changes, relapse rates, and Expanded Disability Status Scale (EDSS). Combining multidimensional patient-reported outcome (PRO) and objective disease impact information independent of EDSS might enhance clinical decision making. As patient-tracking sources expand beyond what is traditionally captured in an office visit, clinicians need tools to help integrate these varied streams of data. Machine learning has the potential to help clinicians predict meaningful patient outcomes from multidimensional and quantified data sources. **Objectives:** To demonstrate the feasibility of predicting clinical outcomes in MS using standard machine learning methods on

multidimensional data including digital cognitive assessments and PROs. **Methods:** Machine learning models were trained on electronic health record data, cognitive domain scores, and PRO data. A prediction model was created given the patient's record. 80% of the dataset was used in training, 20% in testing with an ensemble learning method (random forest classifiers) used to construct a multitude of training decision trees, which then outputted the mean prediction of the individual trees. **Results:** The training set consisted of 258 persons with MS (72.5% female, average age 46.2 ± 10.2) over a 3-year period. Untuned models calculated F1 scores ($2 * [\text{Precision} * \text{Recall}] / [\text{Precision} + \text{Recall}]$), Precision, Recall and Accuracy for multiple models predicting PROs and disease-modifying therapy choice. The most precise and accurate models were for the Driving (0.913, 0.904, 0.942, 0.912), Modified Falls Efficacy Scale (0.789, 0.796, 0.792, 0.829), Depression (0.711, 0.765, 0.714, 0.718), Fatigue (0.716, 0.782, 0.699, 0.755), and Employment (0.672, 0.753, 0.668, 0.705). **Conclusions:** Machine learning combined with objective measures of disease impact and PROs can provide important information to predict economically important and disability-relevant outcomes, potentially enhancing treatment decisions. These results show promising predictive accuracy to be used in a variety of advisory applications and potentially reduce disease-related disability. The results of the other models demonstrate the feasibility of using machine learning in a broader network of clinical sites that will allow for greater accuracy, precision, and recall. The eventual goal is that these models can be used as an aid in the shared decision-making process, and to reduce inappropriate health care costs.

Supported by: None

Disclosure: *Mark Gudesblatt:* Acorda, Amgen, Medtronic, Saol Therapeutics (speakers' bureau); Biogen, EMD Serono, Novartis, Sanofi, Teva (contracted research). *Jared Srinivasan, Olivia Kaczmarek, Daniel Kantor, Daniel Golan, Marijean Buhse, Lori Fafard, Timothy Fratto:* Nothing to disclose. *Myassar Zarif:* Acorda, Biogen, Genzyme, Teva (speakers' bureau). *Barbara Bumstead:* Biogen, Genzyme (speakers' bureau). *Jeffrey Wilken:* Biogen (contracted research); EMD Serono (speakers' bureau); Genzyme (contracted research, speakers' bureau). *Cynthia Sullivan:* Roche (contracted research). *Glen Doniger:* NeuroTrax (salary).

Keywords: Comprehensive care and MS, Machine learning, Natural history of MS

REHABILITATION

(RH101) Significant Structural Neuroplasticity Changes Can Follow Physical Behavioral Change Therapy for Multiple Sclerosis

Victor W. Mark,¹ Brent M. Womble,² Gitendra Usawatte,² David M. Morris,³ Mary H. Bowman,² Staci McKay,² Edward Taub²

¹Physical Medicine and Rehabilitation, University of Alabama at Birmingham (UAB), Birmingham, AL; ²Psychology, UAB, Birmingham, AL; ³Physical Therapy, UAB, Birmingham, AL

Background: Constraint-induced movement therapy (CIMT) is a form of physical behavioral change therapy (BCT) that can significantly improve paretic limb use in the community in progressive multiple sclerosis (MS) for at least 1 year (Mark et al, 2018). Although a few forms of BCT can increase real-life physical activity in MS, none thus far has been examined for whether such treatment can change cerebral cortical grey matter structure. **Objectives:** To evaluate whether CIMT vs dose-matched control physical training can change cortical grey matter structure in progressive MS. **Methods:** Twenty adults with chronic MS matched for unilateral arm disability were randomized to 35 hours/2 weeks of either CIMT or a holistic complementary and alternative medicine (CAM) program (yoga, aquatic therapy, massage, relaxation techniques). Paretic limb use was measured with the Motor Activity Log (MAL), which has been validated against real-world upper limb accelerometry. Pre- and post-treatment 3-T structural brain magnetic resonance imaging scans were performed. Tensor-based morphometry (TBM) and voxel-based morphometry (VBM) were used to evaluate group-level changes in primary motor cortex (M1) structure contralateral to the more-affected arm. Whole-brain statistics were conducted using 1-sample *t* tests within Statistical Parametric Mapping software with a cluster-extent threshold of 10 voxels and false discovery rate of 0.1. **Results:** The 2 groups were identical in high expectancy to benefit. CIMT produced a much larger effect size ($d' = 3.2$) on the MAL than did CAM ($d' = 0.7$). TBM detected an increase in the thickness of M1 after CIMT but not after CAM. VBM detected a change in M1 after CIMT, suggesting an increase in cortical density or volume or both. No change was detected after CAM. **Conclusions:** TBM and VBM suggest that CIMT increases M1 thickness and either volume or density in

MS, unlike dose-matched CAM. The findings suggest for the first time that physical BCT can significantly stimulate cortical neuroplasticity in a degenerative central nervous system disorder. The findings accord with our previous findings of post-CIMT significant white matter structural improvement in progressive MS (Barghi et al, 2018) and grey matter increases in stroke (Gauthier et al, 2008) and cerebral palsy (Sterling et al, 2013). Together, these findings suggest that a specific form of physical BCT can not only stimulate physical activity in the community over the long term but also well improve neurologic structure for progressive MS.

Supported by: None

Disclosure: Nothing to disclose

Keywords: CNS repair, Imaging and MS, Management of activities of daily living in MS

(RH102) A New Look at the Symbol Digit Modalities Test in Multiple Sclerosis and Disabilities

Yilan Liu,¹ Victor W. Mark,² Sean Bowman,¹ Raazen M. Mahmud,¹ Janet Niemeier,¹ Stacey S. Cofield³

¹University of Alabama at Birmingham (UAB), Birmingham, AL; ²Physical Medicine and Rehabilitation, UAB, Birmingham, AL; ³Biostatistics, UAB, Birmingham, AL

Background: Reduced information processing speed (IPS) is the most common cognitive impairment in multiple sclerosis (MS), related to reduced employment and physical ability. The Symbol Digit Modalities Test (SDMT) is the gold standard measure of IPS in MS (Benedict et al, 2017). Although many investigators have suggested that performance on the SDMT involves multiple cognitive processes, there has been little attempt to tease apart what specific cognitive processes may affect the IPS score on the SDMT. Improved understanding of the cognitive processes that contribute to IPS on the SDMT could support future cognitive rehabilitation trials to treat impaired IPS in MS. **Objectives:** Assess specific eye movement measures during the SDMT to suggest what specific cognitive processes may underlie IPS on the SDMT in MS. **Methods:** We recruited a convenience sample of 38 adults with MS, without clinical oculomotor impairment, who performed the SDMT while an infrared eye tracker recorded their eye movements. Eye positions were sampled at 10 Hz. Data were exported to a database for subsequent specific eye movement measures: 1) search organization as calculated by the “best r” metric (index of primarily orthogonal search progress [Mark et al, 2004]); 2) total of upward saccades from the lower area on the page of test symbols to the answer key at the top, inferred as a measure of symbol working memory. Spearman rho was performed to assess correlations between the variables and the SDMT IPS score (correct responses in 90 seconds). **Results:** The mean (SD) SDMT IPS score was 38 (14). Both best r and total of upward saccades significantly positively correlated with the SDMT IPS score (best r, rho = 0.45, P = .004; upward saccades, rho = 0.62, P < .0001). **Conclusions:** Both search organization and total of upward saccades were positively correlated with the IPS score. The latter result was unexpected: the faster on the SDMT, the more often subjects checked the answer key. The results suggest that lower-scoring participants with MS, who less often checked the answer key during the test, may have become lost during their visual search, as reflected by their poorer search organization. Upward saccades may not so much represent working memory but rather as a strategy to assure successful test completion that is not effectively used by lower-scoring persons with MS. Further research will be needed to assess the criterion validity of specific SDMT eye movement measures relative to standard cognitive test assessments.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Information processing speed, Management of activities of daily living in MS

(RH103) Creating a Yoga Program as Part of a Comprehensive Multiple Sclerosis Care Model

Sandra Chapman, Ruth Almen, Carrie M. Hersh, Le H. Hua

Lou Ruvo Center for Brain Health, Cleveland Clinic, Las Vegas, NV

Background: Multiple sclerosis (MS) affects both physical and emotional health. Comprehensive programs addressing health and wellness are essential to improve quality of life. We offer a multidisciplinary approach to treatment, comprising a dedicated medical team, education, social work, rehabilitation, research, and a health and wellness program. The benefits of yoga are increasingly recognized and subsequently requested by patients, yet dedicated programs aimed specifically for persons with MS are limited. MS specialty centers are challenged to explore ways to include yoga into a comprehensive care plan. **Objectives:** We present our experience at the Cleveland Clinic Lou Ruvo Center for Brain Health

in creating the Yoga for MS program as an integral component to comprehensive care. **Methods:** Our center is contracted with a local yoga company to provide weekly sessions, using philanthropy funding. There are 3 instructors: 1 who leads and 2 others who assist, observe, and take notes. Because participants regularly engaged in dialogue before and after sessions, a 60-minute support/community sharing session was added prior to the yoga session. Our social worker attends each session and facilitates as needed. Topics include implications for physical and mental health, as well as coping strategies and the importance of treatment adherence. An instructional video and meditation literature are available for use at home. **Results:** The Yoga for MS Program started in November 2017, initially in a private setting with 1 instructor. In response to our patients’ needs, we hired a yoga company with multiple instructors at a yoga studio in October 2018. Our current space can comfortably support 18 participants, including wheelchairs, scooters, and walkers. To date, we have 12 participants who have a 75% attendance rate in a 12-week session. Average class size in the first year was 4, and now has doubled to 8, the largest being 14. Yoga staff observed over time increased flexibility among the participants. The participants also report reduced stress, increased use of coping strategies, and positive effects on strength, balance, self-esteem, and confidence. Participants also share resources and report feeling like they are “really in this together.” **Conclusions:** A Yoga for MS Program can address the physical and emotional well-being of patients with MS as part of a comprehensive care model. Future studies addressing the direct benefits of yoga on MS outcomes are being explored.

Supported by: None

Disclosure: Sandra Chapman, Ruth Almen: Nothing to disclose. Carrie M. Hersh: Biogen, Genentech (consulting fee, contracted research); EMD Serono (consulting fee); Genzyme, Novartis (consulting fee, speakers’ bureau); PCORI (contracted research). Le H. Hua: Biogen, Celgene, EMD Serono, Genentech, Novartis (consulting fee); Genzyme (consulting fee, speakers’ bureau).

Keywords: Complementary/alternative therapies in MS, Comprehensive care and MS, Yoga

(RH104) Feasibility of “Sit Less, Move More”: An Intervention for Reducing Sedentary Behavior Among African Americans with Multiple Sclerosis

Jessica F. Baird,¹ Jeffer E. Sasaki,² Brian M. Sandroff,³ Robert W. Mott¹

¹Physical Therapy, University of Alabama at Birmingham, Birmingham, AL; ²Department of Sports Sciences, Universidade Federal do Triangulo Mineiro, Bairro Tutunas, Brazil; ³Department of Biostatistics, School of Public Health, University of Alabama at Birmingham, Birmingham, AL

Background: Sedentary behavior (SB) is a major concern in multiple sclerosis (MS), as it may accelerate disease progression and exacerbate physical disability. This is especially concerning in African Americans, a segment of the MS population who present with greater neurologic disability than Caucasians and for whom little MS research data are available. **Objectives:** The current study examined the feasibility of an intervention focused on reducing SB in African Americans with MS. **Methods:** We recruited 30 ambulatory and physically inactive African Americans with MS (age = 44 years) to participate in the “Sit Less, Move More” (SLMM) program. SLMM consisted of a 12-week behavioral intervention that used text-messaging along with Social Cognitive Theory-driven newsletters and behavioral coaching for reducing SB. Feasibility was assessed on 4 domains: process, resource, management, and scientific outcomes. Participants wore activPAL (AP) and ActiGraph (AG) activity monitors at 3 time points (prior to week 1 [T1], during week 6 [T2], and after week 12 [T3]) to measure changes in time spent sitting (AP data) and time spent in sedentary behavior (AG data). Estimates of effect size (Cohen’s d) were calculated to describe the treatment effect of SLMM on SB. **Results:** Process: Of the 64 persons initially contacted, 45 were assessed for eligibility, 31 were sent the informed consent document, and 30 returned a signed informed consent document and were included in the study. Resources: All participants returned T2 testing materials, and 29 (95%) returned T3 testing materials. Twenty-five (83%) participated in all behavioral coaching sessions. Total study costs were \$7242.38 USD including costs for materials, postage, equipment, and participant remuneration. Management: Total personnel time to complete the study was 130 hours. Only 13 participants had valid AP data at all 3 time points, whereas 24 participants had valid AG data at all 3 time points. Scientific outcomes: No adverse events were reported. There was a small treatment effect on time spent sitting (d = -0.13) and sedentary time (d = -0.19). **Conclusions:** The SLMM intervention is safe and feasible for African Americans with MS, and yielded a small reduction in SB. The intervention was low-cost and well-received as an approach for reducing sedentary behavior, and, overall, our results suggest that the SLMM program prog-

ress towards a phase 2 trial to determine its efficacy for reducing SB in African Americans with MS.

Supported by: None

Disclosure: *Jessica F. Baird, Jeffer E. Sasaki, Brian M. Sandroff, Robert W. Motl: Nothing to disclose. Gary Cutter: Avexis Pharmaceuticals, Biogen, BioLineRx, Brainstorm Cell Therapeutics, Click Therapeutics, CSL Behring, Galmed Pharmaceuticals, Genentech, Genzyme, Gilgamesh Pharmaceuticals, GW Pharmaceuticals, Hisun Pharmaceuticals, Horizon Pharmaceuticals, Klein Buendel Inc, MedDay, MedImmune, Merck, Merck/Pfizer, Newrim, National Heart, Lung, and Blood Institute (protocol review committee); National Institute of Child Health and Human Development (OPRU oversight committee); Novartis, Orphazyme, Opko Biologics, Osmotica Pharmaceuticals, Perception Neurosciences, Reata Pharmaceuticals, Receptos/Celgene, Recursion Pharmaceuticals, Roche, Sanofi-Aventis, Somahlution, Teva Pharmaceuticals, TG Therapeutics, Vivus (fees for non-CME/CE services received directly from commercial interest or its agent); MGFA MG Registry, NARCOMS (contracted research); Pythagoras Inc (ownership interest).*

Keywords: Sedentary behavior

(RHI05) The Effect of Aerobic Fitness on Physical and Cognitive Function and Brain Volume in Older Adults with Multiple Sclerosis

Jessica F. Baird,¹ Marcias Bamman,² Cynthia Brown,² John R. Rinker,³ Robert W. Motl¹

¹Physical Therapy, University of Alabama at Birmingham (UAB), Birmingham, AL; ²UAB, Birmingham, AL; ³Neurology, UAB, Birmingham, AL

Background: There is evidence for the beneficial effects of aerobic exercise training on physical and cognitive function in persons with multiple sclerosis (MS). Improvements in function may be associated with the effects of aerobic fitness on deep gray matter (DGM) structures within the brain such as the hippocampus, thalamus, and basal ganglia. To date, we are unaware of research that has examined the effects of aerobic fitness in older adults with MS. Given the aging of the MS population, such an investigation is warranted. **Objectives:** The current study examined the effect of aerobic fitness on physical and cognitive function and DGM brain structures relevant to these outcomes in older adults with MS. **Methods:** We recruited ambulatory adults (age 55+ years) with MS (n = 20, age = 63 years). All participants underwent an assessment of aerobic fitness using a maximal, incremental exercise test on a recumbent stepper, assessments of walking speed (Timed 25-Foot Walk) and walking endurance (6-Minute Walk), assessments of cognitive function (Symbol Digit Modalities Test [SDMT]; Brief Visuospatial Memory Test [BVM]; California Verbal Learning Test [CVLT]), and a 3-T magnetic resonance image of the brain. We dichotomized participants into a "low fit" group (n = 10) and "high fit" group (n = 10) based on the results of the exercise test and calculated effect sizes (Cohen's *d*) between the groups for all outcome measures. **Results:** Aerobic fitness had a large effect on both walking speed ($d = -0.99$) and walking endurance ($d = 1.51$). There was a moderate effect of aerobic fitness on cognitive function (SDMT $d = 0.57$; CVLT $d = 0.48$; BVM $d = 0.67$). The effect of aerobic fitness on DGM brain structures varied by structure. There was little to no effect on the thalamus ($d = 0.19$) and hippocampus ($d = -0.01$), whereas there was a moderate effect on the basal ganglia ($d = 0.53$). **Conclusions:** Our results provide novel evidence demonstrating a positive effect of aerobic fitness on physical and cognitive function in older adults with MS. As aerobic fitness is modifiable by aerobic exercise training, our results suggest that participation in regular physical activity may be an approach to ameliorate the consequences of aging with MS. Our results further suggest that

improvements in function may be mediated by an effect of aerobic fitness on DGM brain structures; however, additional research is warranted to comprehensively investigate the neural adaptations associated with aerobic fitness in this population.

Supported by: None

Disclosure: *Jessica F. Baird, Marcias Bamman, Cynthia Brown, Robert W. Motl: Nothing to disclose. John R. Rinker: Biogen, EMD Serono (contracted research).*

Keywords: Exercise, Imaging and MS

(RHI06) Functional Electrical Stimulation Cycling Exercise Reduces Lower Limb Strength Asymmetry in Persons with Multiple Sclerosis

John W. Farrell III,¹ Thomas Edwards,² Robert W. Motl,³ Lara A. Pilutti¹

¹Interdisciplinary School of Health Sciences, University of Ottawa, Ottawa, ON, Canada;

²School of Human Kinetics, University of Ottawa, Ottawa, ON, Canada; ³Physical Therapy, University of Alabama at Birmingham, Birmingham, AL

Background: Lower limb strength asymmetries (ie, difference $\geq 10\%$ between contralateral muscle groups) have been associated with mobility impairment in persons with multiple sclerosis (PwMS), and may be a target for exercise training interventions aiming to improve mobility. Functional electrical stimulation (FES) cycling is an adapted exercise modality that has demonstrated preliminary benefits for mobility and fitness outcomes in PwMS with mobility impairment, but its potential effects on lower limb strength asymmetry remain unknown. **Objectives:** To assess the effect of FES cycling exercise on lower limb strength asymmetry in PwMS who have mobility impairment (ie, Expanded Disability Status Scale [EDSS] score 5.5-6.5), and to explore associations between change in lower limb strength asymmetry and change in mobility outcomes. **Methods:** Peak torque was recorded bilaterally for knee extensors (KEs) and flexors (KFs) using an isokinetic dynamometer, and was then used to generate lower limb strength asymmetry scores ($1 - (\text{torque}_{\text{weak}} / \text{torque}_{\text{strong}}) \times 100$). Mobility outcomes included the Timed 25-Foot Walk (T25FW) and the 2-Minute Walk (2MW). Participants received 24 weeks (3x/week) of either FES cycling or passive leg cycling (PLC). The FES condition actively cycled while receiving mild electrical stimulation to the quadriceps, hamstrings, and gluteal muscle groups. Exercise intensity was set at 40% to 60% of heart rate reserve, with exercise duration gradually increasing from 10 to 40 minutes per session over the course of the intervention. The PLC condition was identical to the FES condition, but did not receive electrical stimulation and did not actively cycle. **Results:** Eight PwMS (mean [SD] age = 52.9 [7.9]; median [IQR] EDSS score = 6.3 [0.5]) completed the intervention. The FES cycling condition demonstrated a small decrease in KE ($d = -0.33$) and KF ($d = -0.23$) asymmetry compared to the PLC condition. Small-to-moderate associations were observed between change in KE asymmetry and change in T25FW ($p = -0.43$) and 2MW ($p = -0.24$). A moderate association was observed between change in KF asymmetry and change in T25FW ($p = -0.31$), while no association was observed with change in 2MW ($p = -0.07$). **Conclusions:** FES cycling may be efficacious for reducing lower limb strength asymmetry and improving mobility in PwMS who have mobility impairment. These preliminary results will inform future FES cycling investigations with larger sample sizes.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS, Comprehensive care and MS, Functional electrical stimulation

ARE YOU ON TWITTER? IJMCS IS!



We frequently tweet about new IJMCS developments, and we don't want you to be left out. Readers and reviewers: Keep current with IJMCS announcements and the latest free online content. Authors: We're tweeting about your articles; don't miss the opportunity to help spread the word about your research.

Find and follow us @IJMCStweets

POSTERS

COMPLEMENTARY AND ALTERNATIVE THERAPIES

(CAM01) Multiple Sclerosis Imbalance: Visual Rehabilitation

Marcia Baptista Dias,¹ Alice Estevo Dias²

¹Rehabilitation and ²Scientific Research, Brazilian Association of Multiple Sclerosis, Sao Paulo, Brazil

Background: Imbalance is among the most debilitating symptoms in multiple sclerosis (MS), causing falls and reflecting, to a large extent, the dysfunctional integration of visual sensory signals. **Objectives:** This preliminary study aimed to show the effects of visual rehabilitation on balance in a small group of people with MS. **Methods:** Three people with MS presented signs and symptoms of body imbalance. All were evaluated before and after visual rehabilitation by a specialized optometrist, from ocular motility, cover test, and stereoscopy to chromatic and pupil analysis. Rehabilitation consisted of 7 sessions involving balance exercises associated with vision. **Results:** In the initial evaluations, participants presented the same pattern of body imbalance. After visual rehabilitation, improvements in body posture, static and dynamic balance, and overall physical performance were observed in all participants. **Conclusions:** The data obtained revealed that visual function contributes positively to the postural control system and suggests that visual rehabilitation may be an advantageous option for the treatment of imbalance in MS; it involves ocular exercises capable of producing physical and mental stimuli that, as they improve vision, make it possible to decrease the rates of falls and consequent impairment of functional capacity.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS

(CAM02) Acupuncture and Electromagnetotherapy for Chronic Pain Relief in Multiple Sclerosis

Alice Estevo Dias,¹ Teresa Cristina Vieira²

¹Scientific Research and ²Rehabilitation, Brazilian Association of Multiple Sclerosis, Sao Paulo, Brazil

Background: Chronic pain is common in people with multiple sclerosis (PwMS) with approximately 42% to 90% experiencing pain at some stage of the disease course. Pharmacologic treatment in multiple sclerosis (MS)-related pain is usually unsatisfactory and often has severe side effects, and therefore, a need for alternative methods of pain relief is critical. **Objectives:** To evaluate the effectiveness and analgesic efficiency of acupuncture associated with electromagnetotherapy for chronic pain relief in a PwMS group. **Methods:** A total of 12 patients with MS were included in this prospective study: 10 women and 2 men, aged between 40 and 74. Mean Expanded Disability Status Scale score was 4.8; 42% of patients were classified as having relapsing-remitting MS, 33% as secondary-progressive MS, and 25% as primary-progressive MS. All reported pain (10 = back, 2 = legs/feet), used pharmacologic treatment, underwent 15 acupuncture sessions and electromagnetic therapeutic equipment applications (Kenkobie), and answered a structured pain questionnaire. **Results:** The primary end point was reduction in pain intensity or elimination, while the secondary end point improved symptoms and quality of life. This preliminary study revealed that MS-related pain can have a significant impact on health, activity, and participation of people, drastically reducing the quality of life. **Conclusions:** Although our overall results suggest that these nonpharmacologic interventions had beneficial effects on chronic pain and were not harmful, studies with robust methodology are needed to assess safety and possible long-term effects, justifying the use of these interventions on chronic pain in MS.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS

(CAM03) The Effects of Reflexology in People with Multiple Sclerosis

Alice Estevo Dias,¹ Evanda Soares Oliveira²

¹Scientific Research and ²Rehabilitation, Brazilian Association of Multiple Sclerosis, Sao Paulo, Brazil

Background: Multiple sclerosis (MS) is associated with a wide variety of different physical and psychological symptoms that have a profound effect on quality of life. Complementary and alternative medicine (CAM)

is a current treatment that seems effective in relieving symptoms of patients with MS. **Objectives:** To show the opinion of a group of people with MS about the effects of reflexology. **Methods:** This study involved 12 people with MS and healthy feet without injury, damage, thrombosis, inflammation, lesion, or fractures, 7 (58%) women and 5 (42%) men, aged 25 to 72 years, and mean Expanded Disability Status Scale score 4.5. In the group reflexology interventions were performed within 10 weeks, once a week for 45 minutes. Data were collected through a structured questionnaire, immediately after. **Results:** All expressed satisfaction with the interventions, among them, 7 (58%) reported reflexology benefits in both psychological symptoms and pain, and 5 (42%) in exclusively psychological. **Conclusions:** The results showed that, according to participant opinion, reflexology in relieving anxiety, stress, depression, and pain was effective. Therefore, this method, as an efficient technique, can be recommended for people with MS. However, sufficient scientific evidence should support its effectiveness and safety.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS

(CAM04) The Effects of CBD:THC Tincture Oil in Reducing Symptoms and Overall Symptom Management Medication Dosages, in Persons with Multiple Sclerosis

Aryn Sieber,¹ Kristine Werner,² Karen Carera,³ Ben Thrower,⁴ Jacqueline Rosenthal⁴

¹CannaCauses Foundation, Los Angeles, CA; ²Smyrna, GA; ³Brookhaven, GA; ⁴Andrew C. Carlos MS Institute, Shepherd Center, Atlanta, GA

Background: It is now becoming more common for persons with multiple sclerosis (PwMS) to use cannabis to try to alleviate their multiple sclerosis (MS) symptoms. A survey of PwMS published in 2017 found that 47% of respondents considered using cannabis to treat their MS symptoms, 26% used cannabis for their MS symptoms, 20% have spoken with their physician about using cannabis, and 16% currently use cannabis (Cofield et al). Many reviews (Zhornitsky and Poivin, 2012; Jawahar et al, 2013; Koppel et al, 2014; Whiting et al, 2015) agree cannabis might have a positive effect on pain in MS. In addition to the legal status, limited research evidence remains a barrier to understanding the role cannabis can play in PwMS to alleviate symptoms. The amount of scientific research in this area is increasing; however, case reports and anecdotes exceed studies. Thus, data regarding cannabis use to treat pain, spasticity, neuropathy, and sleep quality in PwMS remain limited. **Objectives:** The purpose of this study is to investigate if medicinal cannabis CBD:THC oil tinctures 1) improve symptoms and 2) reduce overall symptom management medication dosages in PwMS. We hypothesize that PwMS will have improvements in these measures while using CBD:THC tinctures. **Methods:** Participants took CBD:THC tincture oil daily. Self-reported symptom and medication assessments rating a scale from 1 to 10 were completed at baseline prior to starting a tincture, and again after an average duration of 3-4 months. **Results:** Sixty-one PwMS aged 25 and older are included in the study. There were significant reductions ($P < .0001$) in the following symptom management scores: pain (from a mean [SD] of 7.4 [2.0] to 3.9 [1.9], $n = 45$), spasticity (from a mean [SD] of 7.2 [1.9] to 3.3 [1.9], $n = 31$), neuropathy (from a mean [SD] of 7.7 [2.0] to 4.5 [2.6], $n = 25$), and sleep (from a mean [SD] of 7.5 [1.9] to 3.0 [2.1], $n = 34$). Gabapentin intake was significantly reduced from a mean [SD] of 1581.3 [1284.6] mg to 625 [739.9] mg ($n = 12$; $P = .036$). There were no significant reductions in baclofen, tizanidine, or benzodiazepine intake. **Conclusions:** Although medicinal cannabis CBD:THC tincture oil shows promise in overall symptom reduction and symptom management medication dosage reduction in PwMS, researchers need to conduct additional studies, including clinical research studies, for PwMS using medicinal cannabis CBD:THC tincture oil. A larger sample size will allow inferential statistics to be performed. This study will further contribute to the evidence related to the efficacy of this intervention.

Supported by: None

Disclosure: Aryn Sieber: CannaCauses Foundation (consulting fee). Kristine Werner, Karen Carera, Ben Thrower, Jacqueline Rosenthal: Nothing to disclose.

Keywords: Complementary/alternative therapies in MS

(CAM05) Challenges and Opportunities in Progressive Multiple Sclerosis Trials: Lessons from Lipoid Acid

Carin S. Waslo,^{1,2} M. Mateo Paz Soldan,^{3,4} Mark S. Freedman,⁵ Pavle Repovic,⁶ Andrew J. Solomon,⁷ John R. Rinker,⁸ Mitchell T. Wallin,⁹ Jodie K. Haselkorn,¹⁰ Rebecca I. Spain¹¹

¹Neurology, VA Portland Health Care System, Portland, OR; ²Neurology, Oregon Health & Science University, Portland, OR; ³Neurology, University of Utah, Salt Lake City, UT; ⁴Neurology, VA Salt Lake City HCS, Salt Lake City, UT; ⁵University of Ottawa and the Ottawa Hospital Research Institute, Ottawa, ON, Canada; ⁶Multiple Sclerosis Center,

Swedish Neuroscience Institute, Seattle, WA; ⁷Neurological Sciences, University of Vermont College of Medicine, Burlington, VT; ⁸Neurology, University of Alabama at Birmingham, Birmingham, AL; ⁹Washington, DC VAMC, MS Center of Excellence, Washington, DC; ¹⁰Department of Epidemiology, University of Washington, Seattle, WA; ¹¹Oregon Health & Science University, Portland, OR

Background: Recruitment for progressive multiple sclerosis (PMS) randomized clinical trials (RCTs) is challenging due to 1) lack of standard definitions of PMS, 2) varying duration of progression prior to entry, 3) restricted age and disability requirements, 4) lack of specific outcome measures with associated sample sizes, 5) poorly understood PMS natural history, 6) integration of prior and/or current treatments with disease-modifying therapies (DMTs), 7) integration of emerging DMTs specific for PMS, and 8) over-the-counter availability of some investigational agents (eg, lipoic acid [LA], biotin). The resulting narrow inclusion criteria makes recruitment for clinical trials difficult, and raises concerns regarding generalizability of the results. **Objectives:** To describe the varying successes of recruitment approaches used in a current multisite RCT for PMS, compare the resulting study population to other PMS trial populations, and consider if the population is reflective of the general PMS population. **Methods:** Describe recruitment methods for Lipoic Acid for Treatment of Progressive MS RCT (LAPMS; trial registration: NCT03161028). Present the demographic characteristics of the subjects and compare to other PMS study populations and to a general reference PMS population. **Results:** LAPMS is a 2-year, multisite, phase 2, double-blind RCT of LA in PMS (n = 118, currently 71% randomized). Enrollment criteria: age ≥ 18 years, non-relapse-related MS clinical worsening in prior 2 years, worsening ≥ 1 year if on concurrent DMT, Expanded Disability Status Scale (EDSS) score 3.0-6.5, and without non-MS-related ambulation deficits. Recruitment sources are direct from clinic (63.1%), physician referral (23.8%), community outreach/MS events (3.6%), local repository (3.6%), Tiny URL promotion/National MS Society website (2.4%), Facebook advertisement (2.4%), and a North American Research Committee on Multiple Sclerosis (NARCOMS) recruitment letter (1.2%). Participants have mean age 59.4 (8.1) years, 36 (49.3%) females, median EDSS score of 6.0 (3.0-6.5), mean 23.0 (11.7) years since symptom onset, and 9 (12.3%) subjects without previous DMT. Mean LAPMS population age and disease duration are higher than those of study populations in the EXPAND SPMS trial and ORATORIO PPMS trial. While similar to the EXPAND trial, LAPMS has higher EDSS levels than the ORATORIO population. Updates and comparison to a general PMS population will be presented. **Conclusions:** Restrictive PMS recruitment criteria create differences in PMS trial populations as well as raise concerns about generalizability of results to wider PMS populations. Approaches to addressing these issues are discussed.

Supported by: None

Disclosure: *Carin S. Waslo, Mark S. Freedman, Mitchell T. Wallin, Jodie K. Haselkorn:* Nothing to disclose. *M. Mateo Paz Soldan:* Biogen (research funding); Novartis (contracted research). *Pavle Repovic:* Alexion (consulting fee, speakers' bureau); Biogen, EMD Serono (consulting fee, contracted research, speakers' bureau); Celgene, Novartis, Sanofi Genzyme, Viela (consulting fee); Genentech (contracted research, speakers' bureau); Genzyme (speakers' bureau). *Andrew J. Solomon:* Actelion, Genentech, Novartis (contracted research); Alexion, Celgene, EMD Serono (consulting fee); Biogen (consulting fee, research funding). *John R. Rinker:* Biogen, EMD Serono (contracted research). *Rebecca I. Spain:* TG Therapeutics (consulting fee).

Keywords: Clinical trials, Complementary/alternative therapies in MS, Disease-modifying treatments in MS

(CAM06) Exercise in Medicine: A Complementary Exercise Promotion Approach Within Comprehensive Multiple Sclerosis Care

Emma V. Richardson,¹ Elizabeth Barstow,² Matthew Fifolt,³ Robert W. Motl¹

¹Department of Physical Therapy, ²Department of Occupational Therapy, and ³Department of Health Care Organization & Policy, University of Alabama at Birmingham, Birmingham, AL

Background: Exercise is one of the only complimentary strategies that improves symptoms of multiple sclerosis (MS) and slows disease progression and functional manifestations. There is further evidence that engaging in exercise regenerates neuroplasticity within the central nervous system. Such evidence supports exercise as a strategy for optimizing independence and quality of life among persons with MS. However, only 20% of the MS population engage in sufficient levels of exercise for accrual of benefits. **Objectives:** The objective of this work was to develop a new, revolutionary, systematic approach for promoting exercise within comprehensive MS care. This system would reduce the burden of exercise promotion on neurologists while allowing patients to receive, timely, accurate information about exercise, and ongoing opportunities

and support to engage in this activity. **Methods:** Over a 6-year period, we conducted interviews with more than 80 persons with MS and more than 70 health care providers to help design, refine, and finalize a new, comprehensive approach to systematically integrate exercise promotion within comprehensive MS care. We also conducted a quality improvement approach to translate the conceptual ideas into a practical context and develop tangible tools to be used to deliver "Exercise in Medicine." **Results:** Through this rigorous line of research, we have built a new, patient-informed, systematic process that integrates exercise promotion within comprehensive MS care: so-called Exercise in Medicine. Persons with MS and health care providers are supportive of this endeavor and continue to be part of its refinement and improvement. **Conclusions:** Exercise in Medicine has the potential to revolutionize the promotion of exercise as a complementary strategy to improve quality of life among persons with MS. This systematic process is now at a stage where it is ready to be tested through clinical trials.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS, Comprehensive care and MS

(CAM07) Changes in Dietary Habits of Individuals Living with Multiple Sclerosis Enrolled in a Day Wellness Program

Zoe Edwards,¹ Brian Hutchinson,² Tiffany Malone¹

¹Multiple Sclerosis Achievement Center, Dignity Health, Citrus Heights, CA; ²Multiple Sclerosis Achievement Center, Dignity Health, Sacramento, CA

Background: Nutrition education for individuals who attend the Multiple Sclerosis Achievement Center (MSAC) focuses on overall health and prevention or management of comorbidities. While there has not been a multiple sclerosis (MS)-specific dietary pattern proven to reduce MS symptoms for all individuals, nutritional choices can affect management strategies. Members of the MSAC attend a weekly program that addresses physical, cognitive, and social well-being. On a monthly basis, nutrition education is provided to discuss diverse aspects of dietary habits and target strategies related to the challenges of living with MS. Some program participants receive additional nutrition education through individualized consultations or small group discussions. **Objectives:** To examine the dietary changes of people with MS who participate in a day wellness program over an 18-month period and determine if changes in habits occur with general nutrition education during the day program. In addition, a comparison in dietary habits between general and individualized nutrition education will be examined. **Methods:** Fifty-two MSAC members have completed Rate Your Plate (RYP), a self-reported food-frequency questionnaire, every 6 months to monitor changes in dietary choices over a year. RYP consists of 27 questions focusing on typical dietary choices within specified categories. The answer to each question is assigned a score of 1, 2, or 3 points with composite scores ranging from 27-81; higher points indicate healthier choices. Between questionnaire administration, members have received monthly nutrition education, with 16 members receiving additional small-group nutrition education. Analysis examining 18-month results is planned. **Results:** Three of four intended rounds of data collection have occurred, with the fourth scheduled in February 2020. The mean (SD) baseline RYP score was 58.48 (8.5). The mean (SD) of the scores at 6 months increased, but not significantly, to 62.33 (7.7) ($P = 1.00$). Increases at 1 year were statistically significant, 63.71 (7.5) ($P < .001$) compared to baseline. Paired t tests were used to identify statistical significance. The category showing the greatest improvement (10%) over the first year was the sugar content of desserts. Another improvement (9%) is the frequency in which members are eating meals out. **Conclusions:** Complete collection and analysis of changes between 4 data collection points will be finalized for presentation of the poster.

Supported by: None

Disclosure: *Zoe Edwards, Tiffany Malone:* Nothing to disclose. *Brian Hutchinson:* Biogen (speakers' bureau).

Keywords: Complementary/alternative therapies in MS, Dietary changes

CASE REPORTS/CASE SERIES

(CRS01) Seasonal Variation and Other Observations in Myelin Oligodendrocyte Glycoprotein (MOG) Antibody-Associated Disease

Allison N. Block, Ahmed Z. Obeidat

Neurology, Medical College of Wisconsin, Milwaukee, WI

Background: Myelin oligodendrocyte glycoprotein (MOG) antibody-associated disorders can mimic multiple sclerosis (MS). MOG antibody seropositivity has been found in subsets of patients with demyelinating disease, especially those with optic neuritis and transverse myelitis. Improved MOG antibody-associated disease understanding is necessary to improve diagnosis timing and provide proper treatment. **Objectives:** To evaluate the characteristics of patients with a confirmed diagnosis of MOG antibody-associated disease. **Methods:** We report on the clinical and imaging characteristics of all patients who presented to our practice with optic neuritis or transverse myelitis, and tested positive for MOG antibody between August 2018 and January 2020. **Results:** We identified 11 patients, of which 9 (82%) were female. Ten (91%) patients presented between September and December. Optic neuritis preceded by a prodromal headache was the presenting symptom in all patients. Sixteen optic nerves in the 11 patients were symptomatic. Of the 16 symptomatic optic nerves, 12 (75%) nerves showed increased signal on diffusion-weighted imaging. Incomplete clinical recovery was observed in 11 (69%) optic nerves and ranged from no light perception to a mild decrease in visual acuity. Spinal cord lesions were present in 4 patients (36%). Poorly demarcated white matter brain T2 signal abnormalities were present in 7 patients (64%). Finally, patients younger than 40 years tended to have higher titers with 5/6 (83%) patients in this age group having a titer of 1:1000. No patients 40 years and older had a titer of 1:1000. All patients received treatment with prednisone or steroid-sparing agents (rituximab or mycophenolate). None of the treated patients relapsed over the duration of the study. One patient was initially misdiagnosed as having MS and was treated with several disease-modifying agents; he continued to relapse (with sustained disability) until the correct diagnosis was confirmed. **Conclusions:** Our findings suggest seasonal variation of MOG antibody-associated disease, with peak clinical presentation during fall and winter months. This may be due to the peak of respiratory viral infections in the fall and winter as preceding infections were reported in association with MOG antibody-associated disease. Increased diffusion-weighted imaging signal of optic nerves may provide insight into the mechanism of optic nerve damage. Incomplete recovery of optic neuritis is common but often mild and rarely resulted in blindness. The significance of higher antibody titer in younger individuals requires further investigation. Our observations contribute to the growing knowledge of MOG antibody-associated disease, a mimicker of MS. Our study was limited by a small sample size.

Supported by: None

Disclosure: *Allison N. Block: Nothing to disclose. Ahmed Z. Obeidat: Alexion (speakers' bureau); Alexion Pharmaceutical, Celgene, EMD Serono, Genentech, Sanofi Genzyme (consulting fee); Biogen, Novartis (consulting fee, speakers' bureau); International Journal of MS Care (editorial board).*

Keywords: *Imaging and MS, Mimickers of MS, Natural history of MS*

(CR502) Multiple Surgeries and Misdiagnosis Before Multiple Sclerosis Diagnosis: A Case Report

Angel Lopez,¹ Joanie Figueroa-Amaro,² Astrid Diaz,¹ Viviana Martinez,¹ Ivonne Vicente,¹ Angel Chinae¹

¹San Juan Multiple Sclerosis Center, Guaynabo, PR; ²Medicine, San Juan Bautista School of Medicine, Caguas, PR

Background: A delay in diagnosis of multiple sclerosis (MS) can increase the amount of disability in patients. **Objectives:** Not applicable. **Methods:** Case report. **Case:** A 50-year-old Puerto Rican man reports onset of symptoms in 1994 with a severe headache that lasted 6 months. A year later the patient fell and afterwards started having gait difficulties, pain, and bowel and bladder problems. After computed tomography was done, he was diagnosed with herniated discs, and treated with physical, massage, and chiropractic therapies. Afterwards, he had frequent falls and facial palsy for 5 years. In these years he had several surgeries performed due to the pain and symptoms. Patient continued with his symptoms after surgeries. Psychiatrist evaluated and in 2017 performed brain and cervical magnetic resonance imaging and referred to neurologist. Patient was diagnosed with MS in 2018 and assigned an Expanded Disability Status Scale score of 6.0. The time from first symptom to diagnosis (24 years) and the amount of surgeries could have influenced the disability that the patient presents. **Results:** Not applicable. **Conclusions:** This case brings attention to the importance of earlier diagnosis and treatment of MS to reduce disability and improve quality of life in patients. Also, it highlights that we still need to increase education to all health care professionals in Puerto Rico about symptoms, tests, and diagnostic criteria for MS.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Comprehensive care and MS, Natural history of MS*

(CR503) Head Trauma as Onset for Multiple Sclerosis Diagnosis: A Case Report

Astrid Diaz, Viviana Martinez, Ivonne Vicente, Cristina M. Rubi, Angel Lopez, Angel Chinae

San Juan Multiple Sclerosis Center, Guaynabo, PR

Background: Recent studies found an increased risk of developing multiple sclerosis (MS) if head trauma was experienced during teenage years, even though the relationship of head injury and risk of developing MS is not clear among the adult population. Previous research in Puerto Rican patients with MS shows that 38.2% reported some type of trauma. **Objectives:** We aim to report a case of head injury and subsequent diagnosis of MS. **Methods:** Case report. **Case:** A 47-year-old Puerto Rican woman with no previous history of comorbidities. The patient was walking, fell, and had head trauma. Concerned, she went to general medicine physician and computed tomography was performed. Months later, the patient started presenting pain in right leg, but was diagnosed with urinary tract infection and treated. After this, patient developed episodes of nausea and high levels of leukocytes. In May 2019, she developed blurry vision that ended with vision loss in right eye. Ophthalmologist evaluated and referred to neuro-ophthalmologist, who ordered magnetic resonance imaging (MRI). She later developed loss of balance and increased falls and was subsequently diagnosed with MS in June 2019. **Results:** Not applicable. **Conclusions:** This case highlights the importance of creating awareness about patients who had head trauma and present neurologic symptoms. Performing tests like MRI as soon as symptoms start could help to achieve an early diagnosis and help reduce disability among patients with MS.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Comprehensive care and MS*

(CR504) Team Approach Yields Surprising Functional Progress and Quality-of-Life Changes in a Challenging Case of Neuromyelitis Optica

Clare T. Hartigan,¹ Christopher Wells²

¹Multiple Sclerosis Outpatient Rehabilitation, Shepherd Center, Atlanta, GA; ²Multiple Sclerosis Wellness Program, Shepherd Center, Atlanta, GA

Background: A 59-year-old female (DR) was diagnosed with neuromyelitis optica (NMO) in February 2016. She developed sudden onset of paralysis and had no voluntary movement from her neck down. DR spent extensive time in a nursing facility and eventually transferred home on a stretcher. At the time of her diagnosis, she was 5'5" tall with body mass index (BMI) of 49.08. A rehabilitation team approach from May 2017 through February 2020 led to surprising functional progress and improved quality of life. **Objectives:** In May 2017 DR was referred to a MS Wellness Program. All mobility was dependent to include Hoyer transfers, sitting balance, bed mobility, and bowel/bladder management in bed. She also had significant neurologic muscle weakness (2-/5 throughout), spasticity, joint contractures, and chronic pain from arthritis in her knees, back, and shoulders. **Methods:** DR attended MS Wellness appointments via stretcher, until she received a power wheelchair. Exercise specialist intervention was 2x/week and included pool program, dryland exercises for stretching, strengthening, and sitting balance. One year later (May 2018) she began slide board transfers. By 2019 transfers were via depression and she started using a hydraulic standing device during exercise specialist sessions. In October 2019 DR was referred to outpatient physical therapy (PT). Her strength was grossly 3-/5, she had contractures at bilateral hips/knees/ankles, and she was unable to stand or walk at home. Her BMI was 43.27. DR attended PT 2-3x/week for 30 sessions to include standing frame and soft tissue work for contractures, transfer training with 3-in-1 bedside commode, progressive gait training, and custom home program. **Results:** In November 2019, DR achieved supervision for transfers on/off bedside commode and began gait training in parallel bars (PBars) with bilateral MalleoLoc braces and heel wedges. Sit/stand and stepping in PBars progressed from Max A to Supervision. By mid-December 2019 she required Mod A to walk outside the PBars with front-wheeled bariatric walker (FWBW). At end of December 2019 she required Min A to walk up to 76 ft for exercise. In early February 2020 patient was able to walk 135 ft at a time with close stand-by assist with FWBW and bilateral solid ankle foot orthoses. She walked over indoor and outdoor terrains including 5-degree ramps. After caregiver training, she became safe to stand and walk at home for exercise. **Conclusions:** After being completely paralyzed after NMO diagnoses in 2016, DR spent nearly 3 years attending a MS Wellness Program and then eventually outpatient PT. She progressed from being totally depen-

dent to requiring supervision for all transfers. She no longer had to use bed pans. Additionally, she learned to walk with stand-by assist indoor/outdoor terrains with FWBW and bilateral ankle foot orthoses. DR's functional gains and improved quality of life would have been very unlikely without a team approach. DR is to be commended for her determination, and as health care professionals we should always keep an open mind.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS, Comprehensive care and MS, Wellness program

(CR505) Differential Diagnosis and Treatment of Tumefactive Demyelination in a Teenaged Girl

Gillian R. Paton, Sarah Hopkins

Division of Neurology, Multiple Sclerosis and Neuroinflammatory Disorders, Children's Hospital of Philadelphia, Philadelphia, PA

Background: Tumefactive demyelination in pediatric patients is rare and associated with high morbidity and mortality. The differential diagnosis includes tumor, abscess, acute hemorrhagic leukoencephalitis, acute disseminated encephalomyelitis, and tumefactive multiple sclerosis (MS). Awareness of the differential and early treatment is essential as this presentation may be associated with death or severe morbidity within days.

Objectives: Review a case of tumefactive demyelination in a teenaged girl, and review the pertinent literature, including differential diagnosis, key clinical characteristics, and treatment options. **Methods:** Medical record review and review of the literature. **Results:** A 16-year-old girl, previously healthy and developmentally normal, presented with encephalopathy progressing quickly to left hemiparesis and global aphasia in the setting of recent upper respiratory infection with cough, headache, and otalgia. Head computed tomography demonstrated vasogenic edema of the left temporal and parietal lobes and 4 mm of midline shift to the right, without obvious underlying mass. Magnetic resonance imaging (MRI) of the brain revealed extensive confluent and expansile-appearing white matter signal abnormality of the left brainstem, internal capsule, and parietal and temporal lobes with associated microhemorrhages and left cerebral peduncle and pontine enhancement, suggestive of severe demyelinating disease, and her cerebrospinal fluid showed a neutrophilic pleocytosis with no identifiable pathogen. A brain biopsy showed no signs of neoplasm. She was promptly treated with high-dose prednisolone, and the course continued for 7 days in conjunction with 9 rounds of plasmapheresis. Cyclophosphamide was initiated with a plan for a 6-month course. Follow-up MRI of the brain 1 month after presentation demonstrated decreased extent of T2/fluid attenuation inversion recovery abnormality and development of cystic encephalomalacia along the left anterior temporal lobe with resolution of midline shift. She was transferred to the inpatient rehabilitation service for a 1 month stay. At discharge she had persistent right-sided weakness with spasticity but was able to ambulate using a single crutch, and had shown improvement in expressive and receptive aphasia. **Conclusions:** Tumefactive demyelination requires rapid evaluation and treatment. The differential diagnosis includes acute hemorrhagic leukoencephalitis, which has a particularly high fatality rate. Early and judicious immunomodulatory treatment in these cases is lifesaving.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Disease-modifying treatments in MS, Imaging and MS, Immunology and MS

(CR506) A Long-Standing Case of Recurrent Transverse Myelitis Due to Myelin Oligodendrocyte Glycoprotein (MOG)-IgG Antibody Mimicking Multiple Sclerosis

Gina S. Perez, Peggy J. Wisdom, Nidhiben Anadani

Neurology, University of Oklahoma Health Sciences Center, Oklahoma City, OK

Background: Myelin oligodendrocyte glycoprotein (MOG) IgG antibody causes central nervous system demyelination and mimics multiple sclerosis (MS). Patients with MOG-IgG typically present with recurrent or monophasic optic neuritis, transverse myelitis, conus medullaris lesion, brainstem encephalitis, steroid-dependent symptoms, and acute disseminated encephalomyelitis. Lack of testing for MOG-IgG in these patients can lead to an incorrect diagnosis of MS, and treatment with certain medications that can worsen MOG-IgG-associated disease. **Objectives:** To report a case of a patient with MOG-IgG antibody-mediated recurrent transverse myelitis who was diagnosed with possible MS for more than 15 years. **Methods:** A 66-year-old woman with a known diagnosis of possible MS presented to our clinic for follow-up. Her history dated back

to 1989 when she had an episode of extreme fatigue, gait imbalance, and numbness in her hands and feet that resolved spontaneously. She had recurrent episodes of symptomatic myelitis in 1997 and 2001. On examination, she had spastic weakness of bilateral iliopsoas and finger extensors (left worse than right), brisk reflexes on the left hemibody with left ankle clonus, relative sensory level at T12, and spastic ataxic gait without assistance. Timed 25-foot walk was 6.86 seconds. **Results:** In 2001, magnetic resonance imaging (MRI) of the spinal cord showed several short segment lesions in the cervical and thoracic spinal cord including the conus medullaris; MRI of the brain was normal. Cerebrospinal fluid (CSF) showed elevated protein, IgG synthesis rate, and CSF-specific oligoclonal bands. Aquaporin 4 antibody (AQP-4) in CSF was negative. She was diagnosed with possible MS given the lack of better explanation and was started on a combination of interferon beta-1a and mycophenolate. Since 2001, she did not have clinical relapse though had new nonspecific lesions in her brain and persistently enhancing short segment lesions in the cervical and thoracic spinal cord in 2014 and 2016. Interferon beta-1a was stopped in 2017 and mycophenolate was continued. Serum MOG-IgG was tested and was positive in February 2019. **Conclusions:** MOG-IgG-mediated disease and MS show a relevant phenotypic, clinical, and radiologic overlap that can potentially lead to misdiagnosis and worse patient outcomes, as some medications used for the treatment of MS might be ineffective or even harmful in MOG-IgG-associated disease. Before the diagnosis of MS, testing for MOG-IgG antibody should be considered in selected patients.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, MOG IgG antibody

(CR507) Case Report of Severe Multiple Sclerosis Relapse Due to B-Cell Reconstitution Post Alemtuzumab

Jennifer Chester,¹ Tyler Kaplan²

¹Kansas City Multiple Sclerosis Center, College Park Specialty Center, Overland Park, KS;

²Neuro-immunology, Rush University Medical Center and University of Utah, Chicago, IL

Background: Alemtuzumab is a pan-lymphocyte-depleting anti-CD52 antibody used in the treatment of multiple sclerosis (MS). However, there have been reports of severely exacerbated central nervous system inflammation following alemtuzumab infusion. Relapse often occurs with the repletion of B cells months after treatment, whereas T cells can take up to 3 years to replenish. B-cell reconstitution occurs when the memory B cells replenish more rapidly than the regulatory T cells. This has previously been seen with the use of rituximab as a B-cell-depleting therapy in disorders such as rheumatoid arthritis. **Objectives:** Here, we present a case of a 52-year-old white woman diagnosed with MS in 2000. She had previously tried multiple disease-modifying therapies including interferon beta-1a, interferon beta-1b, glatiramer, natalizumab, and dimethyl fumarate. She was given round 1 of alemtuzumab in December 2018. In May 2019 she had an exacerbation that caused hospitalization for intravenous methylprednisolone and physical therapy. She made a full recovery to baseline. In July 2019 she had another exacerbation, presenting to the clinic with multiple new symptoms including ataxia, urinary incontinence, weakness, and numbness. She was treated with repository corticotropin injection with no improvement. This was immediately followed by another hospitalization for plasma exchange. Her symptoms continued to progress rapidly. Brain magnetic resonance imaging (MRI) showed development of innumerable enhancing lesions throughout the bilateral cerebral hemispheres and right lateral pons. Cervical spine MRI showed a new 5-mm enhancing lesion. MRI of thoracic spine showed several enhancing thoracic cord lesions consistent with active demyelinating disease. Due to her rapid deterioration, the decision was made to transfer her to another state to receive a higher level of care. After being evaluated it was determined that she likely had active demyelination related to B-cell reconstitution. She was again hospitalized and received 1 dose of rituximab. Unfortunately, as of October 2019, the patient has continued to decline. **Methods:** NA. **Results:** She was discharged home on hospice and passed away a week later. An autopsy was performed and the results are pending at this time. **Conclusions:** This case signifies the importance of strict monitoring of B cells after alemtuzumab due to the risk of relapse as the B-cell population returns. Further analysis is needed for optimal care of these patients.

Supported by: None

Disclosure: Jennifer Chester: Allergan, Biogen, Novartis, Sanofi Genzyme (speakers' bureau). Tyler Kaplan: Nothing to disclose.

Keywords: Disease-modifying treatments in MS, Immunology and MS, Nursing management in MS

(CRS08) Demographics, Clinical Characteristics, and Outcomes of Myelin Oligodendrocyte Glycoprotein (MOG) Antibody Disease Followed Up at Washington University in St. Louis

John R. Ciotti, Anne Cross, Salim Chahin

Washington University in St. Louis, St. Louis, MO

Background: Antibodies to myelin oligodendrocyte glycoprotein (MOG) have been associated with central nervous system demyelination that is distinct from other neuroinflammatory conditions. Our knowledge of the clinical spectrum of MOG antibody disease (MOGAD) is evolving, without clear understanding of prognosis and best treatments. **Objectives:** To report demographics, clinical characteristics, and treatment responses of our MOGAD cohort. **Methods:** Patients at Washington University were identified via diagnosis code and confirmed to have at least 1 positive MOG antibody test. Demographic, clinical course, estimated disability (extracted using published tools), laboratory, and treatment data were collected after institutional review board approval. **Results:** 24 patients with MOGAD were included. They were 75% female and 92% Caucasian with mean onset age of 43.5 (range 16.8-76.0) years. Average duration of follow-up was 4.0 (range 0.4-18.0) years. Initial symptoms exclusively included optic neuritis (17/24, 41% bilateral) or transverse myelitis (9/24), with 2/24 having both occur simultaneously. 12.5% (3/24) had concurrent involvement of other areas (brainstem, cerebral). A total of 42 attacks (including initial onset) were adjudicated; the annualized relapse rate was 0.46. 42% have had only a single attack. Attacks tended to be severe (estimated Δ Expanded Disability Status Scale [EDSS] score +3.3), followed by complete recovery in 35% and no recovery in 21%. Most (81%) attacks were treated with corticosteroids. 38% of all patients have remained stable without long-term treatment, but 79% of those with relapses were on no long-term treatment. Rituximab was associated with a low relapse rate, although breakthrough relapses still occurred in 2 patients. Four patients had 3+ attacks: all had optic neuritis, and 3 were exquisitely sensitive to decreases in their maintenance corticosteroid dose. 54% of patients had multiple MOG antibody titers (average 7.2 months later), of whom 31% became seronegative. None of these patients relapsed after testing negative. Cerebrospinal fluid (CSF) test results were modestly abnormal (median cell count 8, average protein 56.9). No patients had >2 CSF-specific oligoclonal bands. In those with 2+ years of follow-up, 50% remained relapse-free. In those with 5+ years of follow-up, 44% remained relapse-free. Average current EDSS score is 1.9. 58% have an EDSS score \geq 2.0, and 13% have an EDSS score \geq 4.0. **Conclusions:** We report a cohort of Midwestern patients with anti-MOG disease. Our results are largely consistent with those reported in other cohorts of this disease.

Supported by: None

Disclosure: *John R. Ciotti:* Nothing to disclose. *Anne Cross:* Biogen, Celgene, Novartis, TG Therapeutics (consulting fee); EMD Serono, Genentech/Roche (grant and personal fees). *Salim Chahin:* Biogen, Genentech, Novartis, Sanofi Genzyme, Teva Neuroscience (personal fees).

Keywords: MOG antibody disease

(CRS09) A Fatal Case of Alemtuzumab-Induced Immune Thrombocytopenic Purpura in a Patient with Relapsing Multiple Sclerosis

Leila Saadatpour, Rebecca Romero

Neurology, University of Texas Health Science Center at San Antonio, San Antonio, TX

Background: Alemtuzumab is a humanized monoclonal antibody against CD52 approved for relapsing types of multiple sclerosis (MS). Despite being an effective medication, there have been limitations of use for this medication due to various serious adverse effects. Common adverse effects include infections, transfusion reactions, and autoimmune responses such as hemolytic anemia, thrombocytopenia, and autoimmune thyroid and renal disease. **Objectives:** To review a case of refractory immune thrombocytopenic purpura (ITP) as a result of alemtuzumab therapy in a patient who presented with altered mental status and who was found to have multiple foci of intracranial bleeding (intracerebral hemorrhage [ICH]). **Methods:** Case report. **Results:** A 39-year-old woman developed ITP after receiving 2 doses of alemtuzumab. The patient died 2 years after the initial alemtuzumab infusion after dealing with multiple comorbidities due to refractory ITP. According to the literature, ITP occurs in 2%-2.6% of patients treated with alemtuzumab. ITP can be refractory and fatal; however, in the CANMS223 study only 1 of 6 patients with ITP died due to ICH. In addition, in 2 other studies of patients with MS receiving alemtuzumab, no ITP-associated mortality is reported. **Conclusions:** Alemtuzumab is a potent immune regulator that clears up T cells and B

cells. ITP is a rare complication but can be lethal. Our patient's ITP progressed despite her compliance with every-6-month blood count checks and receiving various treatments. The aim of this presentation is to bring the potentially serious complications of alemtuzumab to attention and consider substitute therapies if feasible.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Disease-modifying treatments in MS, Immunology and MS

(CRS10) Colitis Associated with Teriflunomide

Neda Zarghami Esfahani,¹ Gloria von Geldern,¹ Meghan C. Romba,¹ Dhavan Parikh,² Annette Wundes¹

¹Neurology, University of Washington, Seattle, WA; ²Gastroenterology, Everett Clinic, Everett, WA

Background: Teriflunomide is an oral disease-modifying therapy for relapsing-remitting multiple sclerosis (RRMS) that was approved in the United States in 2012. Gastrointestinal (GI) adverse effects occurred in 15%-17% of patients in the clinical trials, and, so far, 3 cases of colitis have been reported. **Objectives:** To report new-onset Crohn disease in a patient with multiple sclerosis while on teriflunomide. **Methods:** Case report and literature review. **Results:** A 49-year-old man with RRMS started teriflunomide in January 2018 after discontinuing glatiramer acetate due to injection fatigue and dimethyl fumarate due to GI intolerance. He developed persistent diarrhea in July 2018; GI workup showed nonbleeding gastric ulcers and mild chronic gastritis without active inflammation in the small and large intestine. GI disease improved with omeprazole and stopping ibuprofen; teriflunomide was reduced to 7 mg daily in December 2018. In August 2019, he had recurrence of significant diarrhea and weight loss. Teriflunomide was stopped, and he was started on colestipol with notable improvement in his diarrhea. Repeat endoscopy showed multiple duodenal ulcers, gastric ulcers, esophagitis, and ulcerations throughout the colon and terminal ileum. Pathology demonstrated inflammatory changes consistent with inflammatory bowel disease. He was started on vedolizumab for Crohn disease in November 2019. **Conclusions:** In 2017, Health Canada released a review on teriflunomide due to postmarketing reports of colitis and concluded that, while no definite link could be established, the patients and providers should be alerted to the occurrence of rare colitis cases. As of October 2019, per Genzyme, 3 cases of colitis while on teriflunomide are reported of which 2 were considered to be related to teriflunomide. We here report a case of new onset of inflammatory colitis with endoscopic and pathologic features of Crohn disease while on teriflunomide and significant improvement after cessation of the drug, and suggest a potential causal relationship between the drug and development of colitis that warrants further investigation. Rare cases of colitis have occurred in patients on teriflunomide including a recent case of new onset of Crohn disease. While the association remains unclear, physicians should be aware of this potential adverse effect. Clinical vigilance and early treatment might be helpful in cessation of colitis progression.

Supported by: None

Disclosure: *Neda Zarghami Esfahani, Gloria von Geldern, Meghan C. Romba, Dhavan Parikh:* Nothing to disclose. *Annette Wundes:* Biogen (consulting fee); Biogen, AbbVie (contracted research).

Keywords: Colitis, Disease-modifying treatments in MS

(CRS11) Remarkable Recovery of Fulminant Multiple Sclerosis After Treatment Induction with Cyclophosphamide

Nicola Carlisle,¹ Sam I. Hooshmand,¹ Michelle Maynard,² Ahmed Z. Obeidat³

¹Neurology, Medical College of Wisconsin and Affiliated Hospitals, Milwaukee, WI;

²Neurosciences, Froedtert and the Medical College of Wisconsin, Milwaukee, WI;

³Neurology, Medical College of Wisconsin, Milwaukee, WI

Background: Fulminant multiple sclerosis (MS) is rare, and the approach to treatment beyond first-line therapies (high-dose steroids, plasma exchange, and intravenous [IV] immunoglobulins) varies. Single case reports of diagnosis and treatment of fulminant MS can provide helpful resources for the clinician. Cyclophosphamide is an alkylating agent that suppresses T- and B-cell function, interleukin 12, and T-helper type 1 (Th1) responses, thereby enhancing Th2, Th3 responses. Review of published reports suggests that cyclophosphamide has utility in early stages of MS during which inflammation predominates over degenerative processes in the central nervous system as seen by gadolinium (Gd)-enhancing lesions. **Objectives:** To describe the clinical course of a case of acute MS successfully treated with cyclophosphamide. **Methods:** Case study. **Results:** A 41-year-old woman with no medical history developed facial

paresthesia, imbalance, and dizziness. Neurologic examination showed hyperreflexia. Magnetic resonance imaging (MRI) showed numerous Gd-enhancing white matter lesions throughout the brain and cervical and thoracic spinal cord. Initial treatment with 1000 mg of IV methylprednisolone did not result in symptom improvement but decreased the burden of enhancing lesions on repeat MRI. On hospital day 8, she clinically deteriorated, and plasma exchange was initiated. She continued to worsen and developed respiratory failure requiring mechanical ventilation, cranial nerve palsies, and quadriplegia. IV immunoglobulin was given with no clinical response. A brain biopsy was obtained due to concern for lymphoma, but it showed severe demyelination with relative axonal preservation. Due to significant clinical deterioration, we proceeded with induction treatment with cyclophosphamide (600 mg/m² daily for 4 doses). Treatment resulted in gradual clinical and radiologic improvement. For maintenance therapy, she received 1 dose of IV rituximab 1000 mg and was discharged to a long-term acute care facility. Subsequently she was weaned off mechanical ventilation, significantly restored upper limb functions, and is able to walk for more than 40 ft with assistance. At day 104 from initial presentation, she continues to improve with no new relapse activity. Follow-up brain and spinal cord MRI 21 days after discharge demonstrated resolution of most of the lesions, interval improvement of some, and no new or enhancing lesions identified. **Conclusions:** Induction therapy with cyclophosphamide combined with supportive care can help improve acute fulminant demyelination.

Supported by: None

Disclosure: *Nicola Carlisle, Sam I. Hooshmand, Michelle Maynard: Nothing to disclose. Ahmed Z. Obeidat: Alexion, Biogen (consulting fee, speakers' bureau); Celgene, EMD Serono, Genentech, Sanofi (consulting fee); International Journal of MS Care (editorial board); Novartis (speakers' bureau).*

Keywords: Disease-modifying treatments in MS, Imaging and MS

(CRS12) Neurofibromatosis Type 1 and Multiple Sclerosis in the Same Patient

Sargon Bet-Shlimon, Annette Wundes, Gloria von Geldern
Neurology, University of Washington, Seattle, WA

Background: Neurofibromatosis type 1 (NF1) is an autosomal-dominant genetic disease involving primarily the skin and peripheral nervous system. Multiple sclerosis (MS) is a demyelinating disease of the central nervous system. White matter lesions on brain magnetic resonance imaging (MRI) can be seen in both diseases. Only a few cases have been reported to date describing patients with both MS and NF1. **Objectives:** To describe 3 cases of comorbid NF1 and MS to raise awareness of the possibility for the rare co-occurrence of both conditions. **Methods:** This is a detailed description of 3 cases and a literature review of co-occurring NF1 and MS. Electronic medical records, neuroimaging, and relevant ancillary tests were reviewed for all cases. **Results:** *Case 1:* A 21-year-old man with NF1 presented with an episode of several days of bilateral distal limb paresthesia and was found to have brain and spine lesions, some of which were enhancing. Optic nerves with likely optic glioma due to NF1 as well as possible prior optic neuritis. Cerebrospinal fluid (CSF) showed positive oligoclonal bands. He was started on glatiramer as disease-modifying therapy, which was advanced to natalizumab based on significant radiographic disease progression. *Case 2:* A 41-year-old woman with NF1 presented with optic neuritis (with improvement with high-dose steroids) and was found to have lesions on brain and spine MRI and positive oligoclonal bands in CSF. She was started on glatiramer acetate with clinical stability and only 1 new non-enhancing lesion on repeat MRI. Due to skin reactions on glatiramer acetate, she was changed to dimethyl fumarate. *Case 3:* A 40-year-old woman with NF1 presented with progressive gait changes. Based on MRI findings more suggestive of MS than NF1 and positive oligoclonal bands in CSF, she was diagnosed with primary progressive MS. **Conclusions:** These cases demonstrate the rare co-occurrence of NF1 and MS as well as the heterogeneity of MS presentations within the NF1 patient population. These cases also demonstrate some of the diagnostic challenges that arise when making a new diagnosis of MS in patients with NF1, including the interpretation of MRI in differentiating suspected demyelinating lesions from lesions that are associated with NF1. It is unclear if the co-occurrence of NF1 and MS is coincidental or if these cases represent an unknown relationship between the 2 diseases.

Supported by: None

Disclosure: *Annette Wundes: AbbVie, Alkermes (contracted research); Biogen (consulting fee, contracted research). Sargon Bet-Shlimon, Gloria von Geldern: Nothing to disclose.*

Keywords: Diagnosis and management of MS, Epidemiology of MS, Imaging and MS

(CRS13) Successful Use of Immunotherapy for Osmotic Demyelination Syndrome

Wijdan Rai,¹ Stephen Kolb²

¹Neurology, Ohio State University, Columbus, OH; ²Ohio State University, Columbus, OH

Background: Osmotic demyelination syndrome (ODS) is a disorder characterized by the destruction of neuronal myelin sheaths in the pons or other susceptible areas, usually due to the rapid correction of hyponatremia, often with irreversible neurologic consequences. The standard of care is supportive treatment. **Objectives:** To highlight the novel use of immunotherapeutic strategies in patients with ODS. **Methods:** Case report. **Results:** A 44-year-old man with chronic alcohol abuse and hyponatremia due to beer potomania presented with acute encephalopathy and rapidly progressive quadriplegia. Initial sodium level was 102 mm/L. Due to septic shock and concern for rhabdomyolysis, he was aggressively fluid resuscitated. Sodium levels increased by 3 points every 2 hours, ultimately normalizing within 24 hours. Brain magnetic resonance imaging showed areas of abnormal fluid attenuation inversion recovery signal hyperintensity within the pons and bilateral basal ganglia without enhancement. He was given 5 treatments of plasmapheresis followed by 5 days of intravenous immunoglobulin. Ten months later, he was ambulatory and independent in his activities of daily living. His only deficit was weakness in left wrist extension and flexion with contractures in the 4th/5th digit. **Conclusions:** Immunotherapy such as plasmapheresis, intravenous immunoglobulin, or a combination may provide a treatment approach for patients with ODS where options are extremely limited. Timing from symptom onset to treatment initiation is crucial, ideally within a week. Recovery can be slow. Patient selection may also inform outcomes as benefit seems to be in those with chronic alcohol abuse or hepatic dysfunction.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS

DISEASE-MODIFYING THERAPY

(DXT01) Maintenance of Working Status and Work Productivity in Persons with Multiple Sclerosis Treated with Dimethyl Fumarate: A 5-Year Analysis of the North American Research Committee on Multiple Sclerosis (NARCOMS) Registry

Amber Salter,¹ Samantha Lancia,¹ Gary Cutter,² Robert J. Fox,³ Ruth Ann Marrie,⁴ Jason P. Mendoza,⁵ James B. Lewin⁵

¹Biostatistics, Washington University School in St. Louis, School of Medicine, St. Louis, MO; ²Biostatistics, University of Alabama at Birmingham, Birmingham, AL; ³Cleveland Clinic, Cleveland, OH; ⁴Health Sciences Centre Winnipeg, University of Manitoba, Winnipeg, MB, Canada; ⁵Biogen, Cambridge, MA

Background: Employment is often affected in persons with multiple sclerosis (MS), and changes in employment status are associated with reduced quality of life. However, there is limited research on the maintenance of employment and work productivity in persons with MS using a disease-modifying therapy. **Objectives:** To evaluate working status and work productivity in persons with relapsing-remitting MS (RRMS) treated with dimethyl fumarate (DMF) for up to 5 years. **Methods:** In this analysis, we included RRMS North American Research Committee on Multiple Sclerosis (NARCOMS) Registry participants from the United States who reported DMF initiation in any semiannual update survey between Fall 2013 and Spring 2018; participants also had to have ≥1 year of follow-up data. The index survey was considered the survey when DMF was initiated. Work productivity was assessed by reported reduction in hours worked (yes/no) and number of work days missed. Time to change in working status (employed full-time to part-time, employed full- or part-time to not working) was evaluated using the Kaplan-Meier method. Participants were censored at last follow-up or DMF discontinuation, whichever came first. **Results:** A total of 608 participants with RRMS initiated DMF within the study period and had follow-up at 1 year. There were 294 (48.4%) participants employed at initiation of DMF (full-time, 73.8%). Most employed participants were female (86.1%) and Caucasian (82.6%) and had a bachelor's degree or higher education level (65.1%), and the mean (SD) age was 47.7 (9.5) years. The mean (SD) age at diagnosis was 36.0 (8.3) years. The median (interquartile range) PDDS level at initiation was 1 (0, 7) and follow-up was 2 (1, 3.5) years. Overall, 49 (16.7%) participants decreased employment; 13 (4%) changed from full-time to part-time status, and 36 (12%) changed from employed (full- or part-time) to not working. During follow-up, 31 (10.5%)

reported reducing their hours worked and there was a median of 3 (1, 6) missed work days per year. Of the 314 participants not employed, 23 (7.3%) participants went from not employed to employed. **Conclusions:** Among NARCOMS Registry participants who were treated with DMF for up to 5 years, most maintained their baseline level of working status and maintained stable levels of work productivity as assessed by missed work days and the proportion reducing work hours. The NARCOMS Registry provides an opportunity to longitudinally assess outcomes in DMF-treated persons with MS.

Supported by: Biogen. NARCOMS is a project of the CMSC.

Disclosure: *Ambler Salter:* Circulation: Cardiovascular Imaging (statistical editor). *Samantha Lancia:* Nothing to disclose. *Gary Cutter:* AMO Pharmaceuticals, BioLineRx, Brainstorm Cell Therapeutics, Galmed Pharmaceuticals, Hisun Pharmaceuticals, Horizon Pharmaceuticals, Merck, Merck/Pfizer, Neurim, National Heart, Lung, and Blood Institute (protocol review committee); National Institute of Child Health and Human Development (OPRU oversight committee); Ophazyme, Opko Biologics, Reata Pharmaceuticals, Receptos/Celgene, Sanofi-Aventis, Teva Pharmaceuticals (data and safety monitoring committees); Biogen, Click Therapeutics, Genentech, Genzyme, GW Pharmaceuticals, Klein Buendel Inc, MedDay, MedImmune, Osmotica Pharmaceuticals, Perception Neuroscience, Recursion Pharmaceuticals, Roche, Somahlution, TG Therapeutics (consulting fee); Novartis (consulting fee, data and safety monitoring committees). *Robert J. Fox:* Actelion, Biogen, Immune, Novartis (advisory committee, consulting fee); Celgene, EMD Serono, Genentech, Teva (consulting fee). *Ruth Ann Marrie:* Multiple Sclerosis Journal-Exp Transl Clin (co-editor). *Jason P. Mendoza, James B. Lewin:* Biogen (ownership interest, salary).

Keywords: Dimethyl fumarate, Disease-modifying treatments in MS

(DXT02) Early Effect of Ofatumumab on B-Cell Counts and Magnetic Resonance Imaging Activity in Relapsing Multiple Sclerosis Patients: Results from the APLIOS Study

Amit Bar-Or,¹ Edward J. Fox,² Alexandra Goodyear,³ Inga Ludwig,⁴ Morten Bagger,⁵ Dieter A. Haering,⁴ Harald Kropshofer,⁴ Martin Merschhemke,⁴ Heinz Wiendl⁵

¹Center for Neuroinflammation and Experimental Therapeutics and Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA;

²Central Texas Neurology Consultants and Dell Medical School, The University of Texas at Austin, Round Rock, TX; ³Novartis Pharmaceuticals Corporation, East Hanover, NJ;

⁴Novartis Pharma AG, Basel, Switzerland; ⁵Department of Neurology, University of Münster, Münster, Germany

Background: B cells play a major role in the pathogenesis of multiple sclerosis (MS). Ofatumumab, the first fully human anti-CD20 monoclonal antibody, with a monthly 20 mg subcutaneous (s.c.) dosing regimen suppressed 94%-98% of the gadolinium-enhancing (Gd+) lesions vs teriflunomide in the phase 3 ASCLEPIOS I/II relapsing MS (RMS) trials. In APLIOS, the onset of ofatumumab effect on B-cell depletion and magnetic resonance imaging (MRI) activity can be determined. **Objectives:** To evaluate the onset of ofatumumab 20 mg s.c. effect on B-cell depletion and suppression of MRI activity in patients with RMS. **Methods:** APLIOS was a 12-week, open-label, phase 2, bioequivalence study in patients with RMS (N = 284) who received ofatumumab 20 mg (0.4 mL) s.c. loading doses on days 1, 7, and 14, and maintenance doses every 4 weeks from week 4 via an autoinjector pen (SensoReady) or a prefilled syringe. Suppression of CD19+ B cells was measured 9 times over 12 weeks. Gd+ lesion counts were assessed at baseline and at weeks 4, 8, and 12. **Results:** Ofatumumab rapidly depleted circulating B cells, from a median B-cell count of 219 cells/ μ L (day 1) to 10 cells/ μ L (day 4) and 1 cell/ μ L by the end of the loading regimen (week 4). The proportion of patients with B-cell counts <10 cells/ μ L was >65% after the first injection by day 7, 94% by week 4, and sustained >95% at all following injections. Ofatumumab treatment reduced the mean number of Gd+ lesions from 1.5 (baseline) to 0.8, 0.3, and 0.1 by weeks 4, 8, and 12, respectively; the proportion of patients free from Gd+ lesions at the corresponding time points were 66.5%, 86.7%, and 94.1%. **Conclusions:** Ofatumumab 20 mg s.c. monthly dosing regimen resulted in a rapid, close-to-complete and sustained B-cell depletion over 12 weeks, leading to a profound reduction of Gd+ lesions in patients with RMS, consistent with the effects observed in the pooled phase 3 ASCLEPIOS I/II patient population.

Supported by: None

Disclosure: *Amit Bar-Or:* Atara Biotherapeutics, Biogen, Celgene/Receptos, Genentech/Roche, GlaxoSmithKline, Mapi, MedImmune, Merck/EMD Serono, Novartis, Sanofi Genzyme (consulting fee, speakers' bureau). *Edward J. Fox:* Biogen, Celgene, Chugai, EMD Serono, Genentech/Roche, MedDay, Novartis, Sanofi, Teva, TG Therapeutics (advisory work, consulting fee, contracted research, speakers' bureau); Genzyme (advisory board, contracted research, speakers' bureau). *Alexandra Goodyear, Inga Ludwig, Morten Bagger, Dieter A. Haering, Harald Kropshofer, Martin Merschhemke:* Novartis (salary). *Heinz Wiendl:*

Bayer HealthCare, Biogen, Fresenius Medical Care, GlaxoSmithKline (consulting fee, honoraria).

Keywords: Disease-modifying treatments in MS, Imaging and MS, Immunology and MS

(DXT03) Analyses of the Effect of Disease Duration on the Efficacy and Safety of Siponimod in Patients with Active Secondary Progressive Multiple Sclerosis from the EXPAND Study

Amit Bar-Or,¹ Stanley L. Cohan,² Patricia K. Coyle,³ Fred D. Lublin,⁴ Xiangyi Meng,⁵ Wendy Su,⁵ Bruce A.C. Cree⁶

¹Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ²Providence Multiple Sclerosis Center, Providence Brain Institute, Portland, OR; ³Stony Brook University, Stony Brook, NY; ⁴Icahn School of Medicine at Mount Sinai, New York, NY; ⁵Novartis Pharmaceuticals Corporation, East Hanover, NJ; ⁶Department of Neurology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA

Background: Siponimod (Mayzent) is a selective sphingosine 1-phosphate receptor (S1P₁ and S1P₂) modulator, approved in the United States for treatment of relapsing forms of multiple sclerosis (MS), including clinically isolated syndrome, relapsing-remitting MS, and active secondary progressive MS (SPMS). In the phase 3 EXPAND registration trial in SPMS, siponimod significantly reduced risk of 3-month (primary end point) and 6-month confirmed disability progression (CDP) by 21% and 26%, respectively. **Objectives:** Assess efficacy and safety of siponimod in patients with active SPMS in subgroups of patients with MS duration (time since onset of first symptoms) of <16 or \geq 16 years (median value) at baseline. **Methods:** Post hoc analyses were performed in patients with active SPMS, defined as a relapse in the 2 years before screening and/or \geq 1 T1 Gd+ lesion at baseline, randomized to siponimod 2 mg every day or placebo. Efficacy end points: time to 3- and 6-month CDP (as per Expanded Disability Status Scale scores). Adverse events (AEs), serious AEs, and AEs leading to treatment discontinuation were assessed. Analyses for hypothesis generation, without adjustment for multiple comparisons. **Results:** There were 779 patients with active SPMS: 427 with MS duration <16 years (siponimod n = 285; placebo, n = 142) and 352 with duration \geq 16 years (siponimod n = 231; placebo, n = 121). For MS duration of <16 years, siponimod reduced 3- and 6-month CDP risk by 32.4% and 42.7%, respectively, vs placebo (3 month: siponimod n = 68 [23.9%], placebo n = 48 [33.8%], hazard ratio [HR] [95% CI]: 0.68 [0.47, 0.98], P = .0378; 6 month: siponimod n = 48 [16.8%], placebo n = 40 [28.2%], HR [95% CI]: 0.57 [0.38, 0.87], P = .0093). For MS duration \geq 16 years, siponimod had a trend towards reduced 3- and 6-month CDP risk of 31.9% and 27.1%, respectively, vs placebo (3 month: siponimod n = 61 [26.4%], placebo n = 43 [35.5%], HR [95% CI]: 0.68 [0.46, 1.01], P = .0540; 6 month: siponimod n = 51 [22.1%], placebo n = 34 [28.1%], HR [95% CI]: 0.73 [0.47, 1.13], P = .1544). Siponimod was generally well tolerated. Any AE rates: <16 years, 84.9% (siponimod), 75.4% (placebo); \geq 16 years, 89.2% (siponimod), 81.8% (placebo). **Conclusions:** In patients with active SPMS and MS duration <16 years, siponimod significantly reduced 3- and 6-month CDP risk compared with placebo. Siponimod showed a trend towards reduced CDP vs placebo in those with duration \geq 16 years. This may reflect the smaller size or more advanced disease in this subgroup.

Supported by: None

Disclosure: *Amit Bar-Or:* Atara Biotherapeutics, Bayer, Bayhill Therapeutics, Berlex, Biogen, BioMS, Celgene/Receptos, Diogenix, Eli Lilly, F. Hoffman-La Roche, Genentech, GlaxoSmithKline, Guthy-Jackson/GGF, Immunic, Janssen/Actelion, Mapi, MedImmune, Merck/EMD Serono, Novartis, Ono Pharmacia, Roche/Genentech, Sanofi Genzyme, Teva Neuroscience, Wyeth (consulting fee). *Stanley L. Cohan:* AbbVie, Adamas, Alexion, MedDay, Sage Bionetworks (contracted research); Biogen, Novartis, Sanofi Genzyme (consulting fee, contracted research, speakers' bureau); EMD Serono, Pear Therapeutics (consulting fee); Genentech, Roche (contracted research, speakers' bureau). *Patricia K. Coyle:* Accordant, Alexion, Bayer, Biogen, Celgene, Genentech/Roche, Sanofi Genzyme, GlaxoSmithKline, Mylan, Novartis, Serono, TG Therapeutics (consulting fee); Actelion, Alkermes, Corrona LLD, MedDay, PCORI (research grant). *Fred D. Lublin:* AbbVie, Acorda Therapeutics, Apitope, Atara Biotherapeutics, Bayer HealthCare Pharmaceuticals, Biogen, Brainstorm Cell Therapeutics, EMD Serono, Forward Pharma, Innate Immunotherapeutics, Mapi Pharma, MedDay Pharma, MedImmune, Novartis, Orion Biotechnology, Polpharma, Receptos/Celgene, Regeneron, Roche Genentech, Sanofi Genzyme, Teva Neuroscience, TG Therapeutics (consulting fee); Actelion (consulting fee, research support); Multiple Sclerosis and Related Disorders (journal editor); National MS Society, Novartis Pharmaceuticals, Sanofi, Teva Neurosciences, Transparency Life Sciences (research support). *Xiangyi Meng, Wendy Su:* Novartis Pharmaceuticals Corporation (sal-

ary). *Bruce A.C. Cree: Akili, Alexion, Atara, Biogen, EMD Serono, Novartis, TG Therapeutics (consulting fee).*

Keywords: Disease-modifying treatments in MS

(DXT04) Siponimod First-Dose Effects in Patients with Secondary Progressive Multiple Sclerosis Receiving Concomitant Selective Serotonin Reuptake Inhibitor Therapy

Amit Bar-Or,¹ Bruce A.C. Cree,² Le H. Hua,³ Amos Katz,⁴ Derrick Robertson,⁵ Brandon Brown,⁶ Desiree Dunlop,⁶ Xiangyi Meng,⁶ Wendy Su⁶

¹Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ²Department of Neurology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA; ³Lou Ruvo Center for Brain Health, Las Vegas, NV; ⁴Linda Cardinale MS Center, Freehold, NJ; ⁵University of South Florida, Multiple Sclerosis Center, Tampa, FL; ⁶Novartis Pharmaceuticals Corporation, East Hanover, NJ

Background: Selective serotonin reuptake inhibitors (SSRIs), citalopram and escitalopram, are associated with prolonged QTc and torsades de pointes; transient heart rate decrease at initiation is a known effect of sphingosine 1-phosphate (S1P) modulators. Siponimod is an S1P receptor type 1,5 modulator, and is metabolized mainly by CYP2C9, followed by CYP3A4. It is approved in the United States for relapsing forms of multiple sclerosis (MS), including clinically isolated syndrome, relapsing-remitting MS, and active secondary progressive MS. First-dose observation with siponimod is only required in certain cardiac conditions, but it is important to understand the cardiac effects in patients receiving concomitant SSRIs. **Objectives:** Evaluate first-dose effects of siponimod in patients receiving concomitant SSRIs during the EXPAND trial. **Methods:** Analyses included data for the overall siponimod group (with or without SSRI), and subgroups of concomitant siponimod and any SSRI at first dose (day 1), and concomitant siponimod and citalopram or escitalopram on day 1. **Results:** In all, 1105 patients were randomized to siponimod; 167 received an SSRI on day 1 and 85 received citalopram or escitalopram. For those with extended monitoring, in the overall siponimod group, and the any SSRI and citalopram/escitalopram subgroups, most were discharged at 6 hours post first dose (91.1%, 91.4%, and 89.6%, respectively). Day 1 after first dose, 4 patients (0.4%) in the overall siponimod group had serious adverse events (AEs), 2 (0.2%) had bradycardia, and 1 (0.1%) had second-degree atrioventricular (AV) block; no serious AEs occurred in the any SSRI or citalopram/escitalopram subgroups. Few patients in the overall siponimod group had cardiac AEs on day 1: 29 patients (2.6%) had bradycardia, 4 (0.4%) had first-degree AV block, 3 (0.3%) had second-degree AV block, and 3 (0.3%) had prolonged QT interval. Incidence of cardiac AEs was low in the any SSRI subgroup: 3 patients (1.8%) had bradycardia and 3 (1.8%) had prolonged QT interval; in the citalopram/escitalopram subgroup, 2 patients (2.4%) had bradycardia and 1 (1.2%) had prolonged QT interval. In the overall siponimod group, 3 patients (0.3%) discontinued drug due to first- or second-degree AV block, or bradycardia. No patient receiving SSRIs had a cardiac AE causing treatment discontinuation. **Conclusions:** Concomitant SSRI use did not appear to affect cardiac outcomes or heart rate changes associated with siponimod treatment initiation.

Supported by: None

Disclosure: *Amit Bar-Or: Atara Biotherapeutics, Bayer, Bayhill Therapeutics, Berlex, Biogen, BioMS, Celgene/Receptos, Diogenix, Eli Lilly, F. Hoffman-La Roche, Genentech, GlaxoSmithKline, Guthy-Jackson/GGF, Immunic, Janssen/Actelion, Mapi, MedImmune, Merck/EMD Serono, Novartis, Ono Pharmacia, Roche/Genentech, Sanofi Genzyme, Teva Neuroscience, Wyeth (consulting fee).* *Bruce A.C. Cree: Akili, Alexion, Atara, Biogen, EMD Serono, Novartis, TG Therapeutics (consulting fee).* *Le H. Hua: Biogen, Celgene, EMD Serono, Genentech, Genzyme, Novartis (consulting fee).* *Amos Katz: Biogen, Genentech, Novartis, Sanofi (consulting fee).* *Derrick Robertson: Acorda (speakers' bureau); Actelion, MedDay, MedImmune, PCORI, TG Therapeutics (grant support); Alexion, Celgene, Teva Neuroscience (consulting fee, speakers' bureau); Biogen, EMD Serono, Genentech, Novartis, Sanofi Genzyme (consulting fee, grant support, speakers' bureau); Mallinckrodt (grant support, speakers' bureau).* *Brandon Brown, Desiree Dunlop, Xiangyi Meng, Wendy Su: Novartis Pharmaceuticals Corporation (salary).*

Keywords: Disease-modifying treatments in MS

(DXT05) Efficacy of Diroximel Fumarate in Highly Active Relapsing-Remitting Multiple Sclerosis: Interim Results from the Phase 3 EVOLVE-MS-1 Study

Annette Wundes,¹ Enrique Alvarez,² Mark S. Freedman,³ Oksana Mokliatchouk,⁴ Hailu Chen,⁴ Shivani Kapadia,⁴ Jordan Messer,⁴ Robert T. Naismith⁵

¹Department of Neurology, University of Washington Medical Center, Seattle, WA; ²Department of Neurology, University of Colorado School of Medicine, Aurora, CO; ³University of Ottawa and the Ottawa Hospital Research Institute, Ottawa, ON, Canada; ⁴Biogen, Cambridge, MA; ⁵Washington University School of Medicine, St. Louis, MO

Background: Diroximel fumarate (DRF) is a novel oral fumarate recently approved in the United States for relapsing forms of multiple sclerosis (MS). Currently, there are no published data demonstrating the efficacy of DRF in patients with highly active relapsing-remitting MS (RRMS). Lesion load are associated with an increased risk of MS progression. **Objectives:** To evaluate clinical and radiologic efficacy of DRF in patients with highly active RRMS. **Methods:** EVOLVE-MS-1 (trial registration: NCT02634307) is an ongoing, open-label, phase 3 study of long-term safety, tolerability, and treatment effect of DRF over 96 weeks in adults with RRMS. Efficacy outcomes were assessed in a post hoc subgroup analysis of patients from EVOLVE-MS-1 with highly active RRMS (≥ 2 relapses in the year before study entry and ≥ 1 gadolinium-enhancing [Gd+] lesion at baseline). Assessments included adjusted annualized relapse rate (ARR), Gd+ lesion count, and 12-week confirmed disability progression (CDP; ≥ 1.5 -point increase in Expanded Disability Status Scale score from a baseline score of 0, or a ≥ 1 -point increase from a baseline score of 1.0-5.5, sustained for 12 weeks). **Results:** A total of 57 patients with highly active RRMS were enrolled in EVOLVE-MS-1 as of November 30, 2018, 14 of whom were newly diagnosed and naive to disease-modifying therapy. Patients had a mean (SD) time since diagnosis of 4.2 (4.6) years and 2.2 (0.6; range 2-5) relapses in the 12 months before study entry. Median (range) DRF exposure for the highly active subgroup was 84 (1-100) weeks; 72% of patients were on treatment for ≥ 1 year. Adjusted ARR was reduced by 83.3% (95% CI 72.0%-90.7%; $P < .0001$) with DRF treatment compared with the 12 months before study entry (0.36 [95% CI 0.20-0.65] vs 2.22 [95% CI 2.06-2.40]), and 73.7% of patients were relapse-free. Estimated proportion of patients with 12-week CDP at week 48 was 8.3%. Mean (SD) time to disability progression for patients who progressed was 5.4 (4.8) months. Significant reductions in mean (SD) Gd+ lesion counts were observed at week 48 (1.0 [4.7]; $n = 41$) compared with baseline (5.0 [7.7]; $n = 53$; 80% reduction; $P = .0003$), and a greater percentage were Gd+ lesion-free (week 48: 80.5% vs baseline: 9.4%). **Conclusions:** DRF demonstrated notable benefits on clinical and radiologic outcomes in patients with highly active RRMS. These data and comparison with similar cohorts (DEFINE/CONFIRM highly active RRMS) support DRF as a treatment option for patients with MS with active disease.

Supported by: Biogen

Disclosure: *Annette Wundes: Alkermes, AbbVie, Biogen (contracted research); Biogen (consulting fee).* *Enrique Alvarez: Actelion, Bayer, Biogen, Celgene, Genentech, Genzyme, Teva, Novartis, TG Therapeutics (consulting fee); Biogen, Genentech, Novartis, Rocky Mountain MS Center (contracted research).* *Mark S. Freedman: Actelion, Bayer, Biogen, Celgene, Chugai, EMD Canada, Genzyme, Merck Serono, Novartis, F. Hoffman La-Roche, Pendopharm, Sanofi-Aventis (consulting fee); Actelion, Bayer, Biogen, Clene, F. Hoffman La-Roche, Merck Serono, MedDay, Novartis, Sanofi-Aventis (participation as a member in a company advisory board, board of directors, or other similar group); Genzyme Canada (contracted research); Sanofi Genzyme (speakers' bureau).* *Oksana Mokliatchouk, Hailu Chen, Shivani Kapadia, Jordan Messer: Biogen (ownership interest, salary).* *Robert T. Naismith: Alexion, Alkermes, Biogen, Celgene, EMD Serono, Genentech, Novartis, Sanofi Genzyme, TG Therapeutics, Viela Bio (consulting fee).*

Keywords: Disease-modifying treatments in MS, Highly active disease

(DXT06) Real-World Effectiveness of Peginterferon Beta-1a Versus Interferon Beta-1a and Glatiramer Acetate in US Multiple Sclerosis Patients

Anthony T. Reder,¹ Nancy Arndt,¹ Caroline Geremakis,² Jason P. Mendoza,² Ray Su,² Charles Makin,² Megan C. Vignos,² Robin L. Avila²

¹University of Chicago Neurology, Chicago, IL; ²Biogen, Cambridge, MA

Background: Interferons (IFNs) and glatiramer acetate (GA) are effective in reducing relapses in relapsing-remitting multiple sclerosis (RRMS). Matching-adjusted indirect comparisons and propensity score (PS) matching analyses of clinical trial data have evaluated the efficacy of disease-modifying therapies (DMTs). However, studies evaluating their real-world effectiveness are limited. **Objectives:** To compare clinical outcomes, effectiveness, and health care resource measures for patients with multiple sclerosis (MS) initiating subcutaneous (SC) peginterferon beta-1a, SC IFN beta-1a three times weekly (SC IFN), or GA in routine clinical practice. **Methods:** This retrospective, observational comparative effectiveness study used data from the Truven MarketScan Commercial Claims Database. Patients aged 18-65 years presenting at least 1 claim for an MS

diagnosis were selected if they initiated peginterferon beta-1a, SC IFN, or GA between November 2014 and March 2017 as standard of care. PS was derived from baseline characteristics and patients were PS matched, including 685 patients for peginterferon beta-1a vs 2035 for SC IFN, and 858 for peginterferon beta-1a vs 2573 for GA. Demographic, clinical, and health care resource measures were evaluated at baseline and post-index measures were evaluated. Annualized relapse rates (ARRs) were compared using a negative binomial model adjusting for disease duration and prior DMTs. The index date was the first peginterferon beta-1a claim, the first SC IFN claim, or the first GA claim. **Results:** Prior DMT utilization was higher with peginterferon beta-1a vs SC IFN (75.1% vs 15.2%, $P < .0001$), and with peginterferon beta-1a vs GA (75.1% vs 13.3%, $P < .0001$). The mean persistence of time a patient was on peginterferon beta-1a (post-index date) was 483 days for the SC IFN comparison and 497 days for the GA comparison. After adjusting for baseline patient demographics and clinical characteristics and PS matching, ARR during the post-index period was lower with peginterferon beta-1a than with SC IFN (0.18 vs 0.23, relative risk [RR] = 0.795, $P = .02$). There was a trend toward decreased ARR with peginterferon beta-1a vs GA (0.20 vs 0.23, RR = 0.876, $P = .18$). **Conclusions:** ARR was lower with peginterferon beta-1a than with SC IFN, and a trend toward decreased ARR with peginterferon beta-1a vs GA was observed. These real-world results contribute to the data available for decision making in routine clinical practice.

Supported by: Biogen

Disclosure: *Anthony T. Reder:* Bayer, Biogen, Novartis, Serono (unrestricted grant support and/or advisory board compensation). *Nancy Arndt:* Biogen, Genzyme, Novartis, Teva (consulting, advisory board compensation, and/or compensation for speaking). *Caroline Geremakis, Jason P. Mendoza, Ray Su, Charles Makin, Megan C. Vignos, Robin L. Avila:* Biogen (employee, may hold stock and/or stock options).

Keywords: Disease-modifying treatments in MS, Real-world evidence

(DXT07) Injection Site Reactions and Risk of Discontinuation Among New and Experienced Peginterferon Beta-1a Users in the Plegridy Observational Program

Ayo Adeyemi,¹ Nicole Tsao,¹ Arman Altincatal,¹ Maria L. Naylor,¹ Marco Salvetti,² Sibyl Wray,³ Gereon Nelles,⁴ Charles Makin¹

¹Biogen, Cambridge, MA; ²Sapienza University, S. Andrea Hospital, Rome, Italy; ³Hope Neurology, Knoxville, TN; ⁴Neurology, NeuroMed Campus Hohenlind, Cologne, Germany

Background: Injection site reactions (ISRs) are a common adverse event associated with peginterferon beta-1a treatment and may lead to discontinuation. This analysis of the Plegridy Observational Program (POP), a 5-year, phase 4 real-world study, assessed the relationship between ISRs and discontinuation of peginterferon beta-1a. **Objectives:** Assess the risk of treatment discontinuation associated with ISRs in new and experienced users of peginterferon beta-1a. **Methods:** Using POP third interim data from November 2014 to September 2018, patients were classified as new users if they initiated peginterferon beta-1a ≤ 31 days prior to, on, or after study consent; and experienced users if they initiated peginterferon beta-1a > 31 days prior to study consent. Treatment discontinuation was based on physician report, and only the first discontinuation was analyzed. Demographics and baseline characteristics were summarized with descriptive statistics. Frequencies and proportions of individuals who experienced ≥ 1 ISR were calculated for each group. Fisher exact test assessed the relative risk (RR) of discontinuation in those with or without ISRs. Kaplan-Meier analysis assessed the cumulative risk of discontinuation. Data from the fourth interim POP analysis (as of September 2019) will be presented. **Results:** Of the 1126 patients included in this analysis, 576 (51%) were new and 550 (49%) were experienced users of peginterferon beta-1a. Among new and experienced users, 280 (49%) and 147 (27%) reported ≥ 1 ISR, respectively. In the new-user cohort, 148 (53%) patients with ISRs and 135 (46%) patients without ISRs discontinued peginterferon beta-1a treatment (RR: 1.16 [95% CI: 0.98, 1.37]). In experienced users, 44 (30%) patients with ISRs and 124 (31%) patients without ISRs discontinued peginterferon beta-1a treatment (RR: 0.97 [95% CI: 0.73, 1.30]). Over 12 months of treatment in the new-user cohort, the cumulative probability of peginterferon beta-1a discontinuation was 0.38 in patients with ≥ 1 ISR and 0.31 in patients without an ISR. **Conclusions:** These preliminary findings suggest an increased risk of discontinuation in new users of peginterferon beta-1a with ISRs, whereas in experienced users no impact of ISRs on discontinuation was observed. New users of peginterferon beta-1a treatment may benefit from additional information on ISR mitigation and management through discussions with their health care professionals.

Supported by: Biogen

Disclosure: *Ayo Adeyemi, Nicole Tsao, Arman Altincatal, Maria L. Naylor, Charles Makin:* Biogen (employee, may hold stock and/or stock options). *Marco Salvetti:* Biogen, Merck, Novartis, Roche, Sanofi, Teva (received grant support and/or speaker honoraria). *Sibyl Wray:* Bayer, Biogen, Celgene, EMD Serono, Genentech/Roche, Sanofi Genzyme, Novartis, Receptos, TG Therapeutics (paid consultant, speaker, and/or contract researcher). *Gereon Nelles:* Bayer, Biogen, Celgene, Merck, Novartis, Roche (speaker honoraria).

Keywords: Disease-modifying treatments in MS, Injection site reactions

(DXT08) Post Hoc Analysis of Efficacy of Cladribine Tablets in Patients with Relapsing-Remitting Multiple Sclerosis Diagnosed Within 3 or 4 Years Prior to the CLARITY Study

Barry Singer,¹ Gavin Giovannoni,² Nicola De Stefano,³ Matt Mandel,⁴ Yann Hyvert,⁵ Julie Aldridge,⁶ Andrew Galazka,⁶ Caroline Lemieux,⁷ Mark S. Freedman⁸

¹Missouri Baptist Medical Center, St Louis, MO; ²Blizard Institute of Cell and Molecular Science, London, United Kingdom; ³University of Siena, Siena, Italy; ⁴EMD Serono Inc, Billerica, MA; ⁵Merck KGaA, Darmstadt, Germany; ⁶Merck, Aubonne, Switzerland; ⁷EMD Serono Inc, Mississauga, ON, Canada; ⁸University of Ottawa and the Ottawa Hospital Research Institute, Ottawa, ON, Canada

Background: Cladribine tablets 3.5 mg/kg (CT3.5, cumulative dose over 2 years; N = 433) treatment significantly improved clinical and magnetic resonance imaging (MRI) outcomes vs placebo (N = 437) in relapsing-remitting multiple sclerosis (RRMS) in the 96-week CLARITY study. Prior post hoc analyses of CLARITY found that patients with shorter disease duration (< 3 years) responded more favorably to CT3.5 in risk reduction of relapse and odds of achieving disease-free activity status vs those with longer disease duration (≥ 3 years). Further analyses are needed to assess the extent of benefit of CT3.5 in patients with a relatively shorter disease duration. **Objectives:** This post hoc analysis further examined efficacy outcomes in CLARITY patients who were early in their disease course (< 3 or < 4 years newly diagnosed [YND]) at study enrollment. **Methods:** Analyses were performed by treatment (CT3.5 vs placebo) and MS-duration groups (< 3 YND, < 4 YND at CLARITY enrollment). End points were relapse, 3- and 6-month confirmed disability progression (CDP, based on Expanded Disability Status Scale score), magnetic resonance imaging (MRI) activity, and no evidence of disease activity (NEDA) status (no relapse, CDP, or MRI activity). P values are nominal. **Results:** These MS-duration group analyses were carried out in 492 patients: < 3 YND: CT3.5 N = 228, placebo N = 207; < 4 YND: CT3.5 N = 256, placebo N = 236. In both MS-duration groups, CT3.5 significantly reduced annualized relapse rate (< 3 YND: 0.14 vs 0.37 [60.7% reduction]; < 4 YND: 0.14 vs 0.36 [61.6% reduction]; $P < .0001$), and increased the number of patients who were relapse-free through week 96 (< 3 YND: 75.4% vs 55.1%; < 4 YND: 75.0% vs 55.9%) vs placebo. CT3.5 treatment increased the probability of being free from 3- or 6-month CDP through week 96 (3-month CDP at week 96 in both groups: 0.84-0.85 vs 0.76 [both groups]; 6-month CDP at week 96: 0.89-0.90 vs 0.83-0.84) vs placebo. CT3.5 also significantly reduced the adjusted mean (95% CI) cumulative number of new T1 gadolinium-enhancing (< 3 YND: 0.11 [0.08-0.16] vs 1.0 [0.73-1.36]; < 4 YND: 0.11 [0.08-0.16] vs 0.93 [0.69-1.24]) and active T2 (< 3 YND: 0.41 [0.33-0.51] vs 1.68 [1.36-2.06]; < 4 YND: 0.41 [0.33-0.50] vs 1.63 [1.34-1.97]) lesions, and increased the number of patients achieving NEDA status using 3-month (< 3 YND: 40.4% vs 11.1%; < 4 YND: 39.5% vs 11.4%) or 6-month (< 3 YND: 41.2% vs 11.6%; < 4 YND: 40.2% vs 12.3%) CDP vs placebo (all $P < .0001$) at 96 weeks. **Conclusions:** Overall, CT3.5 treatment improved clinical and MRI outcomes in CLARITY patients who were early in their disease course.

Supported by: None

Disclosure: *Barry Singer:* AbbVie, Biogen, Novartis, Roche, Sanofi Genzyme (consulting fee, research support); Alexion, Bayer, Celgene, EMD Serono, Genentech, Teva, TG Therapeutics (consulting fee); Alkermes, MedImmune (research support). *Gavin Giovannoni:* AbbVie, Actelion, Almirall, Atara Bio, Bayer Schering Pharma, Five Prime, GlaxoSmithKline, GW Pharma, Merck KGaA, Pfizer Inc, Protein Discovery Laboratories, Sanofi Genzyme, Teva Pharmaceuticals Industries Ltd, UCB, Vertex Pharmaceuticals (consulting fee); Biogen, Ironwood, Merck and Co, Novartis (consulting fee, unrelated research support). *Nicola De Stefano:* Bayer Schering AG, Merck Serono SA, Novartis Pharma AG, Sanofi-Aventis, Serono Symposia International Foundation, Teva Pharmaceutical Industries (consulting fee). *Matt Mandel, Julie Aldridge, Caroline Lemieux:* EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). *Yann Hyvert:* Merck KGaA (salary). *Andrew Galazka:* Merck KGaA, Aubonne, Switzerland (a business of Merck KGaA, Darmstadt, Germany) (salary). *Mark S. Freedman:* Actelion, Bayer HealthCare, Biogen, Chugai, EMD Canada, Genzyme, F. Hoffman La Roche, Novartis, Sanofi, Teva (consulting fee).

Keywords: Disease-modifying treatments in MS

(DXT09) Exploration of Factors Which Influence Treatment Decisions of Patients with Multiple Sclerosis

Belinda Bardsley,¹ John Haynes,² Edith Cinc,³ Elise Heriot,¹ K.-J. Lazarus,¹ Melanie McMurtrie,¹ Richard A.L. Macdonell¹

¹Neurology, Austin Health, Heidelberg, VIC, Australia; ²Faculty of Pharmacy and Pharmaceutical Sciences, Monash University, Parkville, Parkville, VIC, Australia; ³Austin Health and Florey Institute of Neuroscience and Mental Health, Melbourne, VIC, Australia

Background: The past 10 years have brought a wide variety of therapeutic options to Australian patients with relapsing-remitting multiple sclerosis (RRMS). In a complex treatment landscape for an unpredictable disease, it is important to understand how patients view the various factors that contribute to making an informed therapeutic choice. **Objectives:** Identify the factor rated by patients with RRMS (PwRRMS) as having the most influence on treatment choice. **Methods:** This noninterventional, exploratory study prospectively enrolled 78 patients assigned to 1 of 3 groups: 1) initial treatment (n = 25), 2) switching to alternate treatment (n = 27), and 3) stable on treatment (n = 26). Baseline demographic and MS data were collected. Participants completed the survey where they rated factors from most to least important: drug safety, efficacy, ease of use, mode of administration, mechanism of action, concern about disability, requirement for follow-up safety monitoring, balance of risk/benefit, and value of discussion with MS nurse and neurologist. **Results:** The factors ranked first by most participants in influencing treatment choice were (in order): 1) concern about disability (31/78 participants), 2) perception of efficacy (16/78), and 3) perception of safety (11/78). This ranking order was consistent across all groups. 97% of participants were satisfied with the process around choosing treatment, and 92% reported they felt extremely comfortable with their treatment decision. **Conclusions:** Our data indicates that concern about disability is the largest driving factor for PwRRMS choosing between treatments regardless of whether they are starting for the first time, planning a switch in therapy, or are currently stable on an MS medication.

Supported by: None

Disclosure: *Belinda Bardsley:* Biogen Pty Ltd, Roche Pty Ltd (consulting fee); Merck Pty Ltd (contracted research, speakers' bureau, travel support). *John Haynes:* Nothing to disclose. *Edith Cinc:* Biogen Pty Ltd (consulting fee, speakers' bureau); Merck Pty Ltd, Novartis Pty Ltd (consulting fee); Teva Pty Ltd (speakers' bureau). *Elise Heriot:* Clene Nanomedicine, MedDay Pharmaceuticals, Roche Pty Ltd (travel to investigator meeting). *K.-J. Lazarus:* Biogen Pty Ltd (speakers' bureau). *Melanie McMurtrie:* Clene Nanomedicine, MedDay (travel to investigator meeting). *Richard A.L. Macdonell:* Celgene, Genzyme, Novartis, Roche Pty Ltd (consulting fee); Merck Pty Ltd (travel support).

Keywords: Comprehensive care and MS, Disease-modifying treatments in MS, Nursing management in MS

(DXT10) Siponimod Affects Disability Progression in Patients with Secondary Progressive Multiple Sclerosis Independent of Relapse Activity: Results from the Phase 3 EXPAND Study

Bruce A.C. Cree,¹ Robert J. Fox,² Gavin Giovannoni,³ Patrick Vermersch,⁴ Amit Bar-Or,⁵ Ralf Gold,⁶ Nicolas Rouyre,⁷ Goeril Karlsson,⁷ Frank Dahlke,⁷ Ludwig Kappos⁸

¹Department of Neurology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA; ²Mellen Center for Multiple Sclerosis, Cleveland Clinic, Cleveland, OH; ³Queen Mary University of London, Blizard Institute, Barts and the London School of Medicine and Dentistry, London, United Kingdom; ⁴Department of Neurology, University of Lille, CHU Lille, Lille, France; ⁵Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ⁶Department of Neurology, St. Josef-Hospital/Ruhr-University Bochum, Bochum, Germany; ⁷Novartis Pharma AG, Basel, Switzerland; ⁸Departments of Medicine, Clinical Research, Biomedicine, and Biomedical Engineering, Neurologic Clinic and Policlinic, University Hospital Basel, University of Basel, Basel, Switzerland

Background: Siponimod (Mayzent) is a selective sphingosine 1-phosphate receptor (S1P₁, S1P₃) modulator, approved in the United States for the treatment of relapsing forms of multiple sclerosis (MS), including active secondary progressive MS (SPMS). In the phase 3, randomized, double-blind, placebo-controlled EXPAND trial, siponimod reduced the risk of 3- and 6-month confirmed disability progression (CDP) by 21% and 26%, respectively, compared with placebo, in patients with SPMS. Subgroup analyses of EXPAND data suggest that a proportion of the effect of siponimod on CDP was attributable to effects on relapse-independent disability progression. **Objectives:** Assess the impact of siponimod on CDP in patients with/without relapses to uncouple treatment effects on CDP from those on relapses. **Methods:** In EXPAND, patients (aged 18-60 years) with SPMS and Expanded Disability Status Scale score of 3.0-6.5 were included in the study and received once-daily oral siponimod 2 mg

or placebo for up to 3 years. We analyzed the impact of siponimod on CDP by subgroup analysis using the Cox model on time to 3- and 6-month CDP in patients with or without relapses in the 1 and 2 years before study; principal stratum analysis to estimate the effect in patients who would not have relapsed on-study at the month 12, month 18, and month 24 timepoints, regardless of treatment; and Cox model on time to 3-/6-month CDP in the overall population, censoring at time of first relapse. **Results:** For nonrelapsing patients in the 1 and 2 years before study, risk reductions were 18% [hazard ratio [HR], 0.82 [CI: 0.66, 1.02]] and 13% [0.87 [0.68, 1.11]], respectively, for 3-month CDP, and 25% [0.75 [0.59, 0.96]] and 18% [0.82 [0.62, 1.08]], respectively, for 6-month CDP; for relapsing patients, risk reductions were 33% and 33% (3-month CDP), and 30% and 37% (6-month CDP), respectively. In principal stratum estimates, siponimod reduced 3-month CDP by 14%-20% and 6-month CDP by 29%-33% in nonrelapsing patients across the 3 timepoints, suggesting that these patients achieved a large proportion of the effect in the overall population. Cox model censoring at relapse confirmed beneficial effects, reaching nominal statistical significance (6-month CDP: HR 0.77 [0.62, 0.96]). **Conclusions:** Siponimod reduces risk of CDP in patients with SPMS with or without relapses, indicating that the effects on disability are largely independent from those on relapses. Patients with or without relapses may thus benefit from treatment with siponimod.

Supported by: None

Disclosure: *Bruce A.C. Cree:* Akili, Alexion, Atara, Biogen, EMD Serono, Novartis, TG Therapeutics (consulting fee). *Robert J. Fox:* Actelion, EMD Serono, Genentech, Teva (consulting fee); Biogen, Novartis (consulting fee, contracted research). *Gavin Giovannoni:* AbbVie, Bayer Schering, Biogen, Canbex, Eisai, Elan, Five Prime, Genentech, Sanofi Genzyme, GlaxoSmithKline, GW Pharmaceuticals, Ironwood, Merck Serono, Novartis, Pfizer, Roche, Synthron BV, Teva (consulting fee); Multiple Sclerosis and Related Disorders (Eleuver) (co-chief editor); UCB Pharma (grants). *Patrick Vermersch:* Bayer HealthCare, Biogen, Merck Serono, Novartis, Teva Neuroscience (consulting fee, research support); Celgene, Roche, Sanofi, Servier (consulting fee); SAGE, Thieme Verlag (editor). *Amit Bar-Or:* Atara Biotherapeutics, Bayer, Bayhill Therapeutics, Berlex, Biogen, BioMS, Celgene/Receptos, Diogenix, Eli Lilly, F. Hoffman-La Roche, Genentech, GlaxoSmithKline, Guthy-Jackson/GGF, Immunic, Janssen/Actelion, Mapi, MedImmune, Merck/EMD Serono, Novartis, Ono Pharmacia, Roche/Genentech, Sanofi Genzyme, Teva Neuroscience, Wyeth (consulting fee). *Ralf Gold:* Bayer HealthCare, Biogen, Merck Serono, Novartis, Teva Neuroscience (consulting fee, contracted research); SAGE, Thieme Verlag (personal fees for serving as an editor). *Nicolas Rouyre, Goeril Karlsson, Frank Dahlke:* Novartis Pharma AG (salary). *Ludwig Kappos:* Actelion, Alkermes, Allergan, Almirall, Bayer, Biogen, Celgene, CSL Behring, df-mp, European Union, EXCEMED, GeNeuro, Genzyme, Merck, Mitsubishi Tanabe Pharma, Novartis, Pfizer, Receptos/Celgene, Roche, Roche Research Foundations, Sanofi-Aventis, Santhera, Swiss Multiple Sclerosis Society, Swiss National Research Foundation, Teva, UCB Pharma, Vianex (contracted research); Neurostatus products (license fees).

Keywords: Disease-modifying treatments in MS

(DXT11) The Implications of Suboptimal Treatment Outcomes with Disease-Modifying Drugs in Employees with Multiple Sclerosis

Carrie M. Hersh,¹ Richard A. Brook,² Ian A. Beren,³ Nicholas J. Rohrbacher,³ Lori Lebson,⁴ Christian Henke,⁵ Amy L. Phillips⁶

¹Lou Ruvo Center for Brain Health, Cleveland Clinic, Las Vegas, NV; ²Better Health Worldwide, Inc, Newfoundland, NJ; ³The HCMS Group, LLC, Pittsburgh, PA; ⁴EMD Serono, Inc, Rockland, MA; ⁵Merck KGaA, Darmstadt, Germany; ⁶Health Economics & Outcomes Research, EMD Serono, Inc, Rockland, MA

Background: A better understanding of the implications of suboptimal treatment outcomes in employees with multiple sclerosis (MS) may elucidate opportunities for improving care management. **Objectives:** To evaluate suboptimal treatment outcomes with disease-modifying drugs (DMDs) in patients with MS from an employer perspective. **Methods:** US Human Capital Management Services database employees were eligible if they had: ≥ 2 claims with MS diagnoses (ICD-9 CM 340.xx/ICD-10 CM G35) from January 1, 2010–March 31, 2019, ≥ 1 once-/twice-daily oral or any self-injectable DMD claim (first claim=index), continuous eligibility 6-months pre- (baseline) and 1-year post-index (follow-up), no baseline DMD, age 18-64. Suboptimal treatment outcomes were DMD nonadherence (proportion of days covered $< 80\%$), DMD discontinuation (treatment gap > 60 days), DMD switch, or relapse (MS-related hospitalization, emergency room visit, or outpatient visit with corticosteroid ± 7 days). A 2-part logistic-GLM model evaluated costs controlling for age, tenure, marital status, race, exempt status, full-/part-time, salary, location, Charlson Comorbidity Index, smoking, and relapse. **Results:** Of 2173 employees with ≥ 2 MS diagnoses, 488 (22.5%) met eligibil-

ity. Half (n = 247 [50.6%]) had suboptimal treatment outcomes (39.5% nonadherence, 9.8% discontinuation, 10.9% switching, 20.7% relapse; indicators not mutually exclusive). Baseline characteristics were similar for employees with vs without suboptimal treatment outcomes: mean age: 42.60 vs 43.87; female: 73.7% vs 71.4%; white/black/Hispanic: 32.0%/5.3%/3.6% vs 29.5%/5.8%/6.6%; married: 23.1% vs 27.4%; mean salary: \$61,898 vs \$68,737; hypertension: 14.6% vs 16.6%; hyperlipidemia: 11.7% vs 12.9%; gastrointestinal disease: 16.6% vs 12.9%; tobacco use: 3.2% vs 2.5%; baseline magnetic resonance imaging: 57.9% vs 58.9%; baseline relapse: 22.7% vs 19.9%. Employees with vs without suboptimal treatment outcomes had higher all-cause medical (\$12,730 vs \$6428; $P < .0001$), MS-related medical (\$5444 vs \$2652; $P < .0001$), non-DMD pharmacy (\$2920 vs \$2169; $P = .0199$), sick leave (\$1247 vs \$908; $P = .0274$), and short-term disability (\$934 vs \$146; $P = .0001$) costs. Long-term disability (\$751 vs \$0; $P = .1250$) and Workers' Compensation (\$56 vs \$24; $P = .1276$) costs did not differ significantly. **Conclusions:** Half of employees with MS had suboptimal 1-year treatment outcomes (nonadherence, discontinuation, switching, relapse). Suboptimal response to DMD treatment was associated with higher medical, sick leave, and short-term disability costs.

Supported by: None

Disclosure: *Carrie M. Hersb:* Biogen (consulting fee, contracted research, speakers' bureau); EMD Serono, Genentech, Genzyme, Novartis (consulting fee, speakers' bureau); PCORI (contracted research). *Richard A. Brook:* Better Health Worldwide, Inc (salary); EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (study funding). *Ian A. Beren, Nicholas J. Rohrbacker:* EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (study funding); The HCMS Group, LLC (salary). *Lori Lebson:* EMD Serono, Inc (salary). *Christiane Henke:* Merck KGaA (salary). *Amy L. Phillips:* EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary).

Keywords: Disease-modifying treatments in MS, Economic issues and MS

(DXT12) Real-World Effectiveness of Peginterferon Beta-1a Versus Teriflunomide in US Multiple Sclerosis Patients

Cortnee Roman,¹ Caroline Geremakis,² Jason P. Mendoza,² Ray Su,² Charles Makin,² Megan C. Vignos,² Robin L. Avila²

¹Rocky Mountain Multiple Sclerosis Clinic, Salt Lake City, UT; ²Biogen, Cambridge, MA

Background: Both peginterferon beta-1a and teriflunomide (TERI) have been shown in clinical studies to effectively reduce relapses in relapsing-remitting multiple sclerosis (RRMS). In lieu of head-to-head studies, efficacy comparisons between disease-modifying therapies (DMTs) can be assessed through matched-adjusted indirect comparisons and propensity score (PS) matching. However, real-world studies are also valuable in demonstrating the differences in the effectiveness of multiple sclerosis (MS) DMTs. **Objectives:** To compare clinical outcomes and effectiveness for patients with MS initiating subcutaneous (SC) peginterferon beta-1a and TERI in routine clinical practice. **Methods:** Utilizing the Truven MarketScan Commercial Claims Database, a retrospective, observational comparative effectiveness study was performed. Patients aged 18-65 years presenting at least 1 claim for an MS diagnosis were selected if they were initiated on peginterferon beta-1a or TERI between November 2014 and March 2017 as standard of care. PS was derived from the baseline characteristics and patients were PS matched, including 704 patients for peginterferon beta-1a vs 2099 for TERI. Demographic, clinical, and health care resource measures were evaluated at baseline and postindex measures were evaluated. Annualized relapse rate (ARRs) were compared using a negative binomial model adjusting for disease duration and number of prior DMTs. The index date was the first claim for peginterferon beta-1a or TERI, for each respective cohort. **Results:** Significant differences in prior DMT utilization were observed between the peginterferon beta-1a vs TERI cohorts (77.5% vs 56.9%, $P = .0005$). After adjustment for baseline patient demographics and clinical characteristics and PS matching, there was a significant difference in ARR during the postindex period between peginterferon beta-1a-treated and TERI-treated patients (0.26 vs 0.33, $P = .0003$). **Conclusions:** Statistically significant differences in ARR between peginterferon beta-1a and TERI were observed. These findings from real-world data help support decision making in routine clinical practice.

Supported by: Biogen

Disclosure: *Cortnee Roman:* Alexion, Biogen, Celgene, Bristol Myers Squibb, EMD Serono (advisory board and speaker bureau compensation). *Caroline Geremakis, Jason P. Mendoza, Ray Su, Charles Makin, Megan C. Vignos, Robin L. Avila:* Biogen (employee, may hold stock and/or stock options).

Keywords: Disease-modifying treatments in MS, Real-world evidence

(DXT13) Disease-Modifying Therapies: How Confident Are We That We Understand Their Risk?

Cortnee Roman,¹ Christen Kutz,² Timothy West¹

¹Rocky Mountain Multiple Sclerosis Clinic, Salt Lake City, UT; ²Colorado Springs Neurological Associates, Colorado Springs, CO

Background: An increasing number of therapeutic options are available for patients with multiple sclerosis (MS). Each medication has a unique efficacy and safety profile that needs to be critically evaluated to optimize the risk and benefit for each patient's individual treatment plan. While most efficacy data are obtained in phase 3 trials, safety profiles may change over time as patients use these medications in a real-world setting. Participants in clinical trials are often younger and have fewer comorbidities than the general MS population. To understand the evolving nature of serious adverse events (SAEs), it is important to have up-to-date information on the incidence of these events balanced by the number of patients exposed to the medication, as well as the patient-years exposure.

Objectives: To compare rates of SAEs for approved MS disease-modifying therapies in relation to current patient exposures and patient-years of exposure. **Methods:** A retrospective analysis was performed to obtain SAE data from the manufacturers of the most commonly prescribed MS medications. **Results:** As of the writing of this abstract, cases of progressive multifocal leukoencephalopathy have been reported for natalizumab (825), fingolimod (30), dimethyl fumarate (8), ocrelizumab (8), teriflunomide (1), and alemtuzumab (1). Forty-five cases of cryptococcal meningitis have occurred with fingolimod. Multiple cases of Stevens-Johnson and a single fatal case of toxic epidermal necrolysis has been reported with teriflunomide. Thirteen cases of ischemic and hemorrhagic stroke or arterial dissection have been reported with alemtuzumab. This led to a recent update and boxed warning in the US prescribing information for alemtuzumab. The approximate number of patients exposed to alemtuzumab before this update was approximately 22,000, representing approximately 45,000 patient-years. **Conclusions:** These findings suggest that newly approved medications may require 20,000+ patients with 40,000+ patient-years to uncover SAEs in the real-world setting. Current patient exposures for injectable, oral, and infusible MS medications range from 2,000 to over 500,000, while patient-years range from 10,000 to over 2.5 million. A comparison of the above (and additional) SAE rates by patient exposure will be presented.

Supported by: None

Disclosure: *Cortnee Roman:* Alexion, Biogen, Bristol Myers Squibb, Genentech, Novartis, Sanofi Genzyme (speakers' bureau); Alexion, Biogen, Bristol Myers Squibb, EMD Serono, Genentech, Novartis, Sanofi Genzyme (consulting fee); Biogen, Bristol Myers Squibb, Genentech, Novartis, Sanofi Genzyme (advisory board). *Christen Kutz:* Biogen, EMD Serono, Mallinckrodt, Novartis (consulting fee). *Timothy West:* Biogen, EMD Serono, Novartis (consulting fee, speakers' bureau); Genentech (consulting fee); Genzyme (speakers' bureau).

Keywords: Comparative safety, Disease-modifying treatments in MS

(DXT14) Long-Term Safety and Efficacy of Eculizumab in Neuromyelitis Optica Spectrum Disorder

Dean Wingerchuk,¹ Sean J. Pittock,² Achim Berthele,³ Kazuo Fujihara,^{4,5,6} Ho Jin Kim,⁷ Michael Levy,^{8,9} Jacqueline Palace,¹⁰ Ichiro Nakashima,^{4,11} Murat Terzi,¹² Natalia Totolyan,¹³ Shanthy Viswanathan,¹⁴ Kai-Chen Wang,^{15,16} Kerstin Allen,¹⁷ Kenji P. Fujita,¹⁷ Marcus Yountz,¹⁷ Róisín Armstrong¹⁷

¹Mayo Clinic, Scottsdale, AZ; ²Mayo Clinic, Rochester, MN; ³Technical University of Munich, Munich, Germany; ⁴Tohoku University, Sendai, Japan; ⁵Fukushima Medical University, Fukushima City, Japan; ⁶Southern TOHOKU Research Institute for Neuroscience (STRINS), Koriyama, Japan; ⁷National Cancer Center, Goyang, Korea, Republic of (South); ⁸Massachusetts General Hospital and Harvard Medical School, Boston, MA; ⁹Johns Hopkins University, Baltimore, MD; ¹⁰John Radcliffe Hospital, Oxford, United Kingdom; ¹¹Tohoku Medical and Pharmaceutical University, Sendai, Japan; ¹²Ondokuz Mayıs University, Samsun, Turkey; ¹³First Pavlov State Medical University of St Petersburg, St Petersburg, Russian Federation; ¹⁴Kuala Lumpur Hospital, Kuala Lumpur, Malaysia; ¹⁵Cheng-Hsin General Hospital, Taipei, Taiwan; ¹⁶National Yang Ming University, Taipei, Taiwan; ¹⁷Alexion Pharmaceuticals, Boston, MA

Background: Neuromyelitis optica spectrum disorder (NMOSD) relapses can cause significant and irreversible neurologic disability. Eculizumab, a terminal complement inhibitor, reduces the risk of NMOSD relapse in patients with aquaporin-4 immunoglobulin G (AQP4-IgG)-positive NMOSD. In the PREVENT study, eculizumab reduced the risk of relapse by 94.2% vs placebo (hazard ratio 0.058 [95% CI 0.017, 0.197]; $P < .0001$). The rate of adverse events (AEs)/100 patient-years (PYs) was 749.3 and 1160.9 for eculizumab and placebo, respectively. **Objectives:** To present combined long-term safety and efficacy data for eculizumab from the randomized, double-blind, placebo-controlled PREVENT study (trial registration: NCT01892345) and its ongoing

open-label extension (OLE) (NCT02003144) in patients with AQP4-IgG-positive NMOSD. **Methods:** Patients with AQP4-IgG-positive NMOSD received eculizumab 1200 mg/2 weeks (maintenance dose). Eculizumab safety and efficacy data from the PREVENT and OLE studies (data cutoff: October 31, 2018) were combined. **Results:** Overall, 137 patients received eculizumab. These patients were followed up for a median of 107.9 (range 5.1-237.9) weeks and a combined total of 282.3 PYs after eculizumab initiation. The rates of AEs and serious AEs per 100 PYs were 763.1 and 37.6, respectively. The most common AEs included headache (27.0%), upper respiratory tract infection (25.5%), nasopharyngitis (22.6%), urinary tract infection (16.8%), nausea (16.1%), back pain (15.3%), and diarrhea (15.3%). Common serious AEs, excluding NMOSD relapses, were pneumonia (2.9%), urinary tract infection (2.9%), and optic neuritis (2.2%). There was 1 death during PREVENT (pulmonary empyema), and 1 patient developed *Neisseria gonorrhoeae* infection in the OLE. No patient had a meningococcal infection. In total, 8/137 (5.8%) patients treated with eculizumab had an adjudicated on-trial relapse; an estimated 93.9% (95% CI 87.5%, 97.1%) of patients were relapse-free at 192 weeks. **Conclusions:** In this long-term analysis, eculizumab was well tolerated and reported AEs were consistent with its established safety profile in other indications. The percentage of relapse-free patients remained high (approximately 94%) through 192 weeks.

Supported by: None

Disclosure: *Dean Wingerchuk:* MedImmune, Novartis, Biogen, Celgene, Genentech, TG Therapeutics, Arcus Medica, Reistone (consulting fee); *Terumo BCT, Alexion Pharmaceuticals (fees for non-CME/CE services received directly from commercial interest or its agent).* *Sean J. Pittock:* Alexion Pharmaceuticals (all compensation paid directly to Mayo Clinic, consulting fee, contracted research); *Astellas (all compensation paid directly to Mayo Clinic, consulting fee); Autoimmune Encephalitis Alliance (all compensation paid directly to Mayo Clinic, contracted research); Grifols (all compensation paid directly to Mayo Clinic, contracted research); MedImmune (all compensation paid directly to Mayo Clinic, consulting fee, contracted research); UCB (consulting fee, personal compensation for attending UCB advisory board meeting in Stockholm, Sweden, on Sept 10, 2019).* *Achim Berthele:* Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent, speakers' bureau). *Kazuo Fujihara:* Alexion Pharmaceuticals, Biogen, Chugai, Mitsubishi-Tanabe, Novartis (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent); *Asahi Medical, Bayer, Eisai, Roche, Teijin (fees for non-CME/CE services received directly from commercial interest or its agent).* *Ho Jin Kim:* Alexion Pharmaceuticals, Celltrion, Eisai, HanAll BioPharma, Merck Serono, Novartis, Sanofi Genzyme, Teva-Handok, Viela Bio (consulting fee); *Jerkal of Clinical Neurology (associate editor); MedImmune/Viela Bio (steering committee); Multiple Sclerosis Journal (co-editor).* *Michael Levy:* Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent); *Genentech, Quest Diagnostics, Viela Bio (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent).* *Jacqueline Palace:* Biogen, Chugai (contracted research); *LEK, Viela Bio, Guidepoint (consulting fee); Merck Serono (meeting/lecture/workshop participation); Novartis, Roche, Argenx (speakers' bureau); UCB, Viela Bio, Roche (conference/lecture participation).* *Ichiro Nakashima:* Alexion (consulting fee). *Murat Terzi, Shanbhi Viswanathan, Kai-Chen Wang:* Nothing to disclose. *Natalia Totolyan:* Alexion, Janssen, Novartis, Roche, Sanofi, Receptos Inc, Biocad (Russia) (contracted research); *Merck (consulting fee); Roche, Sanofi (fees for lectures).* *Kerstin Allen, Marcus Yountz, Roisin Armstrong:* Alexion Pharmaceuticals (ownership interest, salary). *Kenji P. Fujita:* Alexion Pharmaceuticals, Ablynam Pharmaceuticals (ownership interest, salary).

Keywords: Long-term safety and efficacy of eculizumab in NMOSD

(DXT15) Inebilizumab Reduces Neuromyelitis Optica Spectrum Disorder Disability Worsening: Outcomes and Long-Term Follow-up Data from the N-MOMentum Trial

Dean Wingerchuk,¹ Romain Marignier,² Jeffrey L. Bennett,³ Ho Jin Kim,⁴ Brian Weinschenker,⁵ Sean J. Pittock,⁶ Kazuo Fujihara,⁷ Friedemann Paul,⁸ Gary Cutter,⁹ Ari Green,¹⁰ Orhan Aktas,¹¹ Hans-Peter Hartung,¹² Fred D. Lublin,¹³ Maureen A. Mealy,¹⁴ Jorn Drappa,¹⁴ Gerard Barron,¹⁴ Soraya Madani,¹⁴ Dewei She,¹⁴ Daniel Cimbara,¹⁴ William Rees,¹⁴ John N. Ratchford,¹⁴ Eliezer Katz,¹⁴ Bruce A.C. Cree¹⁵

¹Mayo Clinic, Scottsdale, AZ; ²Lyon University Hospital, Lyon, France; ³School of Medicine, University of Colorado, Aurora, CO; ⁴Research Institute and Hospital of National Cancer Center, Goyang, Korea, Republic of (South); ⁵Neurology, Mayo Clinic, Rochester, MN; ⁶Mayo Clinic, Rochester, MN; ⁷Department of Multiple Sclerosis Therapeutics, Southern Tohoku Research Institute for Neuroscience (STRINS), Koriyama, Japan; ⁸Experimental and Clinical Research Center, Max Delbrueck Center for Molecular Medicine and Charité – Universitätsmedizin Berlin, Berlin, Germany; ⁹BioStatistics, University of Alabama at Birmingham, Birmingham, AL; ¹⁰Department of Neurology and Department of

Ophthalmology, UCSF Weill Institute for Neurosciences, San Francisco, CA; ¹¹Medical Faculty, Heinrich-Heine-University, Dusseldorf, Germany; ¹²Department of Neurology, UKD, Heinrich-Heine-University Dusseldorf, Dusseldorf, Germany; ¹³Icahn School of Medicine at Mount Sinai, New York, NY; ¹⁴Viela Bio, Gaithersburg, MD; ¹⁵Department of Neurology and Department of Ophthalmology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA

Background: Neuromyelitis optica spectrum disorder (NMOSD) is a rare, relapsing, autoimmune, inflammatory disease of the central nervous system. Disability accumulates with repeated attacks, severely affecting quality of life. Inebilizumab, an anti-CD19 monoclonal B-cell-depleting antibody, was assessed in N-MOMentum, a randomized, placebo-controlled, double-masked trial in patients with NMOSD. **Objectives:** To assess the effectiveness of inebilizumab on disability outcomes in N-MOMentum and determine if severity of pre-existing disability influenced efficacy. **Methods:** Adults with NMOSD and an Expanded Disability Status Scale (EDSS) score ≤ 8 were randomized 3:1 to receive inebilizumab 300 mg or placebo on days 1 and 15. The randomized controlled period (RCP) was 28 weeks or up to an adjudicated attack. The proportion of patients with disability worsening (EDSS score change ≥ 2 from a baseline of 0, ≥ 1 from a baseline of 1-5, or ≥ 0.5 from a baseline of ≥ 5.5) was assessed by logistic regression. Change from baseline in modified Rankin Scale scores was analyzed by the Wilcoxon-Mann-Whitney odds approach. Subgroup analysis by baseline EDSS score of the primary outcome (time to adjudicated attack) was performed by Cox proportional hazards regression. **Results:** The median (range) baseline EDSS score of the 174 patients receiving inebilizumab was 3.5 (0-8) and 4.0 (1-8) for the 56 receiving placebo; 18.0% and 30.4% had disability worsening at week 12, respectively. At the end of RCP, 15.5% of patients on inebilizumab and 33.9% on placebo had disability worsening; odds ratio (95% CI): 0.370 (0.185-0.739); $P = .0049$. Of the 9632 paired comparisons, modified Rankin Scale outcomes at end of RCP were better with inebilizumab than placebo in 51.5% of cases and were equal in 21.9% of cases; adjusted odds ratio (95% CI): 1.663 (1.195-2.385); $P = .0023$. Inebilizumab reduced the risk of attack compared with placebo in patients with baseline EDSS score in the lower (<5) or upper half (≥ 5) of the 10-point scale; hazard ratios (95% CI): 0.257 (0.120-0.552); $P = .0005$ and 0.367 (0.137-0.981); $P = .0456$, respectively; the treatment effect was not significantly different (interaction test, $P = .6363$). Long-term, open-label follow-up data will be presented. **Conclusions:** In N-MOMentum, disability outcomes were significantly better with inebilizumab monotherapy than with placebo. Inebilizumab reduced the risk of attack in patients with NMOSD irrespective of the level of pre-existing disability.

Supported by: None

Disclosure: *Dean Wingerchuk:* Alexion Pharmaceuticals, Terumo BCT, Guthy-Jackson Charitable Foundation (contracted research); *BrainStorm Therapeutics, Caladrius Biosciences, Celgene, MedImmune, Novartis, Ono Pharmaceutical (personal fees); Chugai Pharmaceutical (consulting fee); MedImmune, Viela Bio (a clinical trial adjudication committee).* *Romain Marignier:* Biogen, Merck Serono, Novartis, Roche, Sanofi Genzyme, Teva, Viela Bio (funding for travel and honoraria); *MedImmune, Viela Bio (scientific advisory board).* *Jeffrey L. Bennett:* AbbVie, Chugai, Clene Nanomedicine, Equilibrium, Frequency Therapeutics, Genentech, Genzyme, Alexion (personal fees); *Aquaporin (patent issued); EMD Serono, Novartis (grants and personal fees); MedImmune/Viela Bio (consulting fee); Guthy-Jackson Charitable Foundation, Mallinckrodt, National Institutes of Health (grants).* *Ho Jin Kim:* Alexion, Celltrion, Eisai, HanAll BioPharma, MedImmune/Viela Bio, Merck Serono, Novartis (personal fees); *Journal of Clinical Neurology and Multiple Sclerosis Journal (associate editor/co-editor); National Research Foundation of Korea (grants); Sanofi Genzyme, Teva-Handok (grants, personal fees).* *Brian Weinschenker:* Alexion, MedImmune, Viela Bio (royalties, payments); *Mitsubishi Tanabe Pharma (consulting fee); RSR Ltd, Oxford University, Hospices Civils de Lyon and MVZ Labor PD Dr Volkmann and Kollegen GbR (royalty).* *Sean J. Pittock:* Alexion, Autoimmune Encephalitis Alliance, Grifols, Euroimmun, NIH ROI NS065829-01, MedImmune/Viela Bio, Guthy-Jackson Charitable Foundation (consulting fees and research support); *named inventor on filed patents that relate to functional AQP4/NMO-IgG assays and NMO-IgG as a cancer marker (receipt of intellectual property rights/patent holder).* *Kazuo Fujihara:* Alexion, Bayer Schering, Biogen, Chugai, MedImmune, Merck Serono, Mitsubishi Tanabe Pharma, Nihon Pharmaceutical, Novartis, Ono, Viela Bio (scientific advisory boards); *Asahi Kasei Medical, Astellas, Bayer Schering, Biogen, Daiichi Sankyo, Dainippon Sumitomo, Eisai, Mitsubishi Tanabe Pharma, Nihon Pharmaceutical, Novartis, Takeda (funding for travel and speaker honoraria); Asahi Kasei Medical, Bayer Schering, Biogen, Chemo-Sero-Therapeutic Research Institute, Chugai, Genzyme Japan, Mitsubishi Tanabe Pharma, Nihon Pharmaceutical, Ono, Teijin, Teva, Ministry of Education, Culture, Sports, Science and Technology of Japan, Ministry of Health, Welfare and Labor of Japan (contracted research).* *Friedemann Paul:* Bayer, Biogen, Merck

Serono, Novartis, Sanofi Genzyme, Teva (research support, speaker honoraria, travel reimbursement); Guthy-Jackson Charitable Foundation (travel funding); Novartis (serves on steering committee); German Research Council (DFG Exc 257) and German Competence Network for Multiple Sclerosis (financial support). Gary Cutter: AMO Pharma, BioLineRx, Horizon Pharmaceuticals, Merck, Merck/Pfizer, National Heart, Lung, and Blood Institute (protocol review committee); National Institute of Child Health and Human Development (OPRU oversight committee); Neurim, OPKO Biologics, Orphazyme, Reata Pharmaceuticals, Receptos/Celgene, Sanofi-Aventis, Teva Pharmaceuticals (personal fees for participation on data and safety monitoring boards); Atara Biotherapeutics, Axon, Argenix, Biogen, Biotherapeutics, Brainstorm Cell Therapeutics, Charleston Laboratories Inc, Click Therapeutics, Genentech, Genzyme, GW Pharma, Klein Buendel Inc, MedDay, MedImmune, Novartis, Roche, SciFluor, Somahlution, Teva Pharmaceuticals, TG Therapeutics, UTHealth Houston (consulting fee); Pythagoras Inc, AL, USA (president of organization); Teva Neuroscience (grant); University of Alabama at Birmingham (salary). Ari Green: Bionure, Inception Sciences, JAMA Neurology, MedImmune/Viela Bio, Mylan, Synthron, Trims Pharma (other financial relationships [for activities as expert witness, associate editor, advisory board/steering committee participation and end point adjudication]); Conrad N. Hilton Foundation and Tom Sherak MS Hope Foundation (grants); Pipeline Therapeutics (personal fees from and other financial relationships). Orhan Aktas: Almirall, MedImmune, Merck Serono, Roche (personal fees). Bayer HealthCare, Biogen, Genzyme, Novartis, Teva, Viela Bio (grants and personal fees); German Research Foundation (DFG) and German Ministry of Education and Research (BMBF) (grants). Hans-Peter Hartung: Bayer HealthCare, Biogen, CSL Behring, GeNeuro, Genzyme, MedImmune, Merck Serono, Novartis, Opexa, Receptos, Roche, Sanofi, Teva, Viela Bio (consulting, speaking, serving on steering committees). Fred D. Lublin: AbbVie, Acorda, Actelion, Apitope, Atara Biotherapeutics, Bayer HealthCare, Biogen, BrainStorm Cell Therapeutics, EMD Serono, Forward Pharma, GW Pharmaceuticals, Innate Immunotherapeutics, Jazz Pharmaceuticals, MedDay, MedImmune/Viela Bio, Novartis, Orion Biotech, Polpharma, Receptos/Celgene, Roche/Genentech, TG Therapeutics (personal fees); Sanofi Genzyme, Teva Neuroscience (grants and personal fees); Transparency Life Sciences (grants). Maureen A. Mealy, Jorn Drappa, Gerard Barron, Soraya Madani, Dewei She, Daniel Cimborra, William Rees, John N. Ratchford, Eliezer Katz: Viela Bio (salary). Bruce A.C. Cree: Akili, Alexion, Biogen, GeNeuro, Novartis, Sanofi Genzyme, TG Therapeutics (consulting fee).

Keywords: Disease-modifying treatments in MS

(DXT16) Effectiveness of Delayed-Release Dimethyl Fumarate Relative to Duration of Prior Glatiramer Acetate in Patients Enrolled in the RESPOND Study

Derrick Robertson,¹ Pavle Repovic,² Stanley L. Cohan,³ Yang Mao-Draayer,⁴ Ray Su,⁵ James B. Lewin,⁶ Jenna Borowski⁶

¹USF Multiple Sclerosis Center, Tampa, FL; ²Multiple Sclerosis Center, Swedish Neuroscience Institute, Seattle, WA; ³Providence Multiple Sclerosis Center, Providence Brain Institute, Portland, OR; ⁴University of Michigan Medical School, Ann Arbor, MI; ⁵Biogen, Cambridge, MA; ⁶Biogen, Weston, MA

Background: Dimethyl fumarate (DMF) has demonstrated efficacy and a favorable benefit-risk profile in clinical studies of patients with relapsing-remitting multiple sclerosis (RRMS). RESPOND (trial registration: NCT01903291), a phase 4, 12-month study, evaluated outcomes in patients with RRMS prescribed DMF after suboptimal response to glatiramer acetate (GA). Patients may have a suboptimal response to a disease-modifying therapy such as GA early after treatment initiation, or in some cases, after several years. **Objectives:** To evaluate relapse and patient-reported outcomes (PROs) over 12 months in patients with RRMS who switched to DMF after suboptimal response to GA in real-world practice and to explore whether time on prior GA may influence response to DMF. **Methods:** RESPOND was conducted at 63 sites in the United States between August 2013 and February 2016. Patients diagnosed with RRMS with a suboptimal response to GA (perceived insufficient efficacy, intolerance, or poor adherence to GA) were enrolled. The median duration of prior GA was 36 months. This post hoc analysis compared treatment outcomes at 12 months following DMF initiation in subgroups of patients according to duration of prior GA treatment (≤ 36 vs > 36 months). **Results:** Among patients treated with GA for ≤ 36 months ($n = 177$) and > 36 months ($n = 141$), the ARR at 12 months prior to DMF was 0.588 (95% CI 0.49-0.70) and 0.369 (95% CI 0.28-0.49), whereas 12 months after switching to DMF, the ARR was 0.094 (95% CI 0.05-0.18) and 0.121 (95% CI 0.07-0.22), respectively (ARR reductions of 84% and 67%, respectively). The estimated proportion of patients relapsed (PPR) at month 12 on DMF was 6.5% for patients who had received prior GA for ≤ 36 months and 9.8% for patients who had received prior GA for > 36 months. PROs for quality of life, fatigue, disability, treatment satisfaction, work productivity, and depressive symptoms improved or

remained stable in both subgroups. We will also present outcomes for subgroups stratified by reason for discontinuation (insufficient efficacy vs other). **Conclusions:** Improvements in ARR and PPR were observed in patients who switched to DMF earlier (< 36 months prior GA treatment), and in patients who switched to DMF after being treated with GA for > 36 months. ARR and PPR were numerically lower in the group who switched earlier, but these differences were not statistically significant between the 2 subgroups. Both groups demonstrated improvement or stability across several PROs.

Supported by: Biogen

Disclosure: Derrick Robertson: Acorda (speakers' bureau); Actelion, MedDay, PCORI, TG Therapeutics (contracted research); Alexion, Celgene, Teva Neuroscience (consulting fee, speakers' bureau); Biogen, EMD Serono, Genentech, Novartis, Sanofi Genzyme (consulting fee, contracted research, speakers' bureau); Mallinckrodt (contracted research, speakers' bureau). Pavle Repovic: Alexion (consulting fee, speakers' bureau); Biogen, EMD Serono (consulting fee, contracted research, speakers' bureau); Celgene, Novartis, Sanofi Genzyme, Viela (consulting fee); Genentech (contracted research, speakers' bureau); Genzyme (speakers' bureau). Stanley L. Cohan: AbbVie, Adamas, Alexion, Alithios, MedDay, Pear Therapeutics, Roche Genentech, Sage Therapeutics (contracted research); Biogen, Novartis (consulting fee, contracted research, speakers' bureau); EMD Serono, Roche (consulting fee); Genentech (consulting fee, speakers' bureau); Sanofi Genzyme (consulting fee, contracted research); Sanofi Genzyme, Teva Neuroscience (speakers' bureau). Yang Mao-Draayer: Bayer Pharmaceutical, Celgene, EMD Serono, Genentech, Teva (consulting fee); Biogen, Sanofi Genzyme (consulting fee, contracted research); Chugai, National Institute Allergy Infectious Diseases Auto-immune Center of Excellence UMI-A1110557, NIH National Institute of Neurological Disorders and Stroke R01-NS080821, Novartis (contracted research). Ray Su, James B. Lewin: Biogen (ownership interest, salary). Jenna Borowski: Biogen (salary).

Keywords: Dimethyl fumarate, Disease-modifying treatments in MS

(DXT17) Long-term Follow-up Results from the Phase 2 Multicenter Study of Ublituximab (UTX), a Novel Glycoengineered Anti-CD20 Monoclonal Antibody, in Patients with Relapsing Multiple Sclerosis

Edward J. Fox,¹ Sibyl Wray,² Richard Shubin,³ Amy Lovett-Racke,⁴ Deren Huang,⁵ Ann D. Bass,⁶ Michael S. Weiss,⁷ Sean Power,⁷ Jenna Bosco,⁷ Koby Mok⁷

¹Central Texas Neurology Consultants and Dell Medical School, The University of Texas at Austin, Round Rock, TX; ²Hope Neurology, Knoxville, TN; ³Arcadia Neurology Center, Arcadia, CA; ⁴Ohio State University, Columbus, OH; ⁵Mount Carmel Neurology, Westerville, OH; ⁶Neurology Center of San Antonio, San Antonio, TX; ⁷TG Therapeutics, New York, NY

Background: Ublituximab (UTX) is a novel monoclonal antibody targeting a unique epitope on the CD20 antigen and glyco-engineered for enhanced B-cell targeting through antibody-dependent cellular cytotoxicity. Two phase 3 trials, ULTIMATE I and II, are fully enrolled and are investigating the efficacy and safety of UTX in relapsing forms of multiple sclerosis (RMS). **Objectives:** To evaluate the long-term safety and tolerability of UTX treatment in patients with RMS enrolled in the open-label extension (OLE) of a phase 2 trial. **Methods:** TG1101-RMS201 was a 52-week, phase 2, placebo-controlled, multicenter study of UTX in RMS. Subjects who completed RMS201 were eligible to continue treatment in the OLE, receiving 1-hour 450-mg UTX infusions every 24 weeks. **Results:** RMS201 enrolled 48 subjects and the primary end point was to evaluate B-cell depletion. Median B-cell depletion of $> 99\%$ was observed at week 4 and maintained at week 48. At week 48, key observations included: 100% reduction in T1-Gd-enhancing lesions; 10.6% mean decrease in T2 lesion volume; 93% of subjects relapse free, and an annualized relapse rate (ARR) of 0.07. UTX was well tolerated; the most common adverse event (AE) was infusion-related reactions (all grade 1-2). No discontinuations due to AEs were reported. UTX continues to be well tolerated, with a median duration of follow-up of 124.7 weeks, no drug-related discontinuations, and only 1 AE deemed at least possibly related to UTX that occurred in more than 1 patient, which was infusion-related reactions (IRR) all grade 1 or 2 in severity. At the time of presentation, long-term safety information will be presented for all patients on the OLE. **Conclusions:** The phase 2 OLE data support that UTX continues to be safe, well tolerated, and effective with 1-hour infusions. These results support the ongoing phase 3 ULTIMATE program in RMS.

Supported by: None

Disclosure: Edward J. Fox: Biogen, Celgene, Chugai, EMD Serono, MedDay (consulting fee); Genentech/Roche, Sanofi Genzyme, Teva (speakers' bureau) Novartis, TG Therapeutics (contracted research). Amy Lovett-Racke: Novartis, TG Therapeutics (contracted research). Sibyl Wray: Biogen, EMD Serono,

Genentech/Roche (speakers' bureau); Genzyme/Sanofi, Hope Neurology (consulting fee). Richard Shubin: Nothing to disclose. Deren Huang: Biogen, Novartis (consulting fee, speakers' bureau); Celgene, Genentech (consulting fee); Teva (speakers' bureau). Ann D. Bass: Actelion, Genentech-Roche, TG Therapeutics (contracted research); Biogen, EMD Serono, Novartis (speakers' bureau); Mallinckrodt, Sanofi Genzyme (consulting fee). Michael S. Weiss, Sean Power, Jenna Bosco, Koby Mok: TG Therapeutics (salary).

Keywords: Disease-modifying treatments in MS

(DXT18) Adherence and Compliance with Subcutaneous Administration of Ofatumumab in Relapsing Multiple Sclerosis

Edward Fox,¹ Lori Mayer,² Angela Aungst,³ Alexandra Goodyear,⁴ Cecile Kerloeguen,⁵ Linda Mancione,⁴ Nicola Rennie,⁵ Dee Stoneman,⁵ Martin Zalesak,⁵ Marina Ziehn,⁴ Derrick Robertson,⁶ Jeffrey A. Cohen⁷

¹Central Texas Neurology Consultants, Multiple Sclerosis Clinic of Central Texas, Round Rock, TX; ²MS Clinic of Central Texas, Central Texas Neurology Consultants, Round Rock, TX; ³Multiple Sclerosis Division Department of Neurology University of South Florida, Florida, FL; ⁴Novartis Pharmaceuticals Corporation, East Hanover, NJ; ⁵Novartis Pharma AG, Basel, Switzerland; ⁶College of Medicine, Neurology, University of South Florida, Tampa, FL; ⁷Department of Neurology, Mellen MS Center, Neurological Institute, Cleveland Clinic, Cleveland, OH

Background: Ofatumumab (OMB), the first fully human anti-CD20 monoclonal antibody, administered with a monthly 20-mg subcutaneous (s.c.) dosing regimen, demonstrated superior efficacy vs teriflunomide (TER) in the two phase 3 ASCLEPIOS I and ASCLEPIOS II trials in relapsing multiple sclerosis. Patients who completed the double-blind phase of the trials on study drug were eligible for transition to the ongoing open-label extension study ALITHIOS. **Objectives:** To evaluate treatment discontinuation and compliance with OMB and TER treatment in the phase 3 ASCLEPIOS I/II trials and to assess patients' acceptance of transitioning to the ALITHIOS study. **Methods:** In ASCLEPIOS I/II, patients were randomized (1:1) to OMB 20 mg s.c. (loading doses, administered at clinic: days 1, 7, and 14; maintenance doses, administered at home: every 4 weeks from week 4) or TER 14 mg (orally once daily), for up to 30 study months. Here we report on treatment discontinuation and compliance (defined as exposure to study drug [days]/on-treatment period [days] × 100%) in ASCLEPIOS trials and percentage of eligible ASCLEPIOS patients who accepted to transition to the ALITHIOS study and the compliance in this study. **Results:** In ASCLEPIOS I, 759/927 (81.9%) randomized patients (OMB: 400/465 [86.0%]; TER: 359/462 [77.7%]) completed the study on study drug. The proportion of patients discontinuing treatment were OMB, 14.0%; TER, 21.2%. The most common reasons for discontinuation (>2% in any group) were patient/guardian decision (OMB: 4.9%; TER: 8.2%), adverse event (OMB: 5.2%; TER: 5.0%), and physician decision (OMB: 2.2%; TER: 6.5%). In ASCLEPIOS II, 753/955 (78.8%) randomized patients (OMB: 383/481 [79.6%]; TER: 370/474 [78.1%]) completed the study on study drug. Proportion of patients discontinuing treatment were OMB, 20%; TER, 21.5%; reasons for discontinuation were patient/guardian decision (OMB: 7.3%; TER: 7.8%), adverse event (OMB: 5.6%; TER: 4.9%), and physician decision (OMB: 5.2%; TER: 6.8%). In both trials compliance was high (>95% of patients falling in the ≥90% compliance category) across treatment groups. Approximately 90% of eligible patients consented to participate in the open-label study; compliance data will be presented. **Conclusions:** In ASCLEPIOS trials compliance with home-administered s.c. OMB was high and fewer patients discontinued OMB as compared to TER. Most eligible patients accepted transition to the open-label ALITHIOS extension study.

Supported by: None

Disclosure: Edward Fox: Biogen, Celgene, Chugai, EMD Serono, Genentech/Roche, MedDay, Novartis, Sanofi, Genzyme, Teva, TG Therapeutics (advisory work, consulting fee, contracted research, speakers' bureau). Lori Mayer: Biogen, Novartis, Celgene, Genentech, EMD Serono (consulting fee, speakers' bureau). Angela Aungst: Nothing to disclose. Alexandra Goodyear, Cecile Kerloeguen: Novartis (salary). Linda Mancione: Novartis Pharmaceuticals Corporation (salary). Nicola Rennie, Dee Stoneman, Martin Zalesak, Marina Ziehn: Novartis Pharma AG (salary). Derrick Robertson: Biogen, Celgene, EMD Serono, Genentech, Novartis, Sanofi Genzyme, Teva, (consulting fee); Biogen, Celgene, EMD Serono, Genentech, Novartis, Sanofi Genzyme, Teva, Actelion, Mallinckrodt, MedDay, PCORI, TG Therapeutics (contracted research); Biogen, Celgene, EMD Serono, Genentech, Novartis, Sanofi Genzyme, Teva, Mallinckrodt, Accord (speakers' bureau). Jeffrey A. Cohen: Adamas, Convelo, Mylan, Population Council (consulting fee); Multiple Sclerosis Journal (co-editor).

Keywords: Compliance, Disease-modifying treatments in MS

(DXT20) Glatiramer Acetate (GA) Produced by Mapi Pharma Is Equivalent to Commercially Available GA Preparations

Susanna Tchilibon, Nadav Bleich Kimelman, Shai Rubnov, Laura Popper, Uri Danon, Ehud Marom

Mapi Pharma Ltd, Ness Ziona, Israel

Background: Glatiramer acetate (GA) is one of the first disease-modifying treatments approved for relapsing-remitting multiple sclerosis. Recently, several generic equivalents of GA were approved for marketing in the United States. Mapi is developing GA to be used in Glatiramer Acetate Depot (GA Depot) and in Mapi's generic GA. Here, we present key characteristic data from physicochemical (structural) and biological (pharmacodynamic) assays that were conducted to compare Mapi's GA and all US-approved GA equivalents. **Objectives:** To demonstrate through the results (physicochemical and biological) that Mapi's GA is similar to US-approved and commercially available GA equivalents. **Methods:** Mapi's GA is produced using the same starting materials and basic chemistry as Copaxone. At least 5 batches of Copaxone, 1 batch of Glatopa (Sandoz), 1 batch of Glatiramer Acetate Injection (Mylan), and several batches of Mapi's GA were analyzed for physicochemical properties (molecular weight distribution, impurities profile, amino-acid composition, and spectral fingerprint) and various structural signatures. A representative batch of Mapi's GA was compared with three different commercially available GA preparations using a bioassay test (myelin oligodendrocyte glycoprotein-induced EAE in mice). **Results:** The selected tests and data presented here represent a portion of a broader set of physicochemical and biological assays that were conducted. No differences were observed in the physicochemical properties or the structural signatures between Mapi's GA and all other US-approved GA equivalents. Equivalent pharmacodynamic activity of Mapi's GA to 3 commercially available GA preparations (Copaxone, Glatopa, and Glatiramer Acetate Injection) was demonstrated using MOG-induced EAE in mice. **Conclusions:** Mapi's GA is equivalent to commercially available GA products in physicochemical properties, structural signatures, and biological activity as demonstrated by bioassay and complies with the US Food and Drug Administration's guidance for generic GA. These results will support GA Depot and a new generic GA version commercialization by Mapi Pharma.

Supported by: None

Disclosure: Susanna Tchilibon, Nadav Bleich Kimelman, Shai Rubnov, Laura Popper, Uri Danon: Mapi Pharma Ltd (employer/employee relationship). Ehud Marom: Mapi Pharma Ltd (salary).

Keywords: Disease-modifying treatments in MS

(DXT22) Characterization of Incidence and Time-to-Recovery from Grade 3/4 Lymphopenia Lasting ≥6 Months in Patients with Multiple Sclerosis Treated with Cladribine Tablets

Gabriel Pardo,¹ Kottil Rammohan,² Thomas P. Leist,³ Julie Aldridge,⁴ Ciara Rossier,⁵ Axel Nolting,⁶ Matt Mandel,⁴ Andrew Galazka,⁵ Daniel Jones,⁷ Gavin Giovannoni⁸

¹Oklahoma Medical Research Foundation, Oklahoma City, OK; ²Department of Neurology, Multiple Sclerosis Center, University of Miami, Miami, FL; ³Jefferson University Hospital, Philadelphia, PA; ⁴EMD Serono Inc, Billerica, MA; ⁵Merck, Aubonne, Switzerland; ⁶Merck KGaA, Darmstadt, Germany; ⁷EMD Serono Inc, Rockland, MA; ⁸Blizard Institute of Cell and Molecular Science, London, United Kingdom

Background: Patients with multiple sclerosis (MS) treated with cladribine tablets (CTs) are expected to experience lymphopenia because of its mechanism of action (MOA), and transient mild-to-moderate lymphopenia has been observed in most patients. Given that a reduction in overall lymphocyte counts is part of the MOA of cladribine, further studies on the severity of lymphopenia are warranted. **Objectives:** To examine the effect of CT 3.5 mg/kg (CT3.5, cumulative over 2 years) on grade 3/4 lymphopenia lasting ≥6 continuous months postbaseline in subjects with baseline absolute lymphocyte count (ALC) within normal limits from the CT3.5 monotherapy oral cohort, defined as the safety populations of the phase 3 CLARITY, CLARITY Extension, and ORACLE-MS studies of MS, and long-term follow-up in the PREMIERE registry. **Methods:** In this post hoc analysis, combined data from 2 years of phase 3 studies and the follow-up PREMIERE registry were analyzed on the incidence of grade 3/4 lymphopenia (ALC <500/mm³) lasting ≥6 months. Time to an episode and time to recovery were also assessed. Recovery from grade 3/4 lymphopenia is defined as a return to grade ≤2 lymphopenia. **Results:** Of the 923 patients randomized to CT3.5, 891 (96.5%) had baseline ALC within normal limits (grade 0), and in this subpopulation, 218 (23.6%) had at least a single reading of grade 3/4 lymphopenia, and 33 (3.6%)

had grade 3/4 lymphopenia lasting ≥ 6 months (38 episodes). More patients with grade 3/4 lymphopenia lasting ≥ 6 months were female (81.8% vs 66.1%), had used ≥ 1 prior disease-modifying drug (33.3% vs 19.5%), and had more severe disease (≥ 9 T2 lesions [93.9% vs 88.1%], ≥ 1 relapse [75.8% vs 53.9%], higher median score on Expanded Disability Status Scale [3.0 vs 2.0]) at baseline vs the overall patients with baseline ALC within normal limits. Of the 33 CT3.5-treated patients with grade 3/4 lymphopenia lasting ≥ 6 months, more had the episode in year 2 than in year 1 of the core studies (64% vs 18%), with a median (Q1, Q3) time to first episode of 58.9 (51.1, 83.1) weeks. Of the 38 grade 3/4 lymphopenia (≥ 6 months) episodes, 27 (71.1%) lasted 24-48 weeks, and 11 (28.9%) lasted >48 weeks. Median (Q1, Q3) time to recovery from first grade 3/4 lymphopenia (≥ 6 months) episode was 36.3 (28.4, 66.3) weeks. **Conclusions:** The incidence of grade 3/4 lymphopenia lasting ≥ 6 months in patients treated with CT3.5 was low. Most episodes (71.1%) of grade 3/4 lymphopenia lasting ≥ 6 months resolved within 6 months to 1 year.

Supported by: None

Disclosure: *Gabriel Pardo:* Alexion, Biogen, Celgene, EMD Serono, Novartis, Roche/Genentech, Sanofi Genzyme (speaker honoraria and/or consulting fees); *AbbVie, Adamas, Alkermes, Biogen, EMD Serono, Roche/Genentech, Sanofi Genzyme, Novartis, Teva (research support to the institution).* *Kottil Rammohan:* AbbVie, Adamas, Alkermes, Roche/Genentech, Sanofi Genzyme, Teva (research support to institution); *Alexion, Celgene, Roche/Genentech, Sanofi Genzyme (consulting fee); Biogen, EMD Serono, Novartis (consulting fee, research support to the institution).* *Thomas P. Leist:* Acorda, Biogen, EMD Serono, Genzyme, Novartis, Roche/Genentech, Sanofi-Aventis, Teva Neurosciences (speakers' bureau). *Julie Aldridge, Matt Mandel, Daniel Jones:* EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). *Ciara Rossier, Andrew Galazka:* Merck KGaA, Aubonne, Switzerland (a business of Merck KGaA, Darmstadt, Germany) (salary). *Axel Nolting:* Merck KGaA (salary). *Gavin Giovannoni:* AbbVie, Actelion, Almirall, Atara Bio, Bayer Schering Pharma, Five Prime, GlaxoSmithKline, GW Pharma, Merck KGaA, Pfizer Inc, Protein Discovery Laboratories, Sanofi Genzyme, Teva Pharmaceuticals Industries Ltd, UCB, Vertex Pharmaceuticals (consulting fee); *Biogen, Ironwood, Merck and Co, Novartis (consulting fee, unrelated research support).*

Keywords: Disease-modifying treatments in MS, Immunology and MS

(DXT23) Disease-Modifying Therapy Landscape: An Evaluation of Cost and Care

Gail Bridges,¹ Douglas Mager,² Mary Dorholt,² Rochelle Henderson²

¹Accredo, Memphis, TN; ²Express Scripts Inc, St. Louis, MO

Background: Multiple sclerosis (MS) affects ~1 million people in the United States and is the fourth largest specialty drug spend. Over the years many disease-modifying therapies (DMTs) have come to market, resulting in increased drug choice and spend. Patients often struggle with the cost and chronicity of these therapies. **Objectives:** Evaluate DMT utilization, cost, adherence, and switching in a commercial MS population. **Methods:** We analyzed DMT use data of 34.2 million beneficiaries with a pharmacy benefit plan administered by a large pharmacy benefit manager for the 2-year period 2017-2018. Unit cost trend is defined as the rate of cost change due to inflation, discounts, drug mix, and member cost and is determined on a per-member-per-year (PMPY) basis. Adherent patients were defined as an average medication possession ratio of $\geq 80\%$. Switching occurred when an alternative DMT claim occurred after the index DMT drug claim. **Results:** DMT prevalence was 0.09% in 2018. PMPY spend for DMT to treat MS decreased 4.8% in 2018, driven by a 7.8% utilization decrease. Utilization trend was negative year over year. Branded medications Tecfidera (dimethyl fumarate), Gilenya (fingolimod), and Avonex (interferon beta-1a) account for more than 44% of DMTs prescribed in this class and each of these increased in unit cost (range 3.7%-6.0%). Utilization of Copaxone (glatiramer acetate) declined by 59.6%, influenced by generic availability. Generic DMT fill rate rose from 5.2% to 17.0% in year over year. In 2018, just over 27% of patients utilizing MS medications were nonadherent to their DMT. This represents a 1.4% year over year improvement in adherence. Nearly 15% of patients initiating a DMT switched to another medication within 1 year of the initial prescription. Utilization declined for injected interferons, such as Avonex, Rebif (interferon beta-1a), and Betaseron (interferon beta-1b), as market share shifts to oral medications such as Aubagio (teriflunomide), which had a 4.2% utilization increase in 2018. Not all new therapies are oral. Some of the newer DMT options are infrequently administered infused (IV) products, such as Ocrevus (ocrelizumab). Claims for IV DMT may adjudicate through the medical benefit and not be fully appreciated in this analysis. **Conclusions:** Knowledge of DMT trends can assist in reducing costs and identifying opportunities for further study. A significant

portion of patients with MS struggle with adherence. Early identification of nonadherent patients and application of customized support may help optimize therapeutic outcomes.

Supported by: None

Disclosure: *Gail Bridges, Douglas Mager, Mary Dorholt, Rochelle Henderson:* Express Scripts (salary).

Keywords: Adherence, Disease-modifying treatments in MS, Economic issues and MS

(DXT24) Two Expanded Disability Status Scale Subscales Evaluated in Patients with Relapsing-Remitting or Secondary Progressive Multiple Sclerosis

Gary Cutter,¹ Xiangyi Meng,² Jamie L. Weiss,² Ralph H.B. Benedict,³ Stanley L. Cohan,⁴ Bruce A.C. Cree,⁵ Wendy Su,² Florian P. Thomas^{6,7}

¹School of Public Health, University of Alabama, Birmingham, AL; ²Novartis Pharmaceuticals Corporation, East Hanover, NJ; ³University at Buffalo, State University of New York, Buffalo, NY; ⁴Providence Multiple Sclerosis Center, Providence Brain Institute, Portland, OR; ⁵Department of Neurology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA; ⁶Hackensack Meridian School of Medicine, Nutley, NJ; ⁷Department of Neurology, Multiple Sclerosis Center, Hackensack University Medical Center, Hackensack, NJ

Background: In phase 3 trials using the Expanded Disability Status Scale (EDSS), fingolimod and siponimod reduced disability worsening in patients with relapsing-remitting (RRMS) and secondary progressive multiple sclerosis (SPMS), respectively. However, EDSS components may differentially contribute to worsening, and contribution of each component may depend on disease stage. Also, EDSS assessments can be burdensome for patients and clinicians. Using factor analysis, different EDSS functions can be allocated to 2 novel subscales, including parameters most relevant to disease worsening. **Objectives:** Evaluate the effects of fingolimod and siponimod using EDSS subscales derived by factor analysis of phase 3 trial data. **Methods:** PROC FACTOR procedure was used to determine best fit of baseline EDSS data to the following subscales: Motor Integration (MI: ambulation, cerebellar, and pyramidal functions) and Colateral (C: bowel and bladder, brainstem, cerebral, sensory, and visual functions). Treatment effect sizes (ESs) on disability (mean change from baseline vs placebo) were determined overall and by each subscale up to 24 months (M) in patients with RRMS from FREEDOMS and up to 27M in patients with SPMS from EXPAND. Statistical significance was assessed using rank analysis of covariance (FREEDOMS) and a covariance mixed-effect, repeat-measurement model (EXPAND). Subgroup analyses of patients in EXPAND with/without lesion activity or relapses prior to screening were also performed and will be presented. Analyses were for hypothesis generation without multiple comparison adjustment. **Results:** Treatment ESs in FREEDOMS (N = 843: fingolimod, n = 425; placebo, n = 418) at 24 months were -0.14, P = .002 (EDSS); -0.18, P = .002 (MI); and -0.07, P = .081 (C). Significant effects (P < .05) were seen on MI from M6; effects on C were mostly nonsignificant. In EXPAND (N = 1645: siponimod, n = 1099; placebo, n = 546), overall treatment effects were detected over 27M for EDSS (P = .020), MI (P = .014), and C (P = .021). Significant ESs were seen on MI (all P < .01) at M9 (-0.28), M15 (-0.34), and M18 (-0.34), and on C at M18 (-0.24, P < .05) and M27 (-0.54, P < .001). **Conclusions:** The benefits of fingolimod in patients with RRMS mainly affected the MI subscale. For patients with SPMS, the benefits of siponimod were seen on both MI and C. In SPMS, effects on MI appeared earlier than on C, however the largest ES was seen later on C. Defining EDSS subscales with factor analyses may help improve their clinical usefulness.

Supported by: None

Disclosure: *Gary Cutter:* Avexis Pharmaceuticals, BioLineRx, Brainstorm Cell Therapeutics, CSL Behring, Galmed Pharmaceuticals, Hisun Pharmaceuticals, Horizon Pharmaceuticals, Merck, Merck/Pfizer, Neurim, National Heart, Lung, and Blood Institute (protocol review committee); National Institute of Child Health and Human Development (OPRU oversight committee); Ophazyme, Opko Biologics, Reata Pharmaceuticals, Receptos/Celgene, Sanofi-Aventis, Teva Pharmaceuticals, Vivus (data and safety monitoring boards); *Biogen, Click Therapeutics, Genentech, Genzyme, Gilgamesh Pharmaceuticals, GW Pharmaceuticals, Klein Buendel Incorporated, MedDay, MedImmune, Osmotica Pharmaceuticals, Perception Neurosciences, Recursion Pharmaceuticals, Roche, Somablation, TG Therapeutics (consulting fee); Novartis (consulting fee, data and safety monitoring boards); Pythagoras, Inc (president); University of Alabama at Birmingham (salary). Xiangyi Meng, Jamie L. Weiss, Wendy Su:* Novartis Pharmaceuticals Corporation (salary). *Ralph H.B. Benedict:* Acorda, Mallinckrodt, National Institutes of Health, National MS Society (contracted research); *Biogen (CME speaking fees, consulting fee, contracted research); Celgene, EMD Serono (CME speaking fees, consulting fee); Genentech, Genzyme, Novartis (consulting*

fee, contracted research); Psychological Assessment Resources, Inc (royalty); Roche, Sanofi, Takeda, Verasci (consulting fee). **Stanley L. Coban:** AbbVie, Adamas, Alexion, MedDay, Sage Bionetworks (contracted research); Biogen, Novartis, Sanofi Genzyme (consulting fee, contracted research, speakers' bureau); EMD Serono, Pear Therapeutics (consulting fee); Genentech, Roche (contracted research, speakers' bureau). **Bruce A.C. Cree:** Akili, Alexion, Atara, Biogen, EMD Serono, Novartis, TG Therapeutics (consulting fee). **Florian P. Thoms:** Biogen, Teva Neuroscience (contracted research); Genentech, Novartis, Sanofi Genzyme (consulting fee).

Keywords: Disease-modifying treatments in MS

(DXT26) Long-Term Disease Stability Assessed by the Expanded Disability Status Scale in Patients Treated with Cladribine Tablets in the CLARITY and CLARITY Extension Studies

Gavin Giovannoni,¹ Giancarlo Comi,² Kottil Rammohan,³ Peter Rieckmann,⁴ Patrick Vermersch,⁵ Fernando Dangond,⁶ Birgit Keller,⁷ Dominic Jack⁷

¹Queen Mary University of London, London, United Kingdom; ²Department of Neurology and Institute of Experimental Neurology, Università Vita-Salute San Raffaele, Milan, Italy; ³Department of Neurology, Multiple Sclerosis Center, University of Miami, Miami, FL; ⁴Department of Neurology, Medical Park Loipl, and University of Erlangen, Erlangen, Germany; ⁵University of Lille, Lille, France; ⁶EMD Serono Inc, Billerica, MA; ⁷Merck KGaA, Darmstadt, Germany

Background: Treatment with cladribine tablets 10 mg (cumulative dose 3.5 mg/kg [CT3.5] over 2 years) in CLARITY and CLARITY Extension reduced relapse rate and slowed disability progression vs placebo in patients with relapsing-remitting multiple sclerosis (RRMS). **Objectives:** The objective of this post hoc analysis was to evaluate long-term disease stability assessed by the Expanded Disability Status Scale (EDSS) after treatment with CT3.5 in patients with RRMS enrolled in CLARITY and CLARITY Extension. **Methods:** Patients randomized to CT3.5 in CLARITY and then randomized to placebo in CLARITY Extension, with at least 1 postbaseline EDSS measurement, were included (CP3.5; n = 98). This analysis assessed EDSS score over time (from CLARITY randomization to end of follow-up in CLARITY Extension, including the bridging interval between studies) at 6-monthly intervals, and separately time to 3- and 6-month confirmed EDSS score progression from CLARITY baseline. EDSS score worsening or improvement in each year was defined as any increase or decrease, respectively, in minimum EDSS score at 6-monthly intervals; all other cases were classified as stable. An increase or decrease was defined as an EDSS score change of 1.5 points (baseline EDSS 0), 1 point (baseline EDSS ≤4.5), or 0.5 point (baseline EDSS ≥5.0). **Results:** Five years after CLARITY baseline, median EDSS remained stable compared with baseline values. Median EDSS score (95% CI) for patients in the CP3.5 group was 2.5 (2.0-3.5) compared with 3.0 (2.5-3.5) at baseline. In each 12-month period, EDSS score stability was observed in >50% of patients, improvement in 21%-30% of patients, and worsening in 0%-25%. During year 5 in the CP3.5 group, EDSS stability was observed in 53.9% of patients, improvement in 21.3%, and worsening in 24.7%. Less than 30% of patients reached 3- or 6-month confirmed EDSS progression by year 5. **Conclusions:** EDSS score was stable up to 5 years post-CLARITY baseline for the CP3.5 group. Between 20%-30% of patients demonstrated improvement in EDSS score vs baseline each year.

Supported by: None

Disclosure: **Gavin Giovannoni:** AbbVie, Actelion, Almirall, Atara Bio, Bayer Schering Pharma, Five Prime, GlaxoSmithKline, GW Pharma, Merck KGaA, Pfizer Inc, Protein Discovery Laboratories, Sanofi Genzyme, Teva Pharmaceuticals Industries Ltd, UCB, Vertex Pharmaceuticals (consulting fee); Biogen, Ironwood, Merck and Co, Novartis (consulting fee, unrelated research support). **Giancarlo Comi:** Almirall SpA, Biogen, Biogen Italia Srl, Celgene Group, EXCEMED, F. Hoffman-La Roche, Forward Pharma, Genzyme Corporation, Genzyme Europe, MedDay, Merck KGaA, Merck Serono SpA, Novartis, Roche SpA, Sanofi Genzyme, Teva Italia Srl, Teva Pharmaceutical Industries Ltd (consulting fee). **Kottil Rammohan:** Acorda, Biogen, EMD Serono, Genzyme, Novartis, Roche/Genentech, Sanofi-Aventis, Teva Neurosciences (speakers' bureau). **Peter Rieckmann:** Bayer Schering Pharma, Biogen, Boehringer-Ingelheim, Genzyme, Merck, Novartis, Sanofi-Aventis, Serono Symposia International Foundation, Teva Pharmaceutical Industries (speakers' bureau). **Patrick Vermersch:** Almirall, Celgene, Novartis, Roche (consulting fee); Bayer, Biogen, Merck KGaA, Sanofi Genzyme (consulting fee, research support). **Fernando Dangond:** EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). **Birgit Keller, Dominic Jack:** Merck KGaA (salary).

Keywords: Disease-modifying treatments in MS

(DXT27) Integrated Lymphopenia Analysis in Younger and Older Patients with Multiple Sclerosis Treated with Cladribine Tablets

Gavin Giovannoni,¹ Patricia K. Coyle,² Patrick Vermersch,³ Bryan Walker,⁴ Julie Aldridge,⁵ Axel Nolting,⁶ Sana Syed,⁵ Andrew Galazka,⁷ Daniel Jones,⁸ Thomas P. Leis⁹

¹Queen Mary University of London, London, United Kingdom; ²Stony Brook University, Stony Brook, NY; ³University of Lille, Lille, France; ⁴Neurological Department, Duke University School of Medicine, Durham, NC; ⁵EMD Serono Inc, Billerica, MA; ⁶Merck KGaA, Darmstadt, Germany; ⁷Merck, Aubonne, Switzerland; ⁸EMD Serono Inc, Rockland, MA; ⁹Jefferson University Hospital, Philadelphia, PA

Background: The mechanism of action of cladribine is thought to be related to a reduction in lymphocyte counts. As the immune system undergoes changes with aging, it is not currently well understood whether the impact of cladribine tablets (CTs) on lymphocyte counts differs in younger vs older patients. **Objectives:** To examine the effect of age (≤50 and >50 years) at baseline and after treatment with placebo or CT 3.5 mg/kg (CT3.5, cumulative over 2 years) on lymphopenia in patients enrolled in the phase 3 CLARITY, CLARITY Extension, and ORACLE-MS studies, and long-term follow-up in the PREMIERE registry. **Methods:** In this post hoc analysis, combined data from 2 years of phase 3 studies on levels of absolute lymphocyte count (ALC), incidence of grade 3/4 lymphopenia, and time to recovery from severe lymphopenia were analyzed in patients by baseline age group. **Results:** This analysis was carried out in 1564 patients: age ≤50: placebo N = 566, CT3.5 N = 813; age >50: placebo N = 75, CT3.5 N = 110. In both age groups, CT3.5 treatment resulted in a 43.1%-47.6% reduction in mean ALC vs placebo at week 9, which was after completing the year-1 doses. Mean (SD) ALC in the CT3.5 groups at week 9 were: age ≤50: 1.12 (0.50) × 10⁹/L; age >50: 1.00 (0.42) × 10⁹/L. ALC levels in the CT3.5-treated groups gradually increased thereafter up to week 48 [mean (SD) ALC at week 48: age ≤50: 1.27 [0.45] × 10⁹/L; age >50: 1.32 [0.51] × 10⁹/L]; ALC remained 34.5%-35.6% below those treated with placebo. In year 2 at week 55, CT3.5 treatment resulted in a 54.6%-60.8% reduction in mean ALC vs placebo (year 2 doses completed). Mean (SD) ALC in the CT3.5 groups at week 55 were: age ≤50: 0.89 (0.39) × 10⁹/L; age >50: 0.80 (0.33) × 10⁹/L. ALC levels in the CT3.5-treated groups gradually increased thereafter up to week 96 [mean (SD) ALC at week 96: age ≤50: 1.11 [0.42] × 10⁹/L; age >50: 1.11 [0.37] × 10⁹/L]; ALC remained 42.8%-43.7% below those treated with placebo. Incidence of grade 3/4 lymphopenia was higher with CT3.5 vs placebo in year 1 (age ≤50: 8.3%; age >50: 10.0%; vs 0%-0.4% in placebo) and year 2 (age ≤50: 18.7%; age >50: 20.0%; vs 0%-0.2% in placebo). Median time to recovery from grade 3/4 to ≤2 lymphopenia was 1.18 and 1.54 months for CT3.5-treated patients in the age ≤50 and >50 groups, respectively. **Conclusions:** ALC changes in CT3.5-treated patients were similar in older and younger patients relative to placebo-treated patients during 2 years of active treatment as expected. Recovery time from lymphopenia was also similar between the age groups.

Supported by: None

Disclosure: **Gavin Giovannoni:** AbbVie, Actelion, Almirall, Atara Bio, Bayer Schering Pharma, Five Prime, GlaxoSmithKline, GW Pharma, Merck KGaA, Pfizer Inc, Protein Discovery Laboratories, Sanofi Genzyme, Teva Pharmaceuticals Industries Ltd, UCB, Vertex Pharmaceuticals (consulting fee); Biogen, Ironwood, Merck and Co, Novartis (consulting fee, unrelated research support). **Patricia K. Coyle:** Accordant, Acorda Therapeutics, Bayer HealthCare, Biogen, Celgene, EMD Serono, Sanofi Genzyme, Teva Pharmaceuticals USA (consulting fee); Actelion, Alkermes, MedDay, National Institute of Neurological Disorders and Stroke (contracted research); Genentech/Roche, Novartis (consulting fee, contracted research). **Patrick Vermersch:** Almirall, Celgene, Novartis, Roche (consulting fee); Biogen, Merck KGaA, Sanofi Genzyme, Teva (consulting fee, research support). **Bryan Walker:** Biogen, Celgene, EMD Serono, Novartis, Sanofi Genzyme (consulting fee). **Julie Aldridge, Sana Syed, Daniel Jones:** EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). **Axel Nolting:** Merck KGaA (salary). **Andrew Galazka:** Merck KGaA, Aubonne, Switzerland (a business of Merck KGaA, Darmstadt, Germany) (salary). **Thomas P. Leis:** Alkermes, Bayer, Biogen, EMD Serono, Genentech, Genzyme, Novartis (consulting fee).

Keywords: Disease-modifying treatments in MS, Immunology and MS

(DXT28) Effectiveness of Cladribine Tablets in Patients with Relapsing-Remitting Multiple Sclerosis with Baseline Expanded Disability Status Scale Score ≥3.5 or ≤3.0 in CLARITY

Giancarlo Comi,¹ Gabriel Pardo,² Fernando Dangond,³ Julie Aldridge,³ Caroline Lemieux,⁴ Kottil Rammohan⁵

¹Department of Neurology and Institute of Experimental Neurology, Università Vita-Salute San Raffaele, Milan, Italy; ²Oklahoma Medical Research Foundation, Oklahoma City, OK; ³EMD Serono Inc, Billerica, MA; ⁴EMD Serono Inc, Mississauga, ON, Canada; ⁵Department of Neurology, Multiple Sclerosis Center, University of Miami, Miami, FL

Background: In the phase 3 CLARITY study, patients with relapsing-remitting multiple sclerosis (MS) treated with cladribine tablets (CTs) 10 mg (3.5 mg/kg [CT3.5] or 5.25 mg/kg cumulative dose over 2 years), showed significant reductions in annualized relapse rate ($P < .001$), time to 3-month (mo) sustained change in Expanded Disability Status Scale (EDSS) score ($P \leq .03$), and lesion activity on brain magnetic resonance imaging (MRI), all $P < .001$ vs placebo. However, the efficacy of CTs has not been fully characterized in patients transitioning to active secondary progressive MS or not, for which EDSS scores of ≥ 3.5 and ≤ 3.0 , respectively, can be used as a proxy definition. **Objectives:** To examine differences between placebo and CT3.5 on clinical and MRI end points and in attainment of no evidence of disease activity (NEDA) in patients with baseline EDSS scores of ≥ 3.5 or ≤ 3.0 in CLARITY. **Methods:** In this post hoc analysis, week 96 data from CLARITY were retrospectively examined across patients with baseline EDSS score ≥ 3.5 or ≤ 3.0 for relapses, 3- or 6-mo confirmed disability progression (CDP, per EDSS score changes), new T1 gadolinium-enhancing (Gd+) lesions, active T2 lesions, and NEDA. **Results:** Baseline characteristics were evenly distributed across treatment groups. Relapse, T1 Gd+ lesion, and T2 lesion numbers were greater in placebo-treated vs CT3.5-treated patients for both baseline EDSS groups (all $P < .0001$, nominal significance) at week 96. For patients with baseline EDSS score ≥ 3.5 , CT3.5 treatment resulted in improvements in qualifying relapses (Kaplan-Meier estimates at last event: 78.3% vs 60.3%), 3-month CDP (83.5% vs 69.4%), and 6-month CDP (88.1% vs 78.2%) vs placebo. Differences between CT3.5 and placebo treatment in the baseline EDSS ≤ 3.0 group were: 81.1% vs 61.6% in qualifying relapse, 86.3% vs 80.6% in 3-month CDP, and 92.0% vs 87.2% in 6-month CDP. Odds ratios (ORs) favored CT3.5 vs placebo for NEDA based on either 3-month (OR: 4.40) and 6-month CDP (OR: 4.11) in the baseline EDSS ≥ 3.5 group and in the baseline EDSS ≤ 3.0 group (OR for 3-month CDP: 4.23; OR for 6-month CDP: 4.62; all $P < .0001$, nominal significance). **Conclusions:** CT treatment resulted in similar improvements in relapse and MRI outcomes regardless of patient baseline EDSS score. The effect of CTs on NEDA composites was also favorable across patients with baseline EDSS score ≥ 3.5 or ≤ 3.0 .

Supported by: None

Disclosure: Giancarlo Comi: Ammirall SpA, Biogen, Biogen Italia Srl, Celgene Group, EXCEMED, F. Hoffman-La Roche, Forward Pharma, Genzyme Corporation, Genzyme Europe, MedDay, Merck KGaA, Merck Serono SpA, Novartis, Roche SpA, Sanofi Genzyme, Teva Italia Srl, Teva Pharmaceutical Industries Ltd (consulting fee). Gabriel Pardo: AbbVie, Adamas, Alkermes, Sanofi Genzyme, Teva (research support); Alexion, Celgene, Sanofi Genzyme (consulting fee); Biogen, EMD Serono, Novartis, Roche/Genentech (consulting fee, research support). Fernando Dangond, Julie Aldridge, Caroline Lemieux: EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany). Kottil Rammohan: Acorda, Biogen, EMD Serono, Genzyme, Novartis, Roche/Genentech, Sanofi-Aventis, Teva Neurosciences (speakers' bureau).

Keywords: Disease-modifying treatments in MS

(DXT29) ACAPELLA: Real-World Experience with Ocrelizumab: An Observational Study Evaluating Safety in Patients with Relapsing and Progressive Multiple Sclerosis, Year 3 Data

Hannah M Geils, India C. Stribling, Joshua D. Katz, Ellen S. Lathi
The Elliot Lewis Center for Multiple Sclerosis Care, Wellesley, MA

Background: Ocrelizumab (OCR) is a humanized, monoclonal antibody targeting CD20+ B-cells and is approved for the treatment of relapsing-remitting (RRMS) and primary progressive MS (PPMS). The ACAPELLA trial is a prospective study with a primary objective of assessing OCR-associated adverse events in a real-world MS population. ACAPELLA includes patients with preexisting conditions exempted from the phase 2 and 3 clinical trials, such as a prior history of malignancy, prior immunosuppressive treatment, and more advanced age and/or disability. Interim data analyses occur on a biannual basis and findings are reported yearly. This is the third iteration. **Objectives:** We sought to determine the frequency of serious infections and malignancy in a real-world population receiving OCR with characteristics outside the inclusion parameters of the phase 2 and 3 trials. **Methods:** The study includes all subjects treated with OCR at the Elliot Lewis Center since its commercial release in March 2017. Initial assessments include EDSS, brain magnetic resonance imaging (MRI), mammograms (standard of care), collection of medical history including prior serious or recurrent infections, history of malignancy and

exposure to immunosuppressive treatment, JC virus index, and CD19 count. **Results:** As of December 2019, 291 subjects were enrolled, 181 subjects had reached 12 months of treatment, 131 subjects had reached 18 months, and 84 subjects had reached 24 months. Subjects were 29% male, 71% female, with an age range of 18-73. Sixty-three percent had RRMS and 37% PMS (PPMS and progressive RRMS), with an EDSS score range of 0-7.5; 25% had a baseline EDSS score of ≥ 6.0 with a median of 3.0. The rate of infections for all OCR-treated patients was 43% (6% bronchitis, 5% shingles, 56% URIs, 30% urinary tract infections, 9% zoster, and 25% other infections). Four percent of subjects had a serious infection (one that required hospitalization, was felt to be life-threatening, or resulted in death). Three percent of subjects had clinical or MRI relapses. 8% of subjects had a history of prior neoplasm (excluding basal cell carcinoma). Two malignancies have occurred during OCR treatment. **Conclusions:** Thus far, the incidence of adverse events is comparable to that seen in the phase 3 trials and in previously reported ACAPELLA data. Additional topics of interest in the ACAPELLA population include the effect of continued OCR dosing on JC virus index values and immunoglobulin levels, and changes in EDSS and MRI over time.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Disease-modifying treatments in MS, Ocrelizumab

(DXT30) ACAPELLA: Hypogammaglobulinemia and JC Virus Status in Ocrelizumab-Treated Patients, Year 2 Data

Hannah M. Geils, India C. Stribling, Joshua D. Katz, Ellen S. Lathi
The Elliot Lewis Center for Multiple Sclerosis Care, Wellesley, MA

Background: Ocrelizumab (OCR) is a humanized anti-CD20 monoclonal antibody approved for the treatment of relapsing-remitting and primary progressive multiple sclerosis (MS). Immunoglobulin levels were monitored during the phase 3 trials, and 1.5% of patients developed low immunoglobulin G (IgG) values after 2-3 years of OCR treatment, potentially increasing the risk of infections. The JC virus (JCV) antibody index used to stratify PML risk in patients treated with natalizumab was not studied and the impact of long-term B-cell suppression on JCV and IgG titers is unknown. **Objectives:** As part of the ACAPELLA trial, a prospective study with a primary objective of assessing OCR-associated adverse events in a real-world MS population, we sought to evaluate the impact of OCR treatment on immunoglobulin levels and JCV titers over time. **Methods:** The study includes all subjects receiving OCR at the Elliot Lewis Center followed prospectively since March 2017. Subjects are monitored for the occurrence of infections and other serious adverse events and have biannual assessments of serum immunoglobulin levels and JCV antibody titers. **Results:** As of December 2019, 291 patients have been treated with OCR and enrolled in ACAPELLA: 181 have been treated for at least 12 months, 131 have been treated for at least 18 months, and 84 subjects have reached 24 months. Two hundred eighty-one of the total 291 subjects had IgG levels drawn at baseline. Twenty-seven subjects (10%) had IgG levels below the lower limit of normal at baseline. Of the 27 patients with low IgG at baseline, 19 have received treatment for at least 12 months. Of those 19, 4 patients were seen to have a $>10\%$ drop in IgG level after 12 months. Ten patients developed at least 1 low IgG level after 12-24 months of treatment exposure, although many returned to normal. Of the total 291 patients, 281 had a baseline JCV index. Ninety-three (33%) had titers <0.4 , 73 (26%) between 0.4-1.5, and 115 (41%) >1.5 . In our 2-year data, 3 patients had a change in JCV status from positive to negative between 12 and 24 months of treatment duration. Year-3 data are characterized in the poster. **Conclusions:** The frequency of persistent hypogammaglobulinemia was low in this cohort of patients and thus far has not been associated with an increased risk of infection. Three patients had a change in JCV status from positive to negative, and the effect of JCV index in the remaining subjects is further characterized.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Disease-modifying treatments in MS, Immunology and MS, Ocrelizumab

(DXT31) Impact of Eculizumab on Hospitalization Rates and Relapse Treatment in Patients with Neuromyelitis Optica Spectrum Disorder: Phase 3 PREVENT Study

Ho Jin Kim,¹ Sean J. Pittock,² Achim Berthele,³ Kazuo Fujihara,^{4,5,6} Michael Levy,^{7,8} Jacqueline Palace,⁹ Ichiro Nakashima,^{5,10} Murat Terzi,¹¹ Natalia Totolyan,¹² Shanthi Viswanathan,¹³ Kai-Chen Wang,^{14,15} Amy Pace,¹⁶ Kenji P. Fujita,¹⁶ Marcus Yountz,¹⁶ Robin Armstrong,¹⁶ Dean Wingerchuk¹⁷

¹National Cancer Center, Goyang, Korea, Republic of (South); ²Mayo Clinic, Rochester, MN; ³Technical University of Munich, Munich, Germany; ⁴Southern TOHOKU Research Institute for Neuroscience (STRINS), Koriyama, Japan; ⁵Tohoku University, Sendai, Japan; ⁶Fukushima Medical University, Fukushima City, Japan; ⁷Massachusetts General Hospital and Harvard Medical School, Boston, MA; ⁸Johns Hopkins University, Baltimore, MD; ⁹John Radcliffe Hospital, Oxford, United Kingdom; ¹⁰Tohoku Medical and Pharmaceutical University, Sendai, Japan; ¹¹Ondokuz Mayıs University, Samsun, Turkey; ¹²First Pavlov State Medical University of St. Petersburg, St. Petersburg, Russian Federation; ¹³Kuala Lumpur Hospital, Kuala Lumpur, Malaysia; ¹⁴National Yang Ming University, Taipei, Taiwan; ¹⁵Cheng-Hsin General Hospital, Taipei, Taiwan; ¹⁶Alexion Pharmaceuticals, Boston, MA; ¹⁷Mayo Clinic, Scottsdale, AZ

Background: Relapses resulting in hospitalization are common in patients with the rare autoimmune inflammatory disease neuromyelitis optica spectrum disorder (NMOSD). The randomized, double-blind, placebo-controlled PREVENT study (trial registration: NCT01892345) assessed the safety and efficacy of eculizumab in aquaporin-4 immunoglobulin G (AQP4-IgG)-positive NMOSD. Eculizumab significantly reduced the risk of adjudicated relapse compared with placebo (primary end point). **Objectives:** To evaluate rates of relapse-related hospitalization and associated treatment in patients with AQP4-IgG-positive NMOSD receiving eculizumab vs placebo in the PREVENT study. **Methods:** Patients with AQP4-IgG-positive NMOSD were randomized 2:1 to receive eculizumab (maintenance dose, 1200 mg/2 weeks, n = 96) or placebo (n = 47) with/without stable-dose concomitant immunosuppressant therapy (excluding rituximab and mitoxantrone). Hospitalizations were recorded as a component of the adverse event tracking performed throughout the study. The annualized relapse-related hospitalization and treatment rates were defined as the total number of relapses requiring hospitalization, and associated acute treatments, respectively, divided by the total number of patient-years in the study period. **Results:** The median exposure to treatment was 89.43 weeks for eculizumab and 41.29 weeks for placebo. The overall annualized hospitalization rates were 0.26 and 0.78 ($P < .0001$) in the eculizumab and placebo groups, respectively. The annualized relapse-related hospitalization rate was significantly lower in the eculizumab group than in the placebo group: 0.04 vs 0.31, respectively ($P < .0001$). The annualized relapse-related use of intravenous methylprednisolone, plasma exchange, and high-dose oral corticosteroids for eculizumab vs placebo was 0.07 vs 0.42 ($P < .0001$), 0.02 vs 0.19 ($P = .0001$), and 0.04 vs 0.11 ($P = .0733$), respectively. **Conclusions:** Treatment with eculizumab significantly reduced relapse-related hospitalizations and their associated treatment rates in patients with AQP4-IgG-positive NMOSD vs placebo, which may have a favorable effect on health-resource utilization.

Supported by: None

Disclosure: *Ho Jin Kim:* Alexion Pharmaceuticals, Celltrion, Eisai, HanAll BioPharma, Merck Serono, Novartis, Sanofi Genzyme, Teva-Handok, Viela Bio (consulting fee); *Journal of Clinical Neurology* (associate editor); *MedImmune/Viela Bio* (steering committee); *Multiple Sclerosis Journal* (co-editor). *Sean J. Pittock:* Alexion Pharmaceuticals (all compensation paid directly to Mayo Clinic, consulting fee, contracted research); *Astellas* (all compensation paid directly to Mayo Clinic, consulting fee); *Autoimmune Encephalitis Alliance* (all compensation paid directly to Mayo Clinic, contracted research); *Grifols* (all compensation paid directly to Mayo Clinic, contracted research); *MedImmune* (all compensation paid directly to Mayo Clinic, consulting fee, contracted research); *UCB* (consulting fee, personal compensation for attending the UCB advisory board meeting in Stockholm, Sweden, on Sept 10, 2019). *Achim Berthele:* Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent, speakers' bureau). *Kazuo Fujihara:* Alexion Pharmaceuticals, Biogen, Chugai, Mitsubishi-Tanabi, Novartis (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent); *Asahi Medical, Bayer, Eisai, Roche, Teijin* (fees for non-CME/CE services received directly from commercial interest or its agent). *Michael Levy:* Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent); *Genentech, Quest Diagnostics, Viela Bio* (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent). *Jacqueline Palace:* Biogen, Chugai (contracted research); *LEK, Viela Bio, Guidepoint* (consulting fee); *Merck Serono* (meeting/lecture/workshop participation); *Novartis, Roche, Argenx* (speakers' bureau); *UCB, Viela Bio, Roche* (conference/lecture participation). *Ichiro Nakashima:* Alexion (consulting fee). *Murat Terzi, Shanthi Viswanathan, Kai-Chen Wang:* Nothing to disclose. *Natalia Totolyan:* Alexion, Janssen, Novartis, Roche, Sanofi, Receptos Inc, Biocad (Russia) (contracted research); *Merck* (consulting fee); *Roche, Sanofi* (fees for lectures). *Amy Pace, Marcus Yountz, Roisin Armstrong:* Alexion Pharmaceuticals (ownership interest, salary). *Kenji P. Fujita:* Alexion Pharmaceuticals, Ablynam Pharmaceuticals (ownership interest, salary). *Dean Wingerchuk:* MedImmune, Novartis, Biogen, Celgene, Genentech, TG Therapeutics, Arcus

Medica, Reistone (consulting fee); *Terumo BCT, Alexion Pharmaceuticals* (fees for non-CME/CE services received directly from commercial interest or its agent).

Keywords: Hospitalization in NMOSD

(DXT33) ACAPELLA: B-Cell Reconstitution in Ocrelizumab-Treated Patients

India C. Sribling, Hannah M. Geils, Joshua D. Katz, Ellen S. Lathi
The Elliot Lewis Center for Multiple Sclerosis Care, Wellesley, MA

Background: Ocrelizumab (OCR) is a humanized anti-CD20 monoclonal antibody approved for the treatment of relapsing remitting (RRMS) and primary progressive multiple sclerosis (PPMS). In the OPERA trials, circulating CD19+ B-cell counts dropped to 0 within 14 days of OCR infusion. Median time to repletion, defined as >79 cells/ μ L, was 72 (range 27-175) weeks. Up to 5% of patients showed B-cell repletion during treatment. We sought to determine the frequency of patients on OCR who have significant B-cell reconstitution at the time of their next 6-month dose, and to determine if there is a correlation between early B-cell reconstitution and disease breakthrough or adverse events (AEs). **Objectives:** As part of the ACAPELLA trial, a prospective study with a primary objective of assessing OCR-associated AEs in a real-world multiple sclerosis population, we sought to evaluate the frequency and duration of early B-cell reconstitution and its relationship to disease activity and AEs. **Methods:** All subjects receiving OCR at the Elliot Lewis Center since March 2017 who consented to participate had serum immunoglobulin levels, JC virus antibody titers, and lymphocyte subsets on the day of each infusion prior to receiving OCR. Subjects were followed up prospectively and monitored for the occurrence of infections and other serious AEs. **Results:** As of December 2019, 291 patients had been treated with OCR and enrolled in ACAPELLA: 181 had been treated for at least 12 months, 131 had been treated for 18 months, and 84 subjects had reached 24 months. Of the 291 subjects, 207 had CD19 values drawn at an infusion. One hundred eighteen subjects (57%) displayed ≥ 1 cell/ μ L: 81 subjects (39%) had between 1-15 cells/ μ L, 32 (16%) between 16-79 cells/ μ L, and 5 (2%) >79 cells/ μ L. Thirteen patients with B-cell reconstitution at 12 months had early reconstitution with future infusions. Two of the subjects with CD19 values >15 cells/ μ L had clinical or magnetic resonance imaging relapse, compared to 8 subjects who did not have B-cell reconstitution. **Conclusions:** Although many patients displayed some B-cell repopulation prior to their next dose (113 subjects), CD19 counts of >79 cells/ μ L were uncommon (5 subjects). Subjects with early B-cell reconstitution at 1 infusion were likely to continue to show early repopulation at future infusions. Thus far, we have found no significant correlation between B-cell repopulation and either disease activity or adverse events.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Disease-modifying treatments in MS, Ocrelizumab

(DXT34) Revealing the Immune Cell Subtype Reconstitution Profile in Cladribine-Treated Patients at the 96-Week Timepoint (CLARITY) Using Deconvolution Algorithms

Irina Kalatskaya,¹ Gavin Giovannoni,^{2,3} Thomas P. Leist,⁴ Per Soelberg-Sorensen,⁵ Ursula Boschert,⁶ Julie DeMartino,¹ Alex Rolfe¹

¹EMD Serono, Inc, Billerica, MA; ²Bizard Institute of Cell and Molecular Science, London, United Kingdom; ³Queen Mary University of London, London, United Kingdom; ⁴Jefferson University Hospital, Philadelphia, PA; ⁵Department of Neurology, Danish MS Center, University of Copenhagen, Copenhagen, Denmark; ⁶Merck Serono S.A., Eysins, Switzerland

Background: Cladribine tablets (CTs) cumulative licensed dose of 3.5 mg/kg (CT3.5), administered as 2 short oral courses over 2 years, transiently reduces total lymphocyte counts, with median values returning to normal range within 11 months and median B cells by 6 months. Clinical efficacy of CTs is sustained beyond lymphocyte recovery. Flow cytometric observations suggest long-lasting reductions in memory B cells. **Objectives:** Characterize immune cell transcriptomic signatures in peripheral blood from patients with relapsing-remitting multiple sclerosis during immune repopulation at 96 weeks in the CLARITY study using advanced computational algorithms and correlate these signatures with corresponding flow cytometry data of main lymphocyte subtypes. **Methods:** Gene expression data (U133 Plus 2.0 array) in whole blood samples at 96 weeks were available from patients randomized to placebo (n = 57), CT3.5 (n = 62), or CT 5.25 mg/kg (CT5.25, n = 70). These were analyzed with the CIBERSORT deconvolution algorithm (to estimate absolute fractions of 22 immune cell subtypes) and the xCell signature-based method (cell type enrichment analysis for 43 immune cell subtypes).

Wilcoxon rank sum tests compared between treatment arms. Spearman rank correlation coefficient was used to measure the relationship between signatures and cell counts. *P* values < .05 were considered nominally significant. **Results:** At 96 weeks, the relative abundance of naive B cells in CT3.5- and CT5.25-treated patients was significantly higher vs placebo. Plasma cells and class-switched memory B cells were significantly reduced with CTs vs placebo. The M2 macrophage signature was significantly enhanced with CTs vs placebo. Cell abundance of both naive and memory CD4+ and CD8+ was significantly reduced with CTs vs placebo. Deconvolution signature scores were positively and significantly correlated with corresponding flow cytometry data (*r*: 0.68-0.72 CD19+ B cells, 0.71 CD4+ T cells, 0.67-0.69 CD8+ T cells). **Conclusions:** At 96 weeks following CT treatment in year 2, changes in leukocytes suggestive of a shift towards an anti-inflammatory phenotype were detected.

Supported by: None

Disclosure: *Irina Kalatskaya, Julie DeMarrino, Alex Rolfe:* EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). *Gavin Giovannoni:* AbbVie, Actelion, Almirall, Atara Bio, Bayer Schering Pharma, Five Prime, GlaxoSmithKline, GW Pharma, Merck KGaA, Pfizer Inc, Protein Discovery Laboratories, Sanofi Genzyme, Teva Pharmaceuticals Industries Ltd, UCB, Vertex Pharmaceuticals (consulting fee); *Biogen, Ironwood, Merck and Co, Novartis* (consulting fee, unrelated research support). *Thomas P. Leist:* Biogen, EMD Serono, Genentech/Roche, Janssen, Novartis, Teva (consulting fee). *Per Soelberg-Sorensen:* Biogen (advisory board, research support); *Biogen, Sanofi-Aventis* (speakers' bureau); *Genzyme* (research support, speakers' bureau); *GlaxoSmithKline* (advisory board, steering committees); *MedDay Pharmaceuticals* (advisory board); *Merck KGaA, Novartis, Teva* (advisory board, research support, speakers' bureau, steering committees); *Roche* (research support). *Ursula Boschert:* Merck Serono SA (a business of Merck KGaA, Darmstadt, Germany) (salary).

Keywords: Disease-modifying treatments in MS, Immunology and MS

(DXT35) Real-World Experience with Ocrelizumab: A Safety Analysis

Jamie Bolling,¹ Ryan McNiff,² Aaron Carlson,¹ Carlos Vervloet Sollero,¹ Tirisham V. Gyang¹

¹Neurology, University of Florida, Gainesville, FL; ²UF Health, University of Florida, Gainesville, FL

Background: Ocrelizumab is a humanized monoclonal antibody that selectively targets B-lymphocytes, resulting in their depletion. The US Food and Drug Administration approved its use in 2017 for relapsing-remitting and primary-progressive multiple sclerosis (MS). Pooled safety analysis from phase 3 clinical trials revealed an increase in incidence of infection, infusion reaction (IR), and malignancy in ocrelizumab patient groups.

Objectives: We present a real-world safety analysis of ocrelizumab in clinical practice. **Methods:** The University of Florida MS Clinic identified subjects as those treated with ocrelizumab prescribed in clinic through electronic medical records. The study collected longitudinal safety laboratories including complete blood count, lymphocyte subset counts, and immunoglobulin levels. The study also captured clinical data including disease course, prior disease-modifying therapies, IR, and occurrence of major clinical events. Analysis of the data assessed trends in laboratories and occurrence of adverse events (AEs). **Results:** Data from 39 of a potential 200 subjects suggests that white blood cell, neutrophil, lymphocyte, and T-cell counts continuously produce the most abnormal results after initiation of ocrelizumab. Nine subjects had urinary tract infections, and 2 subjects had respiratory tract infections. A case of sepsis and appendicitis resulted in 1 hospitalization. There were 12 IRs reported, and 1 subject discontinued treatment due to bronchospasms. One subject reported a diagnosis of intraductal carcinoma. **Conclusions:** The data reveals that infections and IRs are common among patients treated with ocrelizumab, while malignancies occur but are rare. Of infections, urinary tract infections pose the largest concern, although respiratory infections and secondary infections also occurred. Although IRs were common, they tended to be acute and easily resolved with the exception of 1 discontinuation. There is a need for more data to corroborate trends in laboratory values and potential correlation with AEs. Early findings suggest a significant trend in abnormal laboratory values, reported infections, and IRs.

Supported by: None

Disclosure: *Jamie Bolling, Ryan McNiff, Carlos Vervloet Sollero, Tirisham V. Gyang:* Nothing to disclose. *Aaron Carlson:* Novartis Pharmaceutical (contracted research); *Sanofi Genzyme* (consulting fee).

Keywords: Comprehensive care and MS, Disease-modifying treatments in MS, Treatment safety analysis in MS

(DXT36) Effect of Evobrutinib, a Bruton's Tyrosine Kinase Inhibitor, on Immune Cell and Immunoglobulin Levels over 48 Weeks in a Phase 2 Study in Relapsing Multiple Sclerosis

Xavier Montalban,^{1,2} Jamie Shaw,³ Sana Syed,³ Fernando Dangond,³ Emily C. Martin,³ Roland Grenningloh,³ Martin S. Weber,⁴ on behalf of the Evobrutinib Phase 2 Study Group

¹Vall d'Hebron University Hospital, Barcelona, Spain; ²St. Michael's Hospital, University of Toronto, Toronto, ON, Canada; ³EMD Serono Research & Development Institute, Inc (a business of Merck KGaA, Darmstadt, Germany), Billerica, MA; ⁴Institute of Neuropathology and Department of Neurology, University Medical Center, Göttingen, Germany

Background: Bruton's tyrosine kinase (BTK) plays an important role in proinflammatory pathways potentially involved in multiple sclerosis (MS). Consequently, BTK inhibition is being investigated as a potential therapeutic approach for MS. Evobrutinib, a highly selective BTK inhibitor (BTKi), has a dual mechanism of action, affecting both B cells and macrophages through inhibition of B-cell receptor, Fc receptor, and granulocyte-macrophage colony-stimulating factor receptor signaling, and has demonstrated clinical efficacy in MS in a phase 2 study (trial registration: NCT02975349; Montalban et al, ECTRIMS 2018 [P322]).

Objectives: To examine the effect of evobrutinib on immune cells and immunoglobulins (Igs) over 48 weeks. **Methods:** Patients aged 18-65 years with active relapsing-remitting MS or secondary progressive MS and superimposed relapses were randomized to receive either double-blind evobrutinib (25 mg once daily [qd], 75 mg qd, or 75 mg twice daily), placebo, or open-label dimethyl fumarate 240 mg (reference arm). After 24 weeks, placebo-treated patients were switched to evobrutinib 25 mg qd; other treatment arms continued under original allocation. Safety of evobrutinib, including assessment of B-cell count and Ig level, was a key secondary end point; investigations of the effects of evobrutinib on B-cell subsets, T-cell subsets, and natural killer cells in peripheral blood over 48 weeks were exploratory. **Results:** Of 267 patients randomized to treatment, 227 patients completed 48 weeks of treatment. No clinically relevant changes in the number of total B cells, or of memory B, mature naive B, total T, helper T, cytotoxic T, or natural killer cells, were observed in any evobrutinib treatment group over 48 weeks. No changes in IgG or IgG subtype levels were observed over 48 weeks in any treatment group. At week 48, there were slight increases from baseline in IgA and reductions in IgM for all evobrutinib groups, which were numerically greater than those with placebo at week 24. **Conclusions:** Patients with MS treated with the BTKi evobrutinib showed no evidence of B-cell depletion or change in mature vs naive B-cell subsets over 48 weeks. IgG levels remained stable and slight elevations in IgA levels were observed. These findings demonstrate that, in contrast to genetic deletion of BTK, continued pharmacologic BTK inhibition does not lead to B-cell depletion or significant reductions in circulating Igs.

Sponsored by: EMD Serono, Inc, Rockland, MA, USA, a business of Merck KGaA, Darmstadt, Germany

Disclosure: *Xavier Montalban:* Biogen, Merck Serono, Sanofi Genzyme, Novartis, Sanofi-Aventis, Teva Pharmaceuticals, Roche, Celgene, Actelion, National MS Society, MS International Federation, EXCEMED (speaking honoraria and travel expenses for participation in scientific meetings, steering committee member of clinical trials or participation in advisory boards of clinical trials). *Jamie Shaw, Sana Syed, Fernando Dangond, Emily C. Martin, Roland Grenningloh:* EMD Serono (salary). *Martin S. Weber:* Merck, Merck Serono, Novartis, Roche, Teva, Bayer, Genzyme (travel funding and/or speaker honoraria); *Deutsche Forschungsgemeinschaft* (DFG; WE 3547/5-1), *Novartis, Teva, Biogen, F. Hoffmann-La Roche, Merck, ProFutura Programm of the Universitätsmedizin Göttingen* (research support); *PLoS One* (editor).

Keywords: Clinical trials, Disease-modifying treatments in MS, Immunology and MS

(DXT37) Effect of Triflunomide on Brain Volume Loss in Patients with Relapsing Multiple Sclerosis of Differing Ages in TEMSO

Jiwon Oh,¹ Jens Wuerfel,² Bhupendra O. Khatri,³ Aaron E. Miller,⁴ Jihad Said Inshasi,⁵ Albert Saiz,⁶ Alex L. Lublin,⁷ Elizabeth M. Poole,⁷ Till Sprenger^{8,9}

¹University of Toronto, Toronto, ON, Canada; ²Medical Imaging Analysis Center (MIAC AG) and Department of Biomedical Engineering, University of Basel, Basel, Switzerland;

³Center for Neurological Disorders at Wheaton Franciscan Healthcare, Milwaukee, WI;

⁴Icahn School of Medicine at Mount Sinai, The Corinne Goldsmith Dickinson Center for Multiple Sclerosis, New York, NY; ⁵Dubai Medical College, Dubai, United Arab Emirates;

⁶Department of Neurology, Hospital Clinic and Institut d'Investigacions Biomèdiques August Pi i Sunyer, University of Barcelona, Barcelona, Spain; ⁷Sanofi, Cambridge, MA;

⁸DKD Helios Klinik Wiesbaden, Wiesbaden, Germany; ⁹University Hospital Basel, Basel, Switzerland

Background: Teriflunomide significantly reduced brain volume loss (BVL) vs placebo, assessed post hoc using structural image evaluation using normalization of atrophy (SIENA) in patients with relapsing forms of multiple sclerosis (RMS) enrolled in the phase 3 TEMSO study (trial registration: NCT00134563). It is not known whether the effect of teriflunomide on BVL differs by age. **Objectives:** To analyze the effect of teriflunomide treatment on BVL in patients of different age groups with RMS in TEMSO, with a focus on the >45-year age group. **Methods:** Patients were randomized 1:1:1 to receive placebo or teriflunomide 7 mg or 14 mg for ≤108 weeks. BVL was assessed as annualized percentage brain volume change (PBVC) from baseline using SIENA at years 1 and 2 in patients stratified by age: ≤25 years, >25 to 35 years, >35 to 45 years, and >45 years. Treatment group comparisons of median PBVC values were made via ranked analysis of covariance, adjusted for region, age, Expanded Disability Status Scale stratum, and baseline normalized brain volume. Data are presented at year 2 for all patients treated with teriflunomide 14 mg vs placebo, and for patients aged >45 years treated with teriflunomide 14 mg vs placebo. **Results:** The median annualized PBVC in all patients was 30.6% lower in the teriflunomide 14 mg group (n = 235) vs placebo (n = 234; P = .0001). In patients >45 years, the median annualized PBVC was 35.0% lower in the teriflunomide 14 mg group (n = 49) vs placebo (n = 48; P = .0098). **Conclusions:** Teriflunomide decelerated disease-related brain atrophy in patients with RMS compared with placebo, including in patients aged >45 years.

Supported by: Sanofi

Disclosure: *Jiwon Oh:* Biogen, Roche (consulting or speaking fees, research support); Celgene, EMD Serono, Genzyme, Novartis (consulting or speaking fees); Sanofi (research support). *Jens Wuerfel:* Actelion, Biogen, Novartis, Roche, Sanofi (scientific advisory boards); MIAc AG (Basel, Switzerland) (CEO). *Bhupendra O. Khatri:* Acorda, Celgene, Serono, Teva (consulting/honorarium [consulting work, speaker programs]); Alexion, Biogen, Genentech, Novartis, Sanofi (consulting/honorarium [consulting work, speaker programs], contracted research); Ra Pharmaceuticals (contracted research). *Aaron E. Miller:* Accordant Health Services, Adamas Pharmaceuticals, Celgene, EMD Serono, Genentech/Roche, Mallinckrodt Pharmaceuticals (Questcor), Mapi Pharma (consulting fee); Biogen, Novartis (consulting fee, contracted research); Genentech, Questcor, Roche, Sanofi (contracted research). *Jihad Said Inshasi:* Nothing to disclose. *Albert Saiz:* Bayer, Biogen, Merck-Serono, Novartis, Roche, Sanofi, Teva (remuneration for providing consulting services, lectures). *Alex L. Lublin, Elizabeth M. Poole:* Sanofi (salary). *Till Sprenger:* Actelion, Eli Lilly, Roche, Sanofi, Teva (current institution received payments for advisory board/consultation, speaking activities); EFIC-Grünenthal, Swiss MS Society, Swiss National Research Foundation (received grants); Novartis (current institution received payments for advisory board/consultation, speaking activities; received grants).

Keywords: Disease-modifying treatments in MS, Imaging and MS, Natural history of MS

(DXT38) Effects of Ozanimod on Information Processing Speed: Findings from the Phase 3 SUNBEAM and DAYBREAK Extension Trials

John DeLuca,¹ Jeffrey A. Cohen,² Bruce A.C. Cree,³ Hongjuan Liu,⁴ James K. Sheffield,⁴ Diego Silva,⁴ Giancarlo Comi,⁵ Ludwig Kappos⁶

¹Kessler Foundation, West Orange, New Jersey and Departments of Physical Medicine and Rehabilitation, and Neurology, Rutgers - New Jersey Medical School, Newark, NJ; ²Mellen Center, Cleveland Clinic, Cleveland, OH; ³Department of Neurology and Department of Ophthalmology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA; ⁴Bristol-Myers Squibb, Summit, NJ; ⁵Department of Neurology, San Raffaele Scientific Institute, Vita-Salute San Raffaele University, Milan, Italy; ⁶Neurologic Clinic and Policlinic, Departments of Medicine, Clinical Research, Biomedicine and Biomedical Engineering, University Hospital of Basel, Basel, Switzerland

Background: In the phase 3 SUNBEAM study, ozanimod HCl 1 mg improved information processing speed (IPS), measured with the Symbol Digit Modalities Test (SDMT), a component of a secondary end point, compared with interferon β-1a (IFN). **Objectives:** To evaluate long-term effects of ozanimod, a sphingosine 1-phosphate receptor modulator, on IPS in patients with relapsing multiple sclerosis (RMS). **Methods:** In the double-blind, double-dummy, SUNBEAM study (trial registration: NCT02294058), adults (18-55 years) with RMS were randomized to once-daily oral ozanimod HCl 1 or 0.5 mg, or weekly intramuscular IFN 30 µg. SUNBEAM continued until the last participant was treated for 12 months. Completers were eligible for an open-label extension study (DAYBREAK; NCT02576717) of ozanimod HCl 1 mg. Patients randomized to IFN in SUNBEAM transitioned to ozanimod HCl 1 mg in DAYBREAK 12-24 months after SUNBEAM baseline. This exploratory analysis reports the percentage of participants with clinically meaningful (≥2 point) improvement or worsening of SDMT scores at 12 and 24 months after

SUNBEAM baseline in those initially randomized to ozanimod HCl 1 mg or IFN. **Results:** In SUNBEAM, 447 participants were randomized to ozanimod HCl 1 mg and 448 to IFN [mean [SD] 13.5 [2.9] months of IFN exposure]; of these, 397 and 395, respectively, enrolled in DAYBREAK. Mean (SD) baseline SDMT scores were 47.7 (13.7) and 47.1 (13.5), respectively. At 12 months, 35.6% (152/427) of the ozanimod HCl 1 mg group and 27.9% (119/426) of the IFN group had SDMT improvement; 22.0% (94/427) and 28.2% (120/426), respectively, worsened. At month 24, 41.2% (113/274) of those who received continuous ozanimod HCl 1 mg and 34.5% (91/264) of those originally assigned to IFN in SUNBEAM had SDMT improvement; 21.9% (60/274) and 25.4% (67/264), respectively, worsened relative to SUNBEAM baseline. **Conclusions:** In this exploratory analysis, the percentage of participants with SDMT improvement increased over 24 months of continuous treatment with ozanimod HCl 1 mg. The percentage of participants with SDMT improvement was higher at month 24 than month 12 among those who transitioned from IFN to ozanimod HCl 1 mg during the latter 12 months as part of the DAYBREAK study.

Supported by: None

Disclosure: *John DeLuca:* Biogen, EMD Serono, Canadian MS Society, National MS Society, Consortium of Multiple Sclerosis Centers (CMSC) (grant funding); Celgene Corporation, Biogen, CMSC, Novartis, Sanofi Genzyme, Canadian MS Society, EXCEMED (consulting fee). *Jeffrey A. Cohen:* Convelo, Mylan, Population Council; Multiple Sclerosis Journal (editor, consulting fee). *Bruce A.C. Cree:* Akili, Alexion, Biogen, EMD Serono, Novartis, TG Therapeutics (consulting fee). *Hongjuan Liu, James K. Sheffield, Diego Silva:* Bristol-Myers Squibb (salary). *Giancarlo Comi:* Almirall, Biogen, Celgene Corporation, EXCEMED, Forward Pharma, Genzyme, Merck, Novartis, Roche, Sanofi, Teva (consulting fee). *Ludwig Kappos:* Bayer, Biogen, CSL Bering, Genzyme, Merck, Novartis, Sanofi, Teva (educational activities); Bayer, Biogen, European Union, Inmoswiss, Merck, Novartis, Roche, Swiss MS Society, Swiss National Research Foundation (license fees for Neurostatus products, grants); Bayer, Biogen, Merck, Novartis, Sanofi, Teva (speaking fees); Ludwig Kappos' institution (University Hospital Basel) has received the following, used exclusively for research support: steering committee, advisory board, and consultancy fees from Actelion, Addex, Bayer, Biogen, Biotica, Celgene Corporation; Ludwig Kappos' institution (University Hospital Basel) has received the following, which was used exclusively for research support: steering committee, advisory board, and consultancy fees from Genzyme, Eli Lilly, Merck, Mitsubishi, Novartis, Ono Pharma, Sanofi, Santhera, Siemens, Teva, UCB, Xenopart (consulting fee).

Keywords: Disease-modifying treatments in MS

(DXT40) Effect of the S1P1/5 Receptor Modulator Ozanimod on Cognitive Processing Speed in Subjects with Relapsing Multiple Sclerosis: Design of the ENLIGHTEN Study

Jon V. Riolo,¹ Ling Yang,¹ Thomas Rano,¹ Robert Zivadinov,² Lauren B. Krupp,³ Jonathan Calkwood,⁴ Olaf Stuve,⁵ Robert T. Naismith,⁶ Diego Silva,¹ John DeLuca⁷

¹Bristol-Myers Squibb, Summit, NJ; ²Buffalo Neuroimaging Analysis Center, Department of Neurology, Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, The State University of New York, Buffalo, NY; ³Stony Brook Technology Park, East Setauket, NY; ⁴Minneapolis Clinic of Neurology, Golden Valley, MN; ⁵UT Southwestern Medical Center, Dallas, TX; ⁶Washington University School of Medicine, St. Louis, MO; ⁷Kessler Foundation, West Orange, New Jersey and Departments of Physical Medicine and Rehabilitation, and Neurology, Rutgers - New Jersey Medical School, Newark, NJ

Background: In patients with multiple sclerosis (PwMS), slowed cognitive processing speed emerges as an early deficit. The Symbol Digit Modalities Test (SDMT) is a preferred measure of cognitive performance in PwMS. Ozanimod, an oral sphingosine 1-phosphate (S1P) receptor 1 and 5 modulator, was well tolerated and more effective than weekly intramuscular interferon β-1a (IFN) 30 µg on clinical and magnetic resonance imaging (MRI) end points in the phase 3 RADIANCE and SUNBEAM studies. The SUNBEAM study demonstrated a nominally significant (P < .05) improvement in SDMT with ozanimod HCl 1 or 0.5 mg/day over IFN 30 µg/week; however, the study was not designed to evaluate SDMT as a primary end point. **Objectives:** The primary objective of ENLIGHTEN (trial registration: NCT04140305) is to describe clinically meaningful changes in SDMT (≥4-point or 10% change from baseline) over 3 years in patients with early relapsing multiple sclerosis (RMS) treated with ozanimod HCl 1 mg/day. Secondary objectives are to describe changes from baseline in whole brain and substructure volume; MRI measures of disease activity; patient-reported outcomes (PROs) and quality of life (QOL); disability status based on Timed 25-Foot Walk, 9-Hole Peg Test, and Expanded Disability Status Scale (EDSS); and safety of ozanimod. The study also will explore the correlation between changes in cognitive processing speed and whole brain and substructure volume, and the cor-

relation between changes in cognitive processing speed and PROs and QOL. **Methods:** This ongoing multicenter, open-label study is recruiting 250 patients with RMS (aged 18-65 years) in the United States and Canada. Participants will receive ozanimod HCl 1 mg/day (equivalent to ozanimod 0.92 mg) for 3 years. Key inclusion criteria are diagnosis of multiple sclerosis per 2010 or 2017 McDonald criteria, ≤ 5 years since diagnosis, ≤ 1 approved RMS disease-modifying therapy, EDSS score ≤ 3.5 , and no relapse within 30 days of screening. Key exclusion criteria are prior therapy with ozanimod; clinically relevant cardiac, hepatic, neurologic, pulmonary, or other chronic conditions; >10 gadolinium-enhancing lesions on baseline brain MRI; or any condition or concomitant medication that might affect cognition or confound test performance. **Results:** The ENLIGHTEN study design will be presented. Cognitive performance, disease activity, disability status, and safety will be assessed over 3 years of ozanimod therapy. **Conclusions:** This study will determine if ozanimod has a clinically meaningful benefit on cognitive processing speed in patients with RMS.

Supported by: None

Disclosure: *Jon V. Riolo, Ling Yang, Thomas Rano, Diego Silva:* Bristol-Myers Squibb (salary). *Robert Zivadinov:* Celgene Corporation, EMD Serono, Sanofi Genzyme, Novartis (consulting fee); Celgene Corporation, Sanofi Genzyme, Mapi Pharma, Novartis, Protendis (contracted research). *Lauren B. Krupp:* Biogen, Novartis (consulting fee); Biogen, Lourie Foundation, National Institutes of Health, National Multiple Sclerosis Society, Novartis, US Department of Defense (grant support); Genentech (data safety monitoring board); pharmaceutical and biotechnology companies for use of Fatigue Severity Scale (royalty). *Jonathan Calkwood:* Celgene Corporation (research support, compensation for consulting, advisory boards). *Olaf Stuwe:* Celgene Corporation, EMD Serono, Genentech, Genzyme, TG Therapeutics (consulting fee); Genentech-Roche, Pfizer, TG Therapeutics (data monitoring committees without monetary compensation); Sanofi Genzyme (grant support); Therapeutic Advances in Neurological Disorders (editorial board). *Robert T. Naismith:* Alexion, Alkermes, Bayer AG, Biogen, Celgene Corporation, EMD Serono, Genentech, Genzyme, Novartis, Viela Bio (consulting fee). *John DeLuca:* Alexion, EMD Serono, Canadian MS Society, National MS Society, Consortium of Multiple Sclerosis Centers (CMSC) (grant funding); Celgene Corporation, Biogen, CMSC, Novartis, Sanofi Genzyme, Canadian MS Society, EXCEMED (consulting fee).

Keywords: Cognition and MS, Disease-modifying treatments in MS

(DXT41) Eculizumab Benefits a Broad Range of Patients with Aquaporin-4 Antibody-Positive Neuromyelitis Optica Spectrum Disorder: The Phase 3 PREVENT Study

Kazuo Fujihara,^{1,2,3} Achim Berthele,⁴ Ho Jin Kim,⁵ Michael Levy,^{6,7} Ichiro Nakashima,^{2,8} Celia Oreja-Guevara,⁹ Jacqueline Palace,¹⁰ Sean J. Pittock,¹¹ Murat Terzi,¹² Natalia Totolyan,¹³ Shanthi Viswanathan,¹⁴ Kai-Chen Wang,^{15,16} Amy Pace,¹⁷ Marcus Yountz,¹⁷ Larisa Miller,¹⁷ Imran Tanvir,¹⁷ Roisin Armstrong,¹⁷ Dean Wingerchuk¹⁸

¹Southern TOHOKU Research Institute for Neuroscience (STRINS), Koriyama, Japan; ²Tohoku University, Sendai, Japan; ³Fukushima Medical University, Fukushima City, Japan; ⁴Technical University of Munich, Munich, Germany; ⁵National Cancer Center, Goyang, Korea, Republic of (South); ⁶Massachusetts General Hospital and Harvard Medical School, Boston, MA; ⁷Johns Hopkins University, Baltimore, MD; ⁸Tohoku Medical and Pharmaceutical University, Sendai, Japan; ⁹Hospital Universitario Clinico San Carlos, Madrid, Spain; ¹⁰John Radcliffe Hospital, Oxford, United Kingdom; ¹¹Mayo Clinic, Rochester, MN; ¹²Ondokuz Mayıs University, Samsun, Turkey; ¹³First Pavlov State Medical University of St Petersburg, St Petersburg, Russian Federation; ¹⁴Kuala Lumpur Hospital, Kuala Lumpur, Malaysia; ¹⁵National Yang Ming University, Taipei, Taiwan; ¹⁶Cheng-Hsin General Hospital, Taipei, Taiwan; ¹⁷Alexion Pharmaceuticals, Boston, MA; ¹⁸Mayo Clinic, Scottsdale, AZ

Background: Antibodies to the aquaporin-4 (AQP4) water channel in neuromyelitis optica spectrum disorder (NMOSD) are reported to trigger the complement cascade, which is implicated in neuronal injury. The terminal complement inhibitor eculizumab is the first treatment approved for use in patients with AQP4-immunoglobulin G-positive NMOSD, based on data from the PREVENT study. **Objectives:** To determine whether eculizumab's beneficial effect in reducing relapse risk in patients with NMOSD is associated with time since diagnosis, relapse history, disability burden, or prior immunosuppressant therapy (IST) use, based on data from the phase 3 PREVENT study (trial registration: NCT01892345). **Methods:** In PREVENT, patients received eculizumab (maintenance dose, 1200 mg/2 weeks; n = 96) or placebo (n = 47), with stable-dose concomitant IST (except rituximab and mitoxantrone) permitted. PREVENT was not powered for subgroup analyses; a post hoc descriptive analysis was performed on subgroups defined by time since diagnosis, total number of historical relapses, baseline Expanded Disability Status Scale (EDSS) score, and prior IST use. **Results:** The proportions of patients having an adjudicated relapse were lower with eculizumab than with placebo in all subgroups. Proportions for those receiving eculizumab and

placebo, respectively, were: 2/31 (6.5%) vs 6/12 (50.0%) for <1 year since diagnosis and 1/65 (1.5%) vs 14/35 (40.0%) for ≥ 1 year since diagnosis; 1/39 (2.6%) vs 10/24 (41.7%) for 2-4 historical relapses and 2/57 (3.5%) vs 10/23 (43.5%) for ≥ 5 historical relapses; 0/14 vs 3/6 (50.0%) for baseline EDSS score ≤ 2.0 and 3/82 (3.7%) vs 17/41 (41.5%) for baseline EDSS score ≥ 2.5 to ≤ 7.0 ; 0/15 vs 2/5 (40.0%) for no prior IST use (except corticosteroids alone) and 3/81 (3.7%) vs 18/42 (42.9%) for prior IST use. Relapse risk reductions were consistent and statistically significant in all subgroups. **Conclusions:** The data from this post hoc subgroup analysis suggest that eculizumab reduced relapse risk compared with placebo in patients with AQP4-immunoglobulin G-positive NMOSD, regardless of time since NMOSD diagnosis, relapse history, disability burden, or prior IST use.

Supported by: None

Disclosure: *Kazuo Fujihara:* Alexion Pharmaceuticals, Biogen, Chugai, Mitsubishi-Tanabi, Novartis (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent); Asahi Medical, Bayer, Eisai, Roche, Teijin (fees for non-CME/CE services received directly from commercial interest or its agent). *Achim Berthele:* Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent, speakers' bureau). *Ho Jin Kim:* Alexion Pharmaceuticals, Celltrion, Eisai, HanAll BioPharma, Merck Serono, Novartis, Sanofi Genzyme, Teva-Handok, Viela Bio (consulting fee); Journal of Clinical Neurology (associate editor); MedImmune/Viela Bio (steering committee); Multiple Sclerosis Journal (co-editor). *Michael Levy:* Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent); Genentech, Quest Diagnostics, Viela Bio (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent). *Ichiro Nakashima:* Alexion (consulting fee). *Celia Oreja-Guevara:* Alexion Pharmaceuticals, Merck, Novartis, Roche, Sanofi (consulting fee, contracted research). *Jacqueline Palace:* Biogen, Chugai (contracted research); LEK, Viela Bio, Guidepoint (consulting fee); Merck Serono (meeting/lecture/workshop participation); Novartis, Roche, Argenx (speakers' bureau); UCB, Viela Bio, Roche (conference/lecture participation). *Sean J. Pittock:* Alexion Pharmaceuticals (all compensation paid directly to Mayo Clinic, consulting fee, contracted research); Astellas (all compensation paid directly to Mayo Clinic, consulting fee); Autoimmune Encephalitis Alliance (all compensation paid directly to Mayo Clinic, contracted research); Grifols (all compensation paid directly to Mayo Clinic, contracted research); MedImmune (all compensation paid directly to Mayo Clinic, consulting fee, contracted research); UCB (consulting fee, personal compensation for attending UCB advisory board meeting in Stockholm, Sweden, on Sept 10, 2019). *Murat Terzi:* Shanthi Viswanathan, Kai-Chen Wang: Nothing to disclose. *Natalia Totolyan:* Alexion, Janssen, Novartis, Roche, Sanofi, Receptos Inc, Biocad (Russia) (contracted research); Merck (consulting fee); Roche, Sanofi (fees for lectures). *Amy Pace, Marcus Yountz, Roisin Armstrong:* Alexion Pharmaceuticals (ownership interest, salary). *Larisa Miller:* Alexion Pharmaceuticals (RSU, salary). *Imran Tanvir:* Alexion Pharmaceuticals (salary). *Dean Wingerchuk:* MedImmune, Novartis, Biogen, Celgene, Genentech, TG Therapeutics, Arcus Medica, Reistone (consulting fee); Terumo BCT, Alexion Pharmaceuticals (fees for non-CME/CE services received directly from commercial interest or its agent).

Keywords: Eculizumab in NMOSD

(DXT42) Rationale and Design of CLASSIC-MS Study Evaluating Long-Term Efficacy for Patients with Multiple Sclerosis Treated with Cladribine Tablets

Alexey Boyko,¹ Jorge Correale,² Gilles Edan,³ Mark S. Freedman,⁴ Gavin Giovannoni,⁵ Xavier Montalban,^{6,7} Kottil Rammohan,⁸ Thomas P. Leist,⁹ Dusan Stefoski,¹⁰ Bassem Yamout,¹¹ Belen Garcia-Alonso,¹² Aida Aydemir,¹³ Elisabetta Verdun di Cantogno,¹² CLASSIC-MS Study Group

¹Pirogov Russian National Research Medical University, Moscow, Russia; ²FLENI Institute, Buenos Aires, Argentina; ³Department of Neurology, University Hospital of Rennes, Rennes, TX; ⁴University of Ottawa and the Ottawa Hospital Research Institute, Ottawa, ON, Canada; ⁵Queen Mary University of London, London, United Kingdom; ⁶Centre d'Esclerosi Múltiple de Catalunya (Cemcat), Hospital Universitario Vall d'Hebron, Barcelona, Spain; ⁷St. Michael's Hospital, University of Toronto, Toronto, ON, Canada; ⁸Department of Neurology, Multiple Sclerosis Center, University of Miami, Miami, FL; ⁹Jefferson University Hospital, Philadelphia, PA; ¹⁰Rush Medical College, Chicago, IL; ¹¹Department of Neurology, American University of Beirut Medical Center, Beirut, Lebanon; ¹²Merck KGaA, Darmstadt, Germany; ¹³EMD Serono Research & Development Institute Inc, Billerica, MA

Background: Cladribine tablets 10 mg (CT; cumulative dose 3.5 mg/kg over 2 years) has demonstrated efficacy vs placebo over 2 years in CLARITY, CLARITY Extension, and ORACLE-MS, showing sustained efficacy without further active treatment in CLARITY Extension. **Objectives:** CLASSIC-MS will explore long-term efficacy and real-world treatment patterns in patients who participated in these trials. Long-term safety in this

population has been assessed in the PREMIERE registry. **Methods:** CLASSIC-MS is an exploratory phase 4 study of patients with MS, or those with a first clinical demyelinating event enrolled into the phase 3 trials and who received ≥ 1 course of CT or placebo (N = 1946). Following pre-baseline screening and assessment for eligibility, long-term retrospective data will be obtained from medical records at study visit 1; prospective data will be collected at study visits 1 and 2. Patients will be enrolled for 17 months from approximately Q3 2019 to Q4 2020. Last patient last visit is expected in Q1 2021. Primary objective: evaluation of long-term mobility after treatment with CT or placebo. Secondary objective: assess differences in clinical and magnetic resonance imaging characteristics in long-term responders vs nonresponders. Tertiary end points: real-world treatment patterns, durability of clinical outcomes, quality of life and cognitive outcomes, influence of high-disease activity on long-term response, and differences in genetics between long-term responders and those who are not. **Results:** In 2018, a second feasibility survey was sent to 225 centers; 110 centers provided positive responses and were included, representing 48% of sites originally enrolled in the phase 3 studies. In total 115 centers were not included (81 were not willing to participate; 13 dropped; 16 were nonresponders; 5 were rejected). **Conclusions:** CLASSIC-MS will provide valuable information on the long-term efficacy of CT in patients with MS.

Supported by: None

Disclosure: Alexey Boyko: Biogen, Schering, Merck, Teva, Novartis, Sanofi Genzyme, Actelion, Biocad, Genierum (consulting fee, participated in clinical trials supported by). Jorge Correale: Merck-Serono Argentina, Biogen LATAM, Merck-Serono LATAM, Genzyme (board member); Merck-Serono Argentina, Merck-Serono LATAM, Biogen Argentina, Genzyme Argentina, Teva Argentina (fees for non-CME/CE services received directly from commercial interest or its agent). Gilles Edan: Biogen, Genzyme, Merck Serono, Novartis, Roche, Teva Pharma (consulting fee). Mark S. Freedman: Actelion, Bayer HealthCare, Biogen, Chugai, EMD Canada, Genzyme, F. Hoffman-La Roche, Novartis, Sanofi, Teva (consulting fee). Gavin Giovannoni: AbbVie, Actelion, Atara Bio, Almirall, Bayer Schering Pharma, Biogen, Five Prime, GlaxoSmithKline, GW Pharma, Merck & Co, Merck KGaG, Pfizer Inc, Protein Discovery Laboratories, Teva Pharmaceutical Industries Ltd, Sanofi Genzyme, UCB (speakers' bureau); AbbVie, Actelion, Atara Bio, Almirall, Bayer Schering Pharma, Biogen, Five Prime, GlaxoSmithKline, GW Pharma, Merck & Co, Merck KGaG, Pfizer Inc, Protein Discovery Laboratories, Teva Pharmaceutical Industries Ltd, Sanofi Genzyme, UCB, (consulting fee); Biogen, Merck & Co, Novartis, Ironwood (research support); Vertex Pharmaceuticals, Ironwood, Novartis (consulting fee, speakers' bureau). Xavier Montalban: Biogen, Merck Serono, Genentech, Genzyme, Novartis, Sanofi-Aventis, Teva Pharmaceuticals, Roche, Celgene, Actelion, National MS Society, MS International Federation, EXCEMED (consulting fee, speakers' bureau). Kottil Rammohan: EMD Serono, Biogen, Sanofi-Aventis, Genzyme, Novartis, Teva Neurosciences, Acorda, Roche/Genentech (consulting fee, speakers' bureau). Thomas P. Leist: Acorda, Bayer, Biogen, Daiichi, EMD Serono, Novartis, Ono, Pfizer, Teva Neuroscience (consulting fee). Dusan Stefanski: Acorda, Biogen, Teva Neuroscience (consulting fee, speakers' bureau); Elan, EMD Serono (speakers' bureau); Merck Serono (consulting fee). Bassem Yamout: Bayer, Biogen, Genpharm, Genzyme, Merck-Serono, Novartis; and has received research grants from Bayer, Biogen, Merck-Serono, Novartis, Pfizer (consulting fee, speakers' bureau). Belen Garcia-Alonso, Elisabetta Verdun di Cantogno: Merck KGaA (salary). Aida Aydemir: EMD Serono (salary).

Keywords: Disease-modifying treatments in MS, Management of activities of daily living in MS, Real-world treatment patterns

(DXT43) Analyses of the Effect of Baseline Age on the Efficacy and Safety of Siponimod in Patients with Active Secondary Progressive Multiple Sclerosis from the EXPAND Study

Le H. Hua,¹ Amit Bar-Or,² Fred D. Lublin,³ Xiangyi Meng,⁴ Wendy Su,⁴ Bruce A.C. Cree,⁵ Robert J. Fox⁶

¹Lou Ruvo Center for Brain Health, Las Vegas, NV; ²Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ³Icahn School of Medicine at Mount Sinai, New York, NY; ⁴Novartis Pharmaceuticals Corporation, East Hanover, NJ; ⁵Department of Neurology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA; ⁶Cleveland Clinic, Cleveland, OH

Background: Siponimod (Mayzent) is a selective sphingosine 1-phosphate receptor (S1P₁ and S1P₂) modulator, approved in the United States for treatment of relapsing forms of multiple sclerosis (MS), including clinically isolated syndrome, relapsing-remitting MS, and active secondary progressive MS (SPMS). In the phase 3 EXPAND registration trial in SPMS, siponimod significantly reduced risk of 3-month (primary end point) and 6-month confirmed disability progression (CDP) by 21% and

26%, respectively. **Objectives:** Assess efficacy and safety of siponimod in patients with active SPMS in subgroups of patients aged <45 and ≥ 45 years (median value) at baseline. **Methods:** Post hoc analyses were performed in subgroups of patients with active SPMS, defined as a relapse in the 2 years before screening and/or ≥ 1 T1 gadolinium-enhancing lesion at baseline, randomized to siponimod 2 mg daily or placebo. Efficacy end points included: time to 3- and 6-month CDP (as per Expanded Disability Status Scale scores). Adverse events (AEs), serious AEs, and AEs leading to treatment discontinuation were also assessed. Analyses for hypothesis generation only; no adjustment for multiple comparisons. **Results:** There were 779 patients with active SPMS: 306 patients aged <45 years (siponimod, n = 213; placebo, n = 93) and 473 patients aged ≥ 45 years (siponimod, n = 303; placebo, n = 170). In those <45 years, siponimod reduced risk of 3-month CDP by 31.9% compared with placebo (siponimod, n = 57 [26.8%]; placebo, n = 35 [37.6%]; hazard ratio [HR] [95% CI]: 0.68 [0.45, 1.04]; P = .0734), and reduced 6-month CDP risk by 39.5% (siponimod, n = 44 [20.7%]; placebo, n = 30 [32.3%]; HR [95% CI]: 0.61 [0.38, 0.96]; P = .0339). In the subgroup of patients ≥ 45 years, siponimod reduced the risk of 3-month and 6-month CDP by 31.5% and 33.1%, respectively, vs placebo [3 month: siponimod, n = 72 [23.8%]; placebo, n = 56 [32.9%]; HR [95% CI]: 0.69 [0.48, 0.97]; P = .0340; 6-month: siponimod, n = 55 [18.2%]; placebo, n = 44 [25.9%]; HR [95% CI]: 0.67 [0.45, 1.0]; P = .0471). Siponimod was generally well tolerated in both subgroups. Rates of any AE were similar for siponimod and placebo in patients <45 years (82.6% vs 82.8%), and slightly higher for siponimod in those ≥ 45 years (89.8% vs 75.9%). Rates of AEs and AEs leading to discontinuation were similar between groups. **Conclusions:** In EXPAND, siponimod provided similar clinical effects in reducing CDP risk in patients aged <45 years and ≥ 45 years with active SPMS.

Supported by: None

Disclosure: Le H. Hua: Biogen, Celgene, EMD Serono, Genentech, Genzyme, Novartis (consulting fee). Amit Bar-Or: Atara Biotherapeutics, Bayer, Bayhill Therapeutics, Berlex, Biogen, BioMS, Celgene/Receptos, Diogenix, Eli Lilly, F. Hoffman-La Roche, Genentech, GlaxoSmithKline, Guthy-Jackson/GGF, Immuno, Janssen/Actelion, Mapi, MedImmune, Merck/EMD Serono, Novartis, Ono Pharmacia, Roche/Genentech, Sanofi Genzyme, Teva Neuroscience, Wyeth (consulting fee). Fred D. Lublin: AbbVie, Acorda Therapeutics, Apitope, Atara Biotherapeutics, Bayer HealthCare Pharmaceuticals, Biogen, Brainstorm Cell Therapeutics, EMD Serono, Foward Pharma, Innate Immunotherapeutics, Mapi Pharma, MedDay Pharma, MedImmune, Novartis, Orion Biotechnology, Polpharma, Receptos/Celgene, Regeneron, Roche Genentech, Sanofi Genzyme, Teva Neuroscience, TG Therapeutics (consulting fee); Actelion (consulting fee, research support); Multiple Sclerosis and Related Disorders (journal) (editor); National MS Society, Novartis Pharmaceuticals, Sanofi, Teva Neurosciences, Transparency Life Sciences (research support). Xiangyi Meng, Wendy Su: Novartis Pharmaceuticals Corporation (salary). Bruce A.C. Cree: Akili, Alexion, Atara, Biogen, EMD Serono, Novartis, TG Therapeutics (consulting fee). Robert J. Fox: Actelion, EMD Serono, Genentech, Teva (consulting fee); Biogen (clinical trial contracts, consulting fee); Novartis (clinical trial contracts, consulting fee, contracted research).

Keywords: Disease-modifying treatments in MS

(DXT44) Real-World Patterns of Disease Progression in Patients with Multiple Sclerosis Who Are Adherent Versus Nonadherent to Disease-Modifying Treatments over 6 Years

Lilyana Amezcua,¹ Mitz J. Williams,² Jia Zhou,³ Terrie Livingston³

¹University of Southern California, Los Angeles, CA; ²Joi Life Wellness Group, Newnan, GA; ³EMD Serono Inc, Rockland, MA

Background: Multiple sclerosis (MS) is a chronic disease that requires long-term treatment for most patients. Disease-modifying treatment (DMT) adherence is often an issue for patients with MS, and evidence suggests that nonadherence can affect outcomes. Data are needed to understand the impact of long-term DMT use on clinical outcomes. **Objectives:** To assess the impact of long-term DMT adherence on MS disease progression in the real world. **Methods:** A retrospective cohort analysis of MarketScan Commercial enrollees from 2011-2017 was performed. MS was defined as ≥ 3 ICD-9/10 [340/G35] diagnosis claims or ≥ 1 diagnosis and ≥ 1 DMT claim (age 18 and 65 years at index) with index date being the first diagnosis or DMT claim in 2012. Continuous enrollment started from 1 year preindex, with a follow-up of ≥ 3 years of continuous enrollment and up to 6 years. Adherent-users as medication possession ratio (MPR) ≥ 0.8 in follow-up, and non-adherent users as $0 < \text{MPR} < 0.8$. Propensity score greedy matching was used to balance population characteristics (age, gender, geography, comorbidities, relapses) 1 year preindex. We compared the average number of relapses, defined as a

hospitalization with a primary diagnosis of 340/G35 or an outpatient visit with diagnosis of 340/G35 plus a pharmacy or medical claim for a qualifying corticosteroid within 7 days, between 2 cohorts using Poisson regression model. We also compared the time to first relapse, time to cane/walker use, and time to wheelchair use between 2 cohorts using Cox-proportional hazard model. **Results:** 15,617 patients with MS were identified (42% adherent, 43% nonadherent, 15% non-DMT treated). Of them, 6121 propensity score matched pairs were analyzed. Standardized differences of all baseline characteristics between 2 comparison groups were <0.1. Adherent users had significantly lower average number of relapses (0.153) than nonadherent (0.201) users (annualized relapse rate ratio: 0.76, 95% CI: 0.74-0.79, $P < .001$). Adherent users had significantly longer time to first relapse [hazard ratio [HR] = 0.82, 95% CI: 0.77-0.87, $P < .001$], cane/walker use (HR = 0.81, 95% CI: 0.71-0.93, $P = .003$), and wheelchair use (HR = 0.60, 95% CI: 0.51-0.70, $P < .001$). **Conclusions:** This study highlights the importance of DMT adherence in slowing disease progression. Indicators of MS-related disability were found to be related to adherence, suggesting a lower rate of disability progression over time. Further research is needed to better understand barriers of adherence with DMTs.

Supported by: None

Disclosure: *Lilyana Amezcua:* Biogen (consulting fee, research support); California Community Foundation, MedDay, NIH National Institute of Neurological Disorders and Stroke, National MS Society (research support); EMD Serono, Novartis (consulting fee). *Mitzi J. Williams:* EMD Serono, Inc (employer received funding for study); Joi Life Wellness Group (salary). *Jia Zhou, Terrie Livingston:* EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary).

Keywords: Disease-modifying treatments in MS

(DXT45) Pharmacist-Based Intervention for Improving Baseline Laboratory Monitoring for Patients on Multiple Sclerosis Disease-Modifying Therapies

Lisa Aquillano,¹ Kristina Murphy,² Neil Lava,¹ Diana Vargas¹

¹Neurology and ²Pharmacy, The Emory Clinic, Atlanta, GA

Background: Disease-modifying therapies (DMTs) require baseline laboratory tests (labs) prior to initiation. Under our institution's pharmacotherapy protocol, clinical pharmacy specialists may order labs based on their assessment of the patient's monitoring needs. Currently, there is no standardized clinical practice guideline for ordering appropriate labs prior to initiating medications prescribed for patients with multiple sclerosis (MS). Frequently, baseline laboratory values are not available in the chart, making it challenging to determine treatment and evaluate safety. A standardized process, including the utilization of the pharmacotherapy protocol, would optimize safety and guide clinical decisions regarding the management of patients prescribed DMTs for MS. **Objectives:** The percentage of appropriate labs drawn for the specified DMTs following the implementation of a standardized operating protocol in the MS outpatient neurology clinic. **Methods:** A clinical practice guideline was implemented on October 1, 2019, outlining the baseline labs required within 6 months prior to initiation of specific DMTs. The following DMTs were included in the guideline: interferon beta-1a, peginterferon beta-1a, dimethyl fumarate, and teriflunomide. All patients who had a medication prescribed for the listed DMT from October 1, 2018, to March 31, 2019, were included in the baseline analysis. The clinical pharmacy specialist used the pharmacotherapy protocol to order appropriate labs when a therapy was initiated. A post analysis was performed to compare rates of appropriate labs drawn in all patients who had a prescription for the DMTs listed above, from October 1, 2019, to March 31, 2020. The patients included were those starting DMT for the first time, changing to a new DMT, or restarting a DMT after being off therapy for >6 months. **Results:** (Preliminary) Of the 43 patients prescribed a DMT during the baseline analysis, 13 (30.2%) patients had all appropriate labs drawn within 6 months. Of the remaining 30 (69.8%) patients who did not have all appropriate labs drawn, 18 (60%) patients had no labs documented in the chart within the past 6 months. The most common labs not drawn include differential (83.3%), LFTs (63.3%), and CBC (56.6%). Since the implementation of the clinical practice guideline, 75% of patients have had all appropriate labs drawn within 6 months of therapy initiation. However, baseline labs have been collected on 100% of patients, with only the differential not drawn. **Conclusions:** Integration of a clinical pharmacy specialist in the management of patient safety is crucial to delivering exceptional care. Implementation of a standardized operating procedure and incorporation of pharmacists to assist in the monitoring of patients, contributed to an increase in appropriate labs drawn prior to initiation of DMT. Additionally, the utilization of pharmacists practicing in multidisciplinary settings to further elevate the level of care provided to patients could be extrapolated to other clinic areas and disease states.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Disease-modifying treatments in MS, Pharmacist

(DXT46) Cognitive Performance and Disability Across Age Groups in Teriflunomide-Treated Patients in the Teri-PRO Study

Lori H. Travis,¹ Dennis Chinchilla,² Bhupendra O. Khatri,³ José Meca-Lallana,⁴ Min Su Park,⁵ Patrick Vermersch,⁶ Alex Lublin,⁷ Elizabeth M. Poole,⁷ Patricia K. Coyle⁸

¹Phoenix Neurological Associates, Phoenix, AZ; ²Department of Neurology, CIMA Hospital, San José, Costa Rica; ³Center for Neurological Disorders at Wheaton Franciscan Healthcare, Milwaukee, WI; ⁴Hospital Virgen de la Arrixaca, Universidad Católica San Antonio de Murcia, Murcia, Spain; ⁵Department of Neurology, Yeungnam University Hospital, Daegu, Korea, Republic of (South); ⁶University of Lille, INSERM U995, CHU Lille, Lille, France; ⁷Sanofi, Cambridge, MA; ⁸Stony Brook University, Stony Brook, NY

Background: The Teri-PRO study (trial registration: NCT01895335) evaluated patient-reported outcomes, including cognition, in teriflunomide-treated patients with relapsing forms of multiple sclerosis (MS) in a real-world setting. It is unknown whether treatment effects on cognitive performance are influenced by patient age and/or physical disability. **Objectives:** To evaluate the effects of teriflunomide treatment on cognition across age groups, in the context of disability. **Methods:** Teri-PRO was a phase 4, multicenter, prospective, single-arm, open-label, real-world study assessing treatment satisfaction of teriflunomide 7 mg or 14 mg over 48 weeks using patient-reported outcomes. Cognitive performance, measured using the Symbol Digit Modalities Test (SDMT), and disability, assessed with the Expanded Disability Status Scale (EDSS), were secondary end points. The SDMT was scored as the proportion of correct responses (scale 0-1). Patient improvement was computed as the change in score from baseline to 48 weeks; Spearman correlation assessed the relationship between SDMT and EDSS outcomes. **Results:** SDMT and EDSS data were available in 839 patients (≤25 years: n = 21; >25 to 35 years: n = 105; >35 to 45 years: n = 243; >45 to 55 years: n = 279; and >55 years: n = 191). Baseline mean SDMT scores across all age groups were similar (≤25 years: 0.99; >25 to 35 years: 0.98; >35 to 45 years: 0.98; >45 to 55 years: 0.97; and >55 years: 0.97), and remained stable through week 48 (least squares mean change ranged from -0.01 to +0.01 across groups). At baseline, mean EDSS scores were higher with advancing age, from 1.52 in patients ≤25 years to 4.12 in patients >55 years. Least squares mean EDSS changes from baseline to week 48 were not significant, except in patients >45 to 55 years (+0.13, $P = .0079$). Cognition and disability were not correlated in any age group (Spearman correlations, ≤25 years: -0.28, $P = .2$; >25 to 35 years: 0.02, $P = .9$; >35 to 45 years: -0.09, $P = 0.2$; >45 to 55 years: -0.05, $P = .4$; >55 years: 0.04, $P = .6$) or in the overall population (-0.034, $P = .3$). **Conclusions:** Across all age groups, patients in Teri-PRO had similarly high cognitive function at baseline, and cognitive performance after 48 weeks of teriflunomide treatment was stable. SDMT and EDSS were not significantly correlated.

Supported by: Sanofi

Disclosure: *Lori H. Travis:* Acorda, Biogen, Mallinckrodt, Novartis, Pfizer (consulting fee); Biogen (grant/research support); EMD Serono, Genzyme (consulting fee, grant/research support). *Dennis Chinchilla:* Asofarma, Genzyme, Novartis, Stendhal (educational support); Bayer (educational support, speaker honoraria); Glaxo (speaker honoraria); Merck Serono, Roche (compensation on scientific advisory board, educational support). *Bhupendra O. Khatri:* Acorda, Celgene, Serono, Teva (consulting/honorarium [consulting work, speaker programs]); Alexion, Biogen, Genentech, Novartis, Sanofi (consulting/honorarium [consulting work, speaker programs], contracted research); Ra Pharmaceuticals (contracted research). *José Meca-Lallana:* Almirall, Biogen, Celgene, Genzyme, Merck, Novartis, Roche, Teva (honoraria for consulting or presentations). *Min Su Park:* Bayer, Genzyme, Merck Serono, Novartis (consulting fees, honoraria, or scientific committee support). *Patrick Vermersch:* Almirall, Biogen, Celgene, Merck, Novartis, Roche, Sanofi, Servier, Teva (consulting and/or speaking fees and research support). *Alex Lublin, Elizabeth M. Poole:* Sanofi (salary). *Patricia K. Coyle:* Accordant, Alexion, Bayer, Biogen, Celgene, Sanofi Genzyme, GlaxoSmithKline, Mylan, Serono, TG Therapeutics (consulting fees and/or speaking fees); Actelion, Alkermes, Corona LLD, MedDay, National Institute of Neurological Disorders and Stroke, PCORI (research support); Genentech/Roche, Novartis (consulting fees and/or speaking fees, research support).

Keywords: Aging and MS, Cognition and MS, Disease-modifying treatments in MS

(DXT48) Efficacy of Subcutaneous Interferon Beta-1a in Patients with a First Clinical Demyelinating Event: REFLEX Study – Outcomes in Patients Stratified by 2017 McDonald Criteria

Mark S. Freedman,¹ Ludwig Kappos,² Giancarlo Comi,³ Nicola De Stefano,⁴ Sanjeev Roy,⁵ Delphine Issard⁶

¹University of Ottawa and the Ottawa Hospital Research Institute, Ottawa, ON, Canada; ²Neurologic Clinic and Policlinic, Departments of Medicine, Clinical Research, Biomedicine and Biomedical Engineering, University Hospital of Basel, Basel, Switzerland; ³Department of Neurology and Institute of Experimental Neurology, University Vita-Salute San Raffaele, Ospedale San Raffaele, Milan, Italy; ⁴Department of Neurological and Behavioural Sciences, University of Siena, Siena, Italy; ⁵Merck KGaA, Aubonne, Switzerland; ⁶Cytel Inc, Geneva, Switzerland

Background: The REFLEX (REbif FLEXible dosing in early multiple sclerosis [MS]) trial demonstrated that subcutaneous interferon beta-1a (sc IFN β -1a) reduced conversion to MS (McDonald 2005 criteria) and to clinically definite MS (CDMS) vs placebo in patients with a first clinical event suggestive of MS. A retrospective analysis of the study showed that the overall results were unchanged by the application of the McDonald 2010 MS criteria. The revised 2017 McDonald MS criteria include the presence of cerebrospinal fluid-specific oligoclonal bands, symptomatic lesions, and cortical lesions to aid MS diagnosis. **Objectives:** Assess the effects of sc IFN β -1a on time to McDonald 2005 criteria MS (time to next relapse, Expanded Disability Status Scale [EDSS] progression, or MS-related magnetic resonance imaging lesion or lesions) and CDMS (time to relapse or EDSS progression), and annualized relapse rate (ARR) during REFLEX, stratified by retrospective diagnosis at baseline in patients that either meet or do not meet the updated McDonald 2017 MS criteria. **Methods:** During REFLEX, patients were randomized to either sc IFN β -1a three times weekly (tiw) or once weekly (qw), or placebo, for 2 years. This retrospective analysis stratified patients randomized to the intent-to-treat population in REFLEX into McDonald 2017–positive (defined as those who retrospectively met the 2010 McDonald MS criteria at baseline or those with positive oligoclonal bands) and McDonald 2017–negative subgroups. Kaplan-Meier curves were used to estimate time to McDonald 2005 MS and time to CDMS by treatment group and for each McDonald 2017 subgroup. **Results:** As the detection of oligoclonal bands was optional during REFLEX, only a small number of patients were added from the McDonald 2010 analysis. A total of 235/517 patients were classed as McDonald 2017–positive at baseline (40 of whom were McDonald 2010–negative but had positive oligoclonal bands). In the McDonald 2017–positive subgroup, treatment with sc IFN β -1a tiw or qw significantly delayed time to McDonald 2005 MS (tiw vs placebo hazard ratio [HR] = 0.47, $P < .001$; qw vs placebo HR = 0.58, $P = .002$) and CDMS (tiw vs placebo HR = 0.46, $P = .010$; qw vs placebo HR = 0.42, $P = .003$) vs placebo. Treatment with sc IFN β -1a qw or tiw significantly reduced mean ARR vs placebo in McDonald 2017–positive patients (reductions of 69.1% and 59.3%, respectively; $P < .001$). **Conclusions:** The treatment effects of sc IFN β -1a observed in McDonald 2010 patients on time to McDonald 2005 MS and CDMS were maintained in the McDonald 2017–positive subgroup, although there were only a small number of additional patients when the 2017 criteria were applied.

Supported by: None

Disclosure: *Mark S. Freedman:* Actelion, Bayer HealthCare, Biogen, Chugai, EMD Canada, Genzyme, F. Hoffmann-La Roche, Novartis, Sanofi, Teva (consulting fee); EMD Serono, Genzyme (speakers' bureau). *Ludwig Kappos:* Institutional funding: Actelion, Adxex, Bayer HealthCare, Biogen, Biotica, Genzyme, Eli Lilly, Merck KGaA, Mitsubishi, Novartis, Ono Pharma, Pfizer, Receptos, Sanofi, Santhera, Siemens, Teva, UCB, Xenoport (consulting fee); Institutional funding: Bayer, Biogen, CSL Bering, Genzyme, Merck, Novartis, Sanofi, Teva (educational activities); Institutional funding: Bayer, Biogen, European Union, Imnouis, Merck, Novartis, Roche, Swiss MS Society, Swiss National Research Foundation (contracted research); Institutional funding: Bayer, Biogen, Merck, Novartis, Sanofi, Teva (speakers' bureau); Institutional funding: Neurostatus products (receipt of intellectual property rights/patent holder). *Giancarlo Comi:* MedDay, EXCEMED, Novartis, Teva Pharmaceutical Industries Ltd, Teva Italia Srl, Sanofi Genzyme, Genzyme Corporation, Genzyme Europe, Merck KGaA, Merck Serono S.p.A., Celgene Group, Biogen, Biogen Italia Srl, F. Hoffmann-La Roche, Roche SpA, Almirall SpA, Forward Pharma (consulting fee, speakers' bureau). *Nicola De Stefano:* Biogen, Teva, Novartis, Sanofi Genzyme, Roche, Merck Serono (speakers' bureau); FISM, Novartis (contracted research); Schering, Biogen, Teva, Novartis, Sanofi Genzyme, Roche, Merck Serono (consulting fee); Teva, Novartis, Sanofi Genzyme, Roche, Merck Serono (travel). *Sanjeev Roy:* Merck, Aubonne, Switzerland (salary). *Delphine Issard:* Cytel Inc (salary); Merck KGaA (consulting fee).

Keywords: Diagnosis, Biomarkers, Disease-modifying treatments in MS

(DXT49) Post Hoc Analysis of Efficacy of Cladribine Tablets in Patients with Relapsing-Remitting Multiple Sclerosis Aged Over and Under 30 Years in the CLARITY Study

Mark S. Freedman,¹ Gabriel Pardo,² Nicola De Stefano,³ Julie Aldridge,⁴ Sana Syed,⁴ Yann Hyvert,⁵ Andrew Galazka,⁶ Caroline Lemieux,⁷ Gavin Giovannoni⁸

¹University of Ottawa and the Ottawa Hospital Research Institute, Ottawa, ON, Canada; ²Oklahoma Medical Research Foundation, Oklahoma City, OK; ³University of Siena, Siena, Italy; ⁴EMD Serono Inc, Billerica, MA; ⁵Merck KGaA, Darmstadt, Germany; ⁶Merck, Aubonne, Switzerland; ⁷EMD Serono Inc, Mississauga, ON, Canada; ⁸Blizard Institute of Cell and Molecular Science, London, United Kingdom

Background: Efficacy of cladribine tablets 3.5 mg/kg (CT3.5, cumulative dose over 2 years) has been reported in relapsing-remitting multiple sclerosis (RRMS) in the 96-week CLARITY study. Prior post hoc analyses of CLARITY found that CT3.5 treatment resulted in similar benefits across the age spectrums of the studied patients in risk reduction of relapse and odds of remaining free from disease activity. **Objectives:** This post hoc analysis further examined efficacy outcomes of CT3.5 in CLARITY patients aged ≤ 30 and > 30 years at study enrollment. **Methods:** Analyses were performed by treatment (CT3.5 vs placebo) and age subgroup (≤ 30 and > 30 years), a relatively young age cutoff with adequate N needed for analyses. End points were relapse (relapse-free status and annualized relapse rate [ARR]), 3- and 6-month confirmed disability progression (CDP, based on Expanded Disability Status Scale), magnetic resonance imaging (MRI) activity, and no evidence of disease activity (NEDA) status (no relapse, 3- or 6-month CDP, or MRI activity). P values are nominal. **Results:** This analysis was carried out in 870 patients: ≤ 30 years: CT3.5 N = 109, placebo N = 102; > 30 years: CT3.5 N = 324, placebo N = 335. In both age subgroups, CT3.5 significantly reduced adjusted ARR (95% CI) [≤ 30 years: 0.16 [0.11-0.23] vs 0.48 [0.38-0.60]; > 30 years: 0.15 [0.12-0.18] vs 0.31 [0.27-0.37]; $P < .0001$], and increased the number of patients who were relapse-free through week 96 (≤ 30 years: 73.4% vs 44.1%; > 30 years: 76.2% vs 57.3%) vs placebo. CT3.5 treatment increased the odds of being free from 3- or 6-month CDP through week 96 [3-month CDP at week 96 in both age subgroups: 0.84-0.88 vs 0.76 [both subgroups]; 6-month CDP at week 96: 0.89-0.94 vs 0.83-0.86] vs placebo. CT3.5 also significantly reduced the adjusted mean (95% CI) cumulative number of new T1 gadolinium-enhancing [≤ 30 years: 0.22 [0.15-0.33] vs 1.37 [0.97-1.93]; > 30 years: 0.05 [0.03-0.08] vs 0.77 [0.61-0.98]] and active T2 [≤ 30 years: 0.68 [0.53-0.88] vs 2.20 [1.73-2.81]; > 30 years: 0.26 [0.21-0.32] vs 1.19 [1.01-1.41]] lesions, and increased the number of patients achieving NEDA status using 3-month [≤ 30 years: 30.3% vs 2.9%; > 30 years: 44.4% vs 17.3%] or 6-month [≤ 30 years: 30.3% vs 2.9%; > 30 years: 46.3% vs 17.9%] CDP vs placebo (all $P < .0001$) at 96 weeks. **Conclusions:** CT3.5 treatment improved clinical and MRI outcomes in both younger and older patients in CLARITY. Relapse and disability outcomes appeared mostly similar between the age subgroups; however, the > 30 years subgroup appeared to have a greater reduction in MRI lesion activity and a higher rate of achieving NEDA status.

Supported by: None

Disclosure: *Mark S. Freedman:* Actelion, Bayer HealthCare, Biogen, Chugai, EMD Canada, Genzyme, F. Hoffmann-La Roche, Novartis, Sanofi, Teva Pharmaceuticals USA (consulting fee). *Gabriel Pardo:* AbbVie, Adamas, Alkermes, Sanofi Genzyme, Teva (research support); Alexion, Celgene, Sanofi Genzyme (consulting fee); Biogen, EMD Serono, Novartis, Roche/Genentech (consulting fee, research support). *Nicola De Stefano:* Bayer Schering AG, Merck Serono SA, Novartis Pharma AG, Sanofi-Aventis, Serono Symposia International Foundation, Teva Pharmaceutical Industries (consulting fee). *Julie Aldridge:* Sana Syed, *Caroline Lemieux:* EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). *Yann Hyvert:* Merck KGaA (salary). *Andrew Galazka:* Merck KGaA, Aubonne, Switzerland (a business of Merck KGaA, Darmstadt, Germany) (salary). *Gavin Giovannoni:* AbbVie, Actelion, Almirall, Atara Bio, Bayer Schering Pharma, Five Prime, GlaxoSmithKline, GW Pharma, Merck KGaA, Pfizer Inc, Protein Discovery Laboratories, Sanofi Genzyme, Teva Pharmaceuticals Industries Ltd, UCB, Vertex Pharmaceuticals (consulting fee); Biogen, Ironwood, Merck and Co, Novartis (consulting fee, unrelated research support).

Keywords: Disease-modifying treatments in MS

(DXT50) Prevalence of Serious Adverse Pregnancy Outcomes After Exposure to Interferon Beta Before or During Pregnancy: Stratification by Characteristics of Pregnant Women with Multiple Sclerosis in a Register-Based Cohort Study in Finland and Sweden

Marta Korjagina,¹ Katja Hakkarainen,¹ Sarah Burkill,² Yvonne Geissbuehler,³ Meritxell Sabido,⁴ Achint Kumar,⁵ Kiliana Suzart-Woischnik,⁶ Riho Klement,¹ Jan Hillert,² Auli Verkkoniemi-Ahola,^{7,8} Shahram Bahmanyar,⁹ Scott Montgomery,^{2,9,10} Pasi Korhonen,¹ European Interferon Beta Pregnancy Study Group and Nordic MS Pregnancy & Interferon Beta Study Group

¹StatFinn and EPID Research (an IQVIA company), Espoo, Finland; ²Karolinska Institute, Stockholm, Sweden; ³Novartis Pharma AG, Basel, Switzerland; ⁴Merck KGaA, Darmstadt, Germany; ⁵Biogen, Cambridge, MA; ⁶Bayer AG, Berlin, Germany; ⁷University of Helsinki, Helsinki, Finland; ⁸Helsinki University Hospital, Helsinki, Finland; ⁹Orebro University, Orebro, Sweden; ¹⁰University College London, London, United Kingdom

Background: A recent cohort study in women with multiple sclerosis (MS) reported no increase in the prevalence of adverse pregnancy outcomes after exposure to interferon beta (IFNβ) before or during pregnancy. However, differing prevalence by maternal characteristics is unknown. **Objectives:** To describe the prevalence of serious adverse pregnancy outcomes (SAPOs) among pregnant women with MS exposed to only IFNβ and those unexposed to any MS disease-modifying drugs, with stratification by maternal characteristics. **Methods:** This cohort study extracted register data from Finland (1996-2014) and Sweden (2005-2014) on pregnant women with MS who were 1) dispensed only IFNβ within 6 months before the last menstrual period (LMP) or during pregnancy (IFNβ-exposed, n = 718 pregnancies) and 2) without dispensed MS disease-modifying drugs (unexposed, n = 1397 pregnancies). The prevalence (%) of SAPOs (consisting of elective terminations due to fetal anomaly, major congenital anomalies in live birth, and stillbirth) with 95% CIs was analyzed with stratification by maternal characteristics at LMP: time since MS diagnosis, duration of MS treatment, maternal age, and presence of chronic disease. **Results:** The prevalence of SAPOs appeared similar among the IFNβ-exposed and unexposed groups when MS was diagnosed ≤2 years [0.9% [95% CI 0.1%-3.2%] vs 3.0% [1.6%-5.2%]] or 3-5 years [2.4% [0.9%-5.1%] vs 6.0% [4.0%-8.6%]] before LMP, and was comparable for >5 years [3.3% [1.4%-6.4%] vs 3.0% [1.7%-4.8%]]. When stratified by duration of MS treatment, the prevalence among the IFNβ-exposed vs unexposed with ≤2-year treatment was 1.3% (0.4%-3.4%) vs 4.6% (2.9%-6.9%), 3- to 5-year treatment 1.7% (0.5%-4.4%) vs 4.9% (2.9%-7.7%), and >5-year treatment 4.3% (1.9%-8.3%) vs 2.7% (1.2%-5.0%). The prevalence was similar among the IFNβ-exposed vs unexposed in strata by maternal age (≤20, 21-25, 26-30, 31-35, 36-40, >40 years) and presence of chronic diseases (yes/no). **Conclusions:** In this population-based observational study, the descriptive prevalence of SAPOs appeared similar with IFNβ exposure before or during pregnancy, when pregnant women with MS were stratified by maternal characteristics.

Supported by: None

Disclosure: *Marta Korjagina, Katja Hakkarainen, Riho Klement, Pasi Korhonen:* EMD Serono (contracted research); *StatFinn-EPID Research (salary), Sarah Burkill, Shahram Bahmanyar:* Centre for Pharmacoepidemiology (salary). *Yvonne Geissbuehler:* Novartis Pharma AG (salary). *Meritxell Sabido:* Merck KGaA (salary). *Achint Kumar:* Biogen (salary). *Kiliana Suzart-Woischnik:* Bayer AG (salary). *Jan Hillert:* Bayer Schering, Biogen, Merck Serono, Novartis, Sanofi Genzyme, Teva (consulting fee, speakers' bureau); Bayer Schering, Biogen, Merck Serono, Novartis, Sanofi Genzyme, Teva, Swedish Research Council, Swedish Brain Foundation (contracted research). *Auli Verkkoniemi-Ahola:* Biogen, Genzyme, Grifols, Merck, Novartis, Roche, Sanofi (consulting fee, speakers' bureau); Sanofi (contracted research). *Scott Montgomery:* AstraZeneca, IQVIA, Novartis, Roche (contracted research); Teva (speakers' bureau).

Keywords: Disease-modifying treatments in MS, MS and the caregiver/family, Pregnancy and MS

(DXT51) High Rates of Adherence to Oral Droximel Fumarate and Dimethyl Fumarate Are Observed and Sustained in Relapsing Multiple Sclerosis Patients

Mary Kay Fink,¹ Elzbieta Jasinska,² Pavle Repovic,³ Cortnee Roman,⁴ Hailu Chen,⁵ Shivani Kapadia,⁶ Sibyl Wray⁶

¹Department of Neurology, Washington University School of Medicine, St. Louis, MO; ²Collegium Medicum UJK and Clinical Center, RESMEDICA, Kielce, Poland; ³Multiple Sclerosis Center, Swedish Neuroscience Institute, Seattle, WA; ⁴Rocky Mountain Multiple Sclerosis Clinic, Salt Lake City, UT; ⁵Biogen, Cambridge, MA; ⁶Hope Neurology, Knoxville, TN

Background: Adherence to therapy in multiple sclerosis (MS) leads to improved clinical outcomes. Persistence to therapy and discontinuation rates contribute to adherence. Droximel fumarate (DRF) is a novel oral fumarate recently approved in the United States for relapsing forms of MS, administered as 2 capsules twice daily. **Objectives:** To compare adherence with DRF vs dimethyl fumarate (DMF) in EVOLVE-MS-2 and describe long-term adherence and efficacy outcomes with DRF in EVOLVE-MS-1. **Methods:** Adherence was assessed in patients with relapsing-

remitting MS who received DRF or DMF in the 5-week, randomized EVOLVE-MS-2 study (trial registration: NCT03093324) and in those who received DRF in the ongoing, 96-week, open-label EVOLVE-MS-1 study (NCT02634307). All EVOLVE-MS-2 patients received 2 capsules twice daily. Adherence was determined by pill count (number taken divided by the number prescribed × 100). In patients who discontinued treatment, adherence was based on duration of treatment and not the full study period. Efficacy outcomes were assessed in EVOLVE-MS-1. **Results:** In the completed EVOLVE-MS-2 study, mean adherence was high (DRF 97.1%, n = 252; DMF 97.0%, n = 251); 97.4% of patients were ≥80% adherent to therapy. Discontinuation rates were lower with DRF (3.2%) than DMF (7.2%), primarily due to differences in gastrointestinal tolerability. In EVOLVE-MS-1 as of November 30, 2018 (median [range] exposure, 84 [0-100] weeks; n = 888), mean adherence was 93.4% (n = 877); 92.0% of patients were ≥80% adherent. Overall, 16.3% of patients discontinued treatment. Median (range) exposure for patients with ≥80% vs <80% adherence was 84 (1-100) and 22 (0-97) weeks, respectively. In patients with ≥80% adherence, adjusted annualized relapse rate was 0.15 (95% CI 0.13-0.19), representing a 79.4% (95% CI 75.0-83.0; P < .0001) reduction from the 12 months before study entry. Mean (SD) change in Expanded Disability Status Scale score from baseline to week 96 in patients with ≥80% adherence was 0.07 (0.59; n = 310). The small sample size (n = 70 at baseline, n = 2 at week 96) of patients with <80% adherence limits the ability to draw conclusions on the correlation between adherence and efficacy outcomes. **Conclusions:** High adherence and low discontinuation rates demonstrate documented treatment adherence and persistence to DRF. Adherence rates with DRF and DMF in EVOLVE-MS-2 were sustained in DRF-treated patients from EVOLVE-MS-1 for up to 96 weeks, demonstrating little impact of a 2-capsules-twice-daily dosing regimen. DRF-treated patients in EVOLVE-MS-1 had improved efficacy outcomes from baseline.

Supported by: Biogen

Disclosure: *Mary Kay Fink:* Biogen, EMD Serono, Novartis (speakers' bureau); *Biogen, Novartis, Sanofi Genzyme (scientific advisory board), Elzbieta Jasinska:* Biogen (scientific advisory boards); *Novartis, Allergan, F. Hoffman La-Roche, Teva, Adamed, Polfarma (speakers' bureau), Pavle Repovic:* Alexion, Biogen, Celgene, EMD Serono, Novartis, Sanofi Genzyme, Viela (consulting fee); *Alexion, Biogen, Genzyme, Genentech, EMD Serono (speakers' bureau); Biogen, Genentech, EMD Serono (contracted research), Cortnee Roman:* Alexion, Biogen, Bristol Myers Squibb, Genentech, Novartis, Sanofi Genzyme (speakers' bureau); *Alexion, Biogen, Bristol Myers Squibb, EMD Serono, Genentech, Novartis, Sanofi Genzyme (consulting fee); Biogen, Bristol Myers Squibb, Genentech, Novartis, Sanofi Genzyme (advisory boards), Hailu Chen, Shivani Kapadia:* Biogen (ownership interest, salary), *Sibyl Wray:* Alkermes Inc, Biogen, Celgene, Genentech/Roche, Sanofi Genzyme, Novartis, TG Therapeutics (contracted research); *Biogen, EMD Serono, Genentech/Roche, Sanofi Genzyme (consulting fee, speakers' bureau).*

Keywords: Disease-modifying treatments in MS, Medication adherence

(DXT52) Efficacy and Safety of Eculizumab in Patients with Neuromyelitis Optica Spectrum Disorder Previously Treated with Rituximab: The Phase 3 PREVENT Study

Michael Levy,^{1,2} Achim Berthele,³ Ho Jin Kim,⁴ Kazuo Fujihara,^{5,6,7} Ichiro Nakashima,^{6,8} Celia Oreja-Guevara,⁹ Jacqueline Palace,¹⁰ Sean J. Pittock,¹¹ Murat Terzi,¹² Natalia Totolyan,¹³ Shanthi Viswanathan,¹⁴ Kai-Chen Wang,^{15,16} Amy Pace,¹⁷ Marcus Yountz,¹⁷ Diane Lawson,¹⁷ Eva Laudon-Meyer,¹⁸ Dean Wingerchuk¹⁹

¹Massachusetts General Hospital and Harvard Medical School, Boston, MA; ²Johns Hopkins University, Baltimore, MD; ³Technical University of Munich, Munich, Germany; ⁴National Cancer Center, Goyang, Korea, Republic of (South); ⁵Southern TOHOKU Research Institute for Neuroscience (STRINS), Koriyama, Japan; ⁶Tohoku University, Sendai, Japan; ⁷Fukushima Medical University, Fukushima City, Japan; ⁸Tohoku Medical and Pharmaceutical University, Sendai, Japan; ⁹Hospital Universitario Clinico San Carlos, Madrid, Spain; ¹⁰John Radcliffe Hospital, Oxford, United Kingdom; ¹¹Mayo Clinic, Rochester, MN; ¹²On dokuz Mayis University, Samsun, Turkey; ¹³First Pavlov State Medical University of St Petersburg, St Petersburg, Russian Federation; ¹⁴Kuala Lumpur Hospital, Kuala Lumpur, Malaysia; ¹⁵National Yang Ming University, Taipei, Taiwan; ¹⁶Cheng-Hsin General Hospital, Taipei, Taiwan; ¹⁷Alexion Pharmaceuticals, Boston, MA; ¹⁸Alexion Pharmaceuticals, Zurich, Switzerland; ¹⁹Mayo Clinic, Scottsdale, AZ

Background: In the PREVENT study, eculizumab was associated with a significant reduction in relapse risk vs placebo and was well tolerated. In total, 46 patients (26/96 receiving eculizumab, 20/47 receiving placebo) had been previously treated with the monoclonal antibody rituximab. **Objectives:** To describe the efficacy and safety of eculizumab in patients in the randomized, double-blind, placebo-controlled, phase 3 PREVENT trial (trial registration: NCT01892345) who had previously received rituximab. **Methods:** Adults with aquaporin-4 immunoglobulin G-positive neuromyelitis optica spectrum disorder (NMOSD) received eculizumab (maintenance dose, 1200 mg/2 weeks) or placebo with/

without concomitant immunosuppressive treatment (except rituximab/mitoxantrone). A post hoc descriptive analysis was performed using data from patients with any prior rituximab treatment (within the previous year only for review of adverse events [AEs]) recorded more than 3 months before randomization. **Results:** Baseline characteristics of the prior-rituximab subgroup were similar to the total PREVENT population; however, the subgroup included a lower proportion of Asian patients (10.9% vs 36.4% in total PREVENT study population) and greater representation from the Americas (58.7% vs 30.8%). In the subgroup, median times from last dose of rituximab to meningococcal vaccination and to first dose of study treatment were 31.7 and 38.7 weeks, respectively. Adjudicated relapses occurred in 1/26 (3.8%) and 7/20 (35.0%) patients in the eculizumab and placebo arms, respectively (hazard ratio 0.093; 95% CI 0.011-0.755; $P = .0055$). AE rates in patients receiving eculizumab and placebo within 1 year of previous rituximab use were 1025.8 and 1029.1 events/100 patient-years (both 100% of patients), respectively; rates of serious AEs were 46.9 and 66.0 events/100 patient-years (38.9% and 47.1% of patients), respectively. Serious infections/infestations were recorded in 2/18 (11.1%) and 2/17 (11.8%) patients in the eculizumab and placebo arms, respectively. **Conclusions:** In patients with aquaporin-4 immunoglobulin G-positive NMOSD in PREVENT who had previously received rituximab, the risk of adjudicated relapse was significantly lower with eculizumab than with placebo. Rates of serious infections were similarly low with eculizumab and placebo.

Supported by: None

Disclosure: *Michael Levy:* Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent); Genentech, Quest Diagnostics, Viela Bio (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent). *Achim Berthele:* Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent, speakers' bureau). *Ho Jin Kim:* Alexion Pharmaceuticals, Celtrion, Eisai, HanAll BioPharma, Merck Serono, Novartis, Sanofi Genzyme, Teva-Handok, Viela Bio (consulting fee); Journal of Clinical Neurology (associate editor); MedImmune/Viela Bio (serves on a steering committee); Multiple Sclerosis Journal (co-editor). *Kazuo Fujihara:* Alexion Pharmaceuticals, Biogen, Chugai, Mitsubishi-Tanabi, Novartis (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent); Asahi Medical, Bayer, Eisai, Roche, Teijin (fees for non-CME/CE services received directly from commercial interest or its agent). *Ichiro Nakashima:* Alexion (consulting fee). *Celia Oreja-Guevara:* Alexion Pharmaceuticals, Merck, Novartis, Roche, Sanofi (consulting fee, contracted research). *Jacqueline Palace:* Biogen, Chugai (contracted research); LEK, Viela Bio, Guidepoint (consulting fee); Merck Serono (meeting/lecture/workshop participation); Novartis, Roche, Argenx (speakers' bureau); UCB, Viela Bio, Roche (conference/lecture participation). *Sean J. Pittock:* Alexion Pharmaceuticals (all compensation paid directly to Mayo Clinic, consulting fee, contracted research); Astellas (all compensation paid directly to Mayo Clinic, consulting fee); Autoimmune Encephalitis Alliance (all compensation paid directly to Mayo Clinic, contracted research); Grifols (all compensation paid directly to Mayo Clinic, contracted research); MedImmune (all compensation paid directly to Mayo Clinic, consulting fee, contracted research); UCB (consulting fee, personal compensation for attending UCB advisory board meeting in Stockholm, Sweden, on Sept 10, 2019). *Murat Terzi, Shanthi Viswanathan, Kai-Chen Wang:* Nothing to disclose. *Natalia Tobbyan:* Alexion, Janssen, Novartis, Roche, Sanofi, Receptos Inc, Biocad (Russia) (contracted research); Merck (consulting fee); Roche, Sanofi (fees for lectures). *Amy Pace, Marcus Yountz, Eva Laudon-Meyer:* Alexion Pharmaceuticals (ownership interest, salary). *Diane Lawson:* Alexion Pharmaceuticals (salary). *Dean Wingerchuk:* MedImmune, Novartis, Biogen, Celgene, Genentech, TG Therapeutics, Arcus Medica, Reistone (consulting fee); Terumo BCT, Alexion Pharmaceuticals (fees for non-CME/CE services received directly from commercial interest or its agent).

Keywords: Eculizumab in NMOSD

(DXT53) Multiple Sclerosis Clinical Phenotypes: Using Technology to Educate Patients and Optimize Treatment

Elaine Rudell,¹ Patty Peterson,¹ Andrea Griffin,² Michele F. Ingram,¹ Michelle Fabian³

¹@Point of Care, Livingston, NJ; ²Multiple Sclerosis Association of America, Cherry Hill, NJ; ³Icahn School of Medicine, Corinne Goldsmith Dickinson Center for Multiple Sclerosis, New York, NY

Background: The evolution of new clinical phenotype descriptions for relapsing and progressive forms of multiple sclerosis (MS) by the MS Phenotype Group and new guideline recommendations for disease-modifying therapies have delineated optimal treatments for each clinical phenotype. The Multiple Sclerosis Association of America (MSAA) and @Point of Care have collaborated to develop complementary apps for use by patients and their clinicians that facilitate sharing of data: My

MS Manager, a HIPAA-compliant patient app, and Multiple Sclerosis @ Point of Care, a clinician app. It was important to enhance the My MS Manager app to provide education addressing MS phenotypes and their implications, and a field for patients to enter their phenotype. **Objectives:** Encourage patients with MS to know their phenotype, clinical course, and implications for treatment for discussion with their clinician and caregivers. **Methods:** MSAA and @Point of Care collaborated to enhance the My MS Manager app to include additional fields for the collection of clinical phenotypes, to assess how many did/did not know their clinical phenotype and evaluate other patterns. A video was developed to increase patients' awareness about the importance of knowing their clinical phenotype and how this guides treatment options. **Results:** Of 3765 patients who filled in their clinical phenotype on the My MS Manager app: 20% (744) did not know their clinical phenotype and those unsure of their phenotype were younger (*Implication:* Patients, especially those younger, need to understand the importance of knowing their phenotype and its implications for long-term treatment); thousands of patients with MS viewed the educational videos addressing MS and clinical phenotype information (*Implication:* Patients want to gain insights into their MS and clinical phenotypes); and of 55 patients with primary progressive MS (PPMS), 25% were not receiving a US Food and Drug Administration (FDA)-approved therapy nor were they enrolled in a PPMS clinical trial (*Implication:* Better informed patients with PPMS and clinicians may opt for FDA-approved therapies or clinical trials). **Conclusions:** The My MS Manager patient app facilitates MS patients' ability to record their clinical phenotype, access educational videos (including those addressing phenotype), and empower them to better understand MS phenotypes and implications for treatment choices for discussion with their clinicians. This will ensure optimal treatment choices.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Disease-modifying treatments in MS, MS phenotypes

(DXT54) Assessment of the Discontinuation Rates of Disease-Modifying Therapy in Veterans with Multiple Sclerosis

Naomi Wu,¹ Eric Williamson,² Jeremy Liu,¹ Christine Lava,¹ Hyojin Suh,¹ Andrea Hanssen,² Sunita Dergalust¹

¹Pharmacy and ²Neurology, VA Greater Los Angeles Healthcare System, Los Angeles, CA

Background: Disease-modifying therapy (DMT) is the preferred treatment approach for multiple sclerosis (MS) and has been shown to reduce the rate of relapse and slow disease progression. Poor adherence is associated with an increased risk of relapse leading to an increase in morbidity outcomes as well as overall costs in patients with MS. We conducted a retrospective chart review to improve our understanding of the reasons for discontinuation of DMTs and the impact of adverse drug reactions on the rate discontinuation of DMTs and to identify barriers to adherence.

Objectives: To examine the discontinuation rates and the reasons for discontinuation of DMTs among veterans with MS. **Methods:** We conducted a retrospective chart review of veterans with MS seen in the MS Clinic and General Neurology Clinic at Veterans Affairs Greater Los Angeles Healthcare System who were on DMTs from January 1, 2010, to December 31, 2019. Demographic data and the following data points were collected: past medical history, date of diagnosis, duration of MS, characteristics related to DMT use such as prescription refill history, reason for discontinuation, duration of medication use, and response to DMTs. **Results:** To date we have screened 100 electronic medical records of veterans with MS on DMTs. We documented 220 trials of DMT in 100 patients enrolled in our study. Among these 100 patients, the most commonly prescribed DMTs were interferon-beta and glatiramer acetate. Adherence rates observed were highest among veterans on infused DMTs and lowest among veterans on injectable DMTs. Approximately 20% of patients discontinued the injectable DMTs due to inefficacy, compared to 10% for oral DMTs and 10% for infused DMTs. Injectable DMTs were discontinued in 20% of veterans due to adverse drug reactions, compared to 18% for oral DMTs and 4% for infused DMTs. Data collection is ongoing and may help us identify barriers to DMT adherence in veterans with MS. **Conclusions:** Preliminary results of our study suggest differences in adherence to DMTs and possible reasons for discontinuation of DMTs in veterans with MS.

Supported by: None

Disclosure: *Naomi Wu, Jeremy Liu, Christine Lava, Hyojin Suh, Andrea Hanssen, Sunita Dergalust:* Nothing to disclose. *Eric Williamson:* Novartis Pharmaceuticals (contracted research).

Keywords: Disease-modifying treatments in MS, Medication adherence

(DXT55) Herpes Zoster Virus (HZV) Infections Among Multiple Sclerosis Patients Treated with Various Disease-Modifying Therapies

Nicola Carlisle,¹ Sam I. Hooshmand,¹ Michelle Maynard,² Leah Hoffman,³ Ahmed Z. Obeidat⁴

¹Neurology, Medical College of Wisconsin and Affiliated Hospitals, Milwaukee, WI; ²Neurosciences, Froedter and the Medical College of Wisconsin, Milwaukee, WI;

³Pharmacy, Froedter and the Medical College of Wisconsin, Milwaukee, WI; ⁴Neurology, Medical College of Wisconsin, Milwaukee, WI

Background: Disease-modifying therapies (DMTs) for multiple sclerosis (MS) may increase the risk for opportunistic infections, including herpes zoster (HZV). The relative frequency of HZV infection in the treated MS population is unknown. Furthermore, the relative distribution of reported cases per age group and gender is unknown. **Objectives:** To stratify the frequency of voluntarily reported HZV infections by DMT, age, and gender in patients with MS. **Methods:** We queried the Food and Drug Administration Adverse Event Reporting System (FAERS) for adverse events ("herpes zoster" and "varicella") reported in patients with MS between January 1999 and June 2019 receiving interferon beta (IFN β), glatiramer acetate (GA), natalizumab (NAT), fingolimod (FIN), teriflunomide (TER), dimethyl fumarate (DMF), alemtuzumab (ALE), and ocrelizumab (OCR). We excluded reports where the "suspect drug" included 2 or more DMTs. We stratified the reports for each DMT, by year of report, age, and gender. **Results:** 3352 reports met our inclusion criteria. Mean (SD) annual report rates were highest for patients treated with NAT at 81.9 (91.9) and lowest for GA at 2.1 (2.5). Other DMTs: FIN 70.3 (27.3); DMF 66 (21); OCR 55.3 (27.8); ALE 22.8 (15); IFN β 22.6 (18); and TER 10.4 (4.7). Reports were 4.7x more in females (ranging from 2.3x for ALE to 8.2x for IFN β). The highest percentage of reports was in the sixth decade of life for all DMTs except ALE (fourth decade). Several reports were in individuals younger than 40 (25.0%). **Conclusions:** Reports of HZV infections varied based on the DMT used, patient age, and gender. HZV reports were nearly fivefold more frequent in females than males, and reports among patients younger than 40 were higher than expected. Database limitations precluded calculations of incidence. We encourage further investigations of the incidence and risk mitigation strategies (including vaccination practices) of HZV in patients with MS on DMTs regardless of age of the patient.

Supported by: None

Disclosure: Nicola Carlisle, Sam I. Hooshmand, Michelle Maynard, Leah Hoffman: Nothing to disclose. Ahmed Z. Obeidat: Alexion, Biogen (consulting fee, speakers' bureau); Celgene, EMD Serono, Genentech, Sanofi (consulting fee); International Journal of MS Care (editorial board); Novartis (speakers' bureau).

Keywords: Disease-modifying treatments in MS

(DXT56) Potential Weight Changes Among Patients with Multiple Sclerosis Undergoing Treatment with Ocrevus (Ocrelizumab)

Olivia Wei,¹ Elizabeth S. Gromisch,^{2,3,4} Lindsay O. Neto,^{2,5} Jennifer A. Ruiz,^{2,4} Peter Wade^{2,3}

¹Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT;

²Mandell Center for Multiple Sclerosis, Mount Sinai Rehabilitation Hospital, Trinity Health of New England, Hartford, CT; ³Department of Neurology, University of Connecticut School of Medicine, Farmington, CT; ⁴Departments of Rehabilitative Medicine and Medical Sciences, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT; ⁵Department of Rehabilitative Medicine, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT

Background: Ocrelizumab (Ocrevus), approved for use in 2017, has quickly become a formidable disease-modifying therapy option for multiple sclerosis (MS). While effective, potential adverse effects from ocrelizumab are still being explored, such as weight changes. **Objectives:** 1) To examine if patients have had weight changes while receiving ocrelizumab and 2) to explore associations between changes in weight and patients' demographic and clinical characteristics. **Methods:** Data were extracted from the medical records of 152 patients who had recorded weights before and after receiving ocrelizumab, including their weight in kilograms, age, gender, race, ethnicity, and MS subtype, duration, and disability [Expanded Disease Status Scale [EDSS]]. There was an average of 10.96 \pm 5.76 months between the 2 weight measurements. Changes in weight were examined using a 1-sample Wilcoxon signed rank test, while associations between weight changes and demographic and clinical variables were assessed with Spearman rho (ρ) correlations and Mann-Whitney U tests. **Results:** Over 52% of patients (n = 80) gained weight after receiving ocrelizumab (median: 2.36 kg), while nearly 39% (n = 59) lost weight (median: -1.90 kg). In the overall sample (n = 152), there was a significant change from pretreatment weight (z = 2.51, P =

.012), with a median change of 0.37 kg (range: -7.72 to 17.30 kg). Changes in weight were negatively correlated with EDSS scores (p = -0.18, P = .045), with patients having weight gain having a median EDSS score of 3 (range: 0-7.0) and patients having weight loss having a median EDSS score of 4.0 (range: 1.5-7.5). There were no other significant associations. **Conclusions:** Findings from this preliminary study suggest that weight changes after ocrelizumab are frequently seen, with most patients either gaining or losing weight. As the EDSS score was the only variable associated with weight changes, further investigation is warranted to understand the underlying phenomenon, as well as the normal distribution of weight changes for persons with MS under more controlled time frames and across all levels of EDSS.

Supported by: None

Disclosure: Olivia Wei, Elizabeth S. Gromisch, Lindsay O. Neto, Jennifer A. Ruiz: Nothing to disclose. Peter Wade: Biogen, Celgene, EMD Serono, Genentech, Mallinckrodt, Novartis, Sanofi Genzyme (speakers' bureau).

Keywords: Disease-modifying treatments in MS, Weight changes

(DXT57) FAST: Faster and Safe Administration of Tysabri

Patricia A. Pagnotta

Neurology Associates, PA, and MS Center of Greater Orlando, Maitland, FL

Background: Natalizumab (Tysabri) has been US Food and Drug Administration approved since 2004, and the incidence of infusion reactions is extremely low and has not increased over the 10 years of observational study done by Biogen. Patients sometimes comment on the burden of being infused monthly and being in the infusion center for 2 hours. In fact, many patients refuse the hour of observation after they are comfortable with infusions. Therefore, is it possible to reduce the amount of infusion time for natalizumab safely? **Objectives:** To determine if natalizumab can safely be given over 30 minutes vs the standard 60-minute intravenous infusion. **Methods:** Observational study of patients in MS Center of Greater Orlando's Infusion Center who consented to be infused over 30 minutes. Patients who were recruited for study had been on natalizumab for greater than 6 months. **Results:** Of 25 patients, 22 (88%) did very well with no effect. In the other 3 persons effects were mild, and they did not require any additional treatment. **Conclusions:** Administering natalizumab over 30 minutes is a reasonable and safe option for most patients. Further studies are suggested to ensure validity.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Disease-modifying treatments in MS, Tolerability

(DXT58) Reduction of Risk of Secondary Progressive Multiple Sclerosis within 2 Years of Treatment with Cladribine Tablets: An Analysis of the CLARITY Study

Patrick Vermersch,¹ Gavin Giovannoni,^{2,3} Per Soelberg-Sorensen,⁴ Kottil Rammohan,⁵ Stuart Cook,⁶ Birgit Keller,⁷ Sanjeev Roy⁸

¹University of Lille, Lille, France; ²Blizard Institute of Cell and Molecular Science, London, United Kingdom; ³Queen Mary University of London, London, United Kingdom;

⁴Department of Neurology, Danish MS Center, University of Copenhagen, Copenhagen, Denmark; ⁵Department of Neurology, Multiple Sclerosis Center, University of Miami, Miami, FL; ⁶Rutgers, The State University of New Jersey, New Jersey Medical School, Newark, NJ; ⁷Merck KGaA, Darmstadt, Germany; ⁸Merck KGaA, Aulendorf, Switzerland

Background: Cladribine tablets (CTs) 10mg, cumulative dose 3.5 mg/kg (CT3.5; N = 433) over 2 years showed efficacy vs placebo (N = 437) in patients with relapsing multiple sclerosis (MS) in the CLARITY study. **Objectives:** Explore (post hoc) the relationship between baseline Expanded Disability Status Scale (EDSS) score and risk of progression to secondary progressive MS (SPMS) or to EDSS score \geq 6.0 in CLARITY. **Methods:** As progression to SPMS was not recorded during the trial, a proxy composite definition was used: confirmed disability progression (CDP), CDP within the leading EDSS-defined functional score (FS), EDSS score postbaseline \geq 4.0, pyramidal FS \geq 2, all conditions met for at least 3 months (mo) in the absence of a relapse. Patients progressing to EDSS score \geq 6.0 were defined by having \geq 1 postbaseline EDSS score \geq 6.0 with 3- or 6-month CDP. In this post hoc analysis, odds ratios (ORs) and corresponding CIs are estimated by a logistic regression model with treatment and baseline EDSS score (\leq 3.0 or \geq 3.5) as fixed effects. **Results:** Overall, proxy SPMS progression was seen in 6.7% of CT3.5 patients vs 13.5% for placebo (OR 0.46 [95% CI: 0.28, 0.76]; P = .0024). In the baseline EDSS \leq 3.0 subgroup (CT3.5 n = 257; placebo n = 235), proxy SPMS progression occurred in 3.5% vs 7.7% (CT3.5 vs placebo; OR 0.44 [95% CI: 0.19, 0.99]; P = .0471). In the baseline EDSS \geq 3.5 subgroup (CT3.5 n = 148; placebo n = 157), proxy SPMS progression occurred in 12.2% vs 22.4% (CT3.5 vs placebo; OR 0.48 [95% CI: 0.26, 0.9]; P = .0212). Similar effects have been observed for each

proxy SPMS component vs placebo. Proportions of patients with at least 1 EDSS value ≥ 6.0 postbaseline were 6.4% vs 14.5% (CT3.5 vs placebo; OR 0.4 [95% CI: 0.24, 0.66]; $P = .0004$). Corresponding proportions for patients with 3-month CDP with EDSS score ≥ 6.0 were 3.5% vs 8.0% (CT3.5 vs placebo; OR 0.42 [95% CI: 0.22, 0.82]; $P = .0114$) and for patients with 6-month CDP with EDSS ≥ 6.0 were 2.8% vs 5.8% (CT3.5 vs placebo; OR 0.48 [95% CI: 0.22, 1.02]; $P = .0566$). Subgroup analysis by baseline EDSS score showed that in patients with baseline EDSS score ≤ 3.0 , 0.8% vs 4.3% had at least 1 EDSS score ≥ 6.0 (CT3.5 vs placebo; OR 0.18 [95% CI: 0.04, 0.81]; $P = .0262$). In patients with baseline EDSS score ≥ 3.5 , the corresponding proportions were 16.2% vs 29.9% (CT3.5 vs placebo; OR 0.45 [95% CI: 0.26, 0.79]; $P = .0051$). **Conclusions:** The risks of progressing to SPMS (proxy) within 2 years of treatment, or having EDSS score ≥ 6.0 , are significantly reduced with CT3.5 vs placebo, regardless of baseline EDSS score (≤ 3.0 or ≥ 3.5).

Supported by: None

Disclosure: *Patrick Vermersch:* Almirall, Celgene, Novartis, Roche (consulting fee); Biogen, Merck KGaA, Sanofi Genzyme, Teva (consulting fee, research support). *Gavin Giovannoni:* AbbVie, Actelion, Almirall, Atara Bio, Bayer Schering Pharma, Five Prime, GlaxoSmithKline, GW Pharma, Merck KGaA, Pfizer Inc, Protein Discovery Laboratories, Sanofi Genzyme, Teva Pharmaceuticals Industries Ltd, UCB, Vertex Pharmaceuticals (consulting fee); Biogen, Ironwood, Merck and Co, Novartis (consulting fee, unrelated research support). *Per Soelberg-Sorensen:* Biogen (advisory board, research support); Biogen, Sanofi-Aventis (speakers' bureau); Genzyme (research support, speakers' bureau); GlaxoSmithKline (advisory board, steering committees); MedDay Pharmaceuticals, Merck HealthCare KGaA (advisory board); Merck KGaA (research support, speakers' bureau, steering committees); Novartis, Teva (advisory board, research support, speakers' bureau, steering committees); Roche (research support). *Kottil Rammohan:* Acorda, Biogen, EMD Serono, Genzyme, Novartis, Roche/Genentech, Sanofi-Aventis, Teva Neurosciences (speakers' bureau). *Stuart Cook:* Actinobac Biomed Inc, Biogen, Merck KGaA, Teva Pharmaceuticals (advisory board); Bayer HealthCare (advisory board, grant support); Neurology Reviews, Sanofi Aventis (consulting fee). *Birgit Keller:* Merck KGaA (salary). *Sanjeev Roy:* Merck KGaA, Aubonne, Switzerland (salary).

Keywords: Disease-modifying treatments in MS

(DXT59) The CLARITY Study: Efficacy Outcomes Among Patients Who Received Disease-Modifying Drugs Prior to Treatment with Cladribine Tablets

Patrick Vermersch,¹ Kottil Rammohan,² Doris Damian,³ Dominic Jack,⁴ Gerard Harty,³ Schifflon L Wong³

¹University of Lille, Lille, France; ²Department of Neurology, Multiple Sclerosis Center, University of Miami, Miami, FL; ³EMD Serono Inc, Billerica, MA; ⁴Merck KGaA, Darmstadt, Germany

Background: Cladribine tablets (CTs) 10 mg, (cumulative dose 3.5 mg/kg [CT3.5] over 2 years) showed efficacy vs placebo in patients with relapsing-remitting multiple sclerosis (RRMS) in the pivotal phase 3 CLARITY study. The CLARITY study included patients treated with 0-2 disease-modifying drugs (DMDs) prior to study entry (patients treated with >2 DMDs prior to study entry were excluded). A total of 433 patients were randomized to CT3.5 and 437 patients to placebo. **Objectives:** To report clinical outcomes and magnetic resonance imaging (MRI) lesion counts in the subgroup of patients from CLARITY who had used a DMD at any time prior to randomization. **Methods:** Post hoc analysis of efficacy, annualized relapse rate (ARR), relapse-free rate, MRI activity, and time to 3-month and 6-month confirmed EDSS score progression (CDP) stratified by the cohort of patients who had received a prior DMD treatment before entering the CLARITY study. P values less than 0.05 were considered nominally significant. **Results:** Of those patients who received a prior DMD (interferon beta [IFN β]-1a, IFN β -1b, glatiramer acetate, or natalizumab), 110 were randomized to CT3.5 and 132 received placebo. Among patients with prior DMD use, CT3.5, compared to placebo, resulted in a nominally significant reduction in ARR (CT3.5, 0.22; placebo, 0.42; $P < .005$), a higher relapse-free rate (CT3.5, 70.4%; placebo, 55.9%; $P = .0204$), a numerically lower risk of 3-month (hazard ratio = 0.64, $P = .1589$) and 6-month (hazard ratio = 0.62, $P = .2071$) CDP, and reductions in the brain lesion counts ($P < .001$ for each type of lesion). **Conclusions:** Among patients who were pretreated with either IFN β -1a, IFN β -1b, glatiramer acetate, or natalizumab, efficacy outcomes were similar to those seen in the full CLARITY active RRMS population, wherein patients who received CT3.5 showed statistically significant improvements in efficacy outcomes compared to placebo.

Supported by: None

Disclosure: *Patrick Vermersch:* Almirall, Celgene, Novartis, Roche (consulting

fee); Biogen, Merck KGaA, Sanofi Genzyme, Teva (consulting fee, research support). *Kottil Rammohan:* Acorda, Biogen, EMD Serono, Genzyme, Novartis, Roche/Genentech, Sanofi-Aventis, Teva Neurosciences (speakers' bureau). *Doris Damian, Gerard Harty, Schifflon L Wong:* EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). *Dominic Jack:* Merck KGaA (salary).

Keywords: Disease-modifying treatments in MS

(DXT60) Correlations Between Four Common Measures of Cognition in Patients with Secondary Progressive Multiple Sclerosis

Ralph H.B. Benedict,¹ Xiangyi Meng,² Amit Bar-Or,³ Bruce A.C. Cree,⁴ Robert J. Fox,⁵ Patrick Vermersch,⁶ Wendy Su,² Ludwig Kappos⁷

¹University at Buffalo, State University of New York, Buffalo, NY; ²Novartis Pharmaceuticals Corporation, East Hanover, NJ; ³Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ⁴Department of Neurology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA; ⁵Mellen Center for Multiple Sclerosis, Cleveland Clinic, Cleveland, OH; ⁶Department of Neurology, University of Lille, CHU Lille, Lille, France; ⁷Departments of Medicine, Clinical Research, Biomedicine, and Biomedical Engineering, Neurologic Clinic and Policlinic, University Hospital Basel, University of Basel, Basel, Switzerland

Background: Cognitive impairment is common in patients with secondary progressive multiple sclerosis (SPMS). The oral Symbol Digit Modalities Test (SDMT), Paced Auditory Serial Addition Test (PASAT), and Brief Visuospatial Memory Test Revised (BVMTR) were used to track cognition changes in the EXPAND study. This analysis evaluates whether these tests are redundant or complementary in progressive disease. **Objectives:** Evaluate relationship between common cognitive tests in patients with SPMS and identify whether these tests provide unique insights on disease progression. **Methods:** EXPAND was a 36-month, randomized, placebo-controlled trial of siponimod (Mayzent) 2 mg/day in patients with SPMS. Cross-sectional, pairwise Pearson correlations (r) between SDMT, PASAT, and BVMTR total learning (-TL) and delayed recall (-DR) indices were calculated by treatment group and combined. Correlations were examined for baseline and change in scores from first postbaseline measurement to month 24. Correlations were strong (>0.6), intermediate (0.4-0.6), or weak (<0.4); $P < .05$ indicates that correlations were statistically different to 0. Analyses for hypothesis generation; no adjustment for multiple comparisons. **Results:** Data were analyzed for 1644 patients (siponimod, $n = 1098$; placebo, $n = 546$). In both treatment groups at baseline, strong correlations (0.89) were found between BVMTRL and BVMTR-DR; intermediate correlations (0.45-0.59) were observed between all other tests (all $P < .0001$). For change from baseline to month 24, strong correlations persisted between BVMTR-TL and BVMTR-DR (siponimod, 0.68; placebo, 0.72; both $P < .0001$), but correlations were weak between all other tests with both treatments (0.050-1.7; $P > .05$ in many cases). **Conclusions:** Cognitive outcomes were correlated at baseline, confirming large overlap in variance for BVMTR indices and intermediate shared variance for other measures at a single time point. Weak correlations of change to month 24 between SDMT, PASAT, and BVMTR may suggest that each test tracks different aspects of cognitive decline and/or has different test characteristics when applied repeatedly in patients with SPMS.

Supported by: None

Disclosure: *Ralph H.B. Benedict:* Acorda, Mallinckrodt, National Institutes of Health, National MS Society (contracted research); Biogen (CME speaking fees, consulting fee, contracted research); Celgene, EMD Serono (CME speaking fees, consulting fee); Genentech, Genzyme, Novartis (consulting fee, contracted research); Psychological Assessment Resources, Inc (royalty); Roche, Sanofi, Takeda, Verasci (consulting fee). *Xiangyi Meng, Wendy Su:* Novartis Pharmaceuticals Corporation (salary). *Amit Bar-Or:* Atara Biotherapeutics, Bayer, Bayhill Therapeutics, Berlex, Biogen, BioMS, Celgene/Receptos, Diogenix, Eli Lilly, F. Hoffman-La Roche, Genentech, GlaxoSmithKline, Guthy-Jackson/GGF, Immunic, Janssen/Actelion, Mapi, MedImmune, Merck/EMD Serono, Novartis, Ono Pharmacia, Roche/Genentech, Sanofi Genzyme, Teva Neuroscience, Wyeth (consulting fee). *Bruce A.C. Cree:* Akili, Alexion, Atara, Biogen, EMD Serono, Novartis, TG Therapeutics (consulting fee). *Robert J. Fox:* Actelion, EMD Serono, Genentech, Teva (consulting fee); Biogen, Novartis (consulting fee, contracted research). *Patrick Vermersch:* Biogen, Celgene, Merck, Teva (consulting fee); La Revue des Microbiotes (personal fees for serving as editor); Novartis, Roche, Sanofi Genzyme (consulting fee, contracted research). *Ludwig Kappos:* Actelion, Alkermes, Allergan, Almirall, Bayer, Biogen, Celgene, CSL Behring, df-mp, European Union, EXCEMED, GeNeuro, Genzyme, Merck, Mitsubishi Tanabe Pharma, Novartis, Pfizer, Receptos/Celgene, Roche, Roche Research Foundations, Sanofi-Aventis, Santhera, Swiss Multiple Sclerosis Society, Swiss National Research Foundation, Teva, UCB Pharma, Vianex (contracted research); Neurostatus products (license fees).

Keywords: Disease-modifying treatments in MS

(DXT61) Injection-Related Reactions with Subcutaneous Administration of Ofatumumab in Relapsing Multiple Sclerosis: Pooled Analysis of the Phase 3 ASCLEPIOS I and II Trials

Ratnakar Pingili,¹ Amit Bar-Or,² Jeffrey A. Cohen,³ Patricia K. Coyle,⁴ Anne H. Cross,⁵ Stephen L. Hauser,⁶ Ludwig Kappos,⁷ Cecile Kerloeguen,⁸ Ayan Das Gupta,⁹ Valentine Jehl,⁸ Dieter A. Haering,⁸ Krishnan Ramanathan,⁸ Martin Merschhemke⁸

¹Novartis Pharmaceuticals Corporation, East Hanover, NJ; ²Center for Neuroinflammation and Experimental Therapeutics and Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ³Department of Neurology, Mellen MS Center, Neurological Institute, Cleveland Clinic, Cleveland, OH; ⁴Stony Brook University, Stony Brook, NY; ⁵Department of Neurology, Division of Neuroimmunology, Washington University School of Medicine, St Louis, MO; ⁶University of California, San Francisco, San Francisco, CA; ⁷Neurologic Clinic and Polyclinic, Departments of Medicine, Clinical Research, Biomedicine and Biomedical Engineering, University Hospital of Basel, Basel, Switzerland; ⁸Novartis Pharma AG, Basel, Switzerland; ⁹Novartis Healthcare Pvt Ltd, Hyderabad, India

Background: Ofatumumab, the first fully human anti-CD20 monoclonal antibody, with a monthly 20 mg subcutaneous (s.c.) dosing regimen, demonstrated superior efficacy (reductions in clinical relapses by 51%-59%, disability worsening by 33%-34%, and gadolinium-enhancing lesions by 94%-98%) vs teriflunomide in the two phase 3 ASCLEPIOS I/II relapsing multiple sclerosis (RMS) trials. Injection-related reactions (IRRs) were the most common adverse events (AEs) observed. **Objectives:** To characterize the risk of IRRs (systemic and local site reactions) observed with ofatumumab in patients with RMS. **Methods:** In the pooled ASCLEPIOS I/II trials, patients were randomized (1:1) to receive s.c. ofatumumab 20 mg (n = 946) (loading dose: days 1, 7 and 14; maintenance dose: every 4 weeks from week 4) or oral teriflunomide 14 mg once daily (n = 936), for up to 30 months. Patients in the teriflunomide group received matching placebo injections. All patients received the first 4 injections at the clinic and subsequent injections at home. Premedication was recommended, but not mandatory. Both systemic (during and within 24 hours postinjection) and local site IRRs (at any time) were reported. **Results:** 20.6% (n = 195) of the patients in the ofatumumab group and 15.3% (n = 143) in the teriflunomide group had ≥1 systemic IRR. Incidence of systemic IRRs with the first injection was 14.4% with ofatumumab vs 7.5% with teriflunomide. The incidence of systemic IRRs decreased with subsequent doses and was similar to the matching placebo injections in the teriflunomide group. Most IRRs (99.8%) were grade 1/2 in severity; grade 3 IRRs were observed in 2 patients (0.2%) with ofatumumab at the first injection (1 of which was reported as a serious AE) vs none with teriflunomide. One additional IRR (grade 1) was also reported as a serious AE with ofatumumab. The serious IRRs (0.2%) were manageable, and patients continued treatment with no recurrences. No life-threatening IRRs were reported during the study. The most frequent (≥2%) IRR symptoms observed with ofatumumab were fever, headache, myalgia, chills, and fatigue. Most local site IRRs were mild to moderate in severity and nonserious in nature; the most frequently reported symptoms (≥2%) included erythema, pain, itching, and swelling. **Conclusions:** Systemic and local IRRs with ofatumumab 20 mg s.c. were mostly mild to moderate in severity. Beyond the first injection, IRRs were no more frequent with ofatumumab vs matching placebo injections.

Supported by: None

Disclosure: Ratnakar Pingili, Cecile Kerloeguen, Ayan Das Gupta, Valentine Jehl, Dieter A. Haering, Krishnan Ramanathan, Martin Merschhemke: Novartis (salary). Amit Bar-Or: Atara Biotherapeutics, Biogen, Celgene, EMD Serono, Genentech/Roche, GlaxoSmithKline, Mapi, MedImmune, Merck, Novartis, Receptos, Sanofi Genzyme (consulting fee, grant support, speakers' bureau). Jeffrey A. Cohen: Adamas, Convelo, Mylan, Population Council (consulting fee); Multiple Sclerosis Journal (co-editor). Patricia K. Coyle: Accordant, Alexion, Bayer, Biogen, Celgene, Sanofi Genzyme, GlaxoSmithKline, Mylan, Serono, TG Therapeutics (consulting fee); Actelion, Alkermes, Corrona LLD, MedDay, National Institute of Neurological Diseases and Stroke, PCORI (contracted research); Genentech/Roche, Novartis (consulting fee, contracted research). Anne H. Cross: Academic CME (CE provider) (speaking and preparation of slides for CME talk); Biogen, Celgene, TG Therapeutics (consulting fee); EMD Serono (consulting fee, contracted research); Genentech/Roche (consulting fee, contracted research, speakers' bureau); Novartis (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent, scientific advisory board for ASCLEPIOS I and II); Projects in Knowledge (CE provider) (preparation of educational manuscripts; activities); Rockpointe (Potomac Center for Medical Education) (CE provider) (speakers' bureau). Stephen L. Hauser: Alector (scientific advisory board member [stock options received]); Annexon, Bionure, Molecular Stethoscope, Symbiotix (scientific advisory board member [stock options previously received]); F. Hoffmann-La Roche (travel support/reimbursement and

writing support for anti-CD20 meetings and presentations); Neurona (board of trustees member [stock options previously received]); Novartis (travel support/reimbursement for anti-CD20 meetings and presentations). Ludwig Kappos: Actelion, Minoryx, Receptos, Santhera (consulting fee, steering committee; advisory board); Allergan, Allmirall, CSL Behring, Pfizer (support for educational activities); Bayer HealthCare (consulting fee, speakers' bureau, steering committee; advisory board; support for educational activities; license fees for Neurostatus products; grants); Biogen, Merck (consulting fee, steering committee; advisory board; support for educational activities; license fees for Neurostatus products; grants); Celgene, Genzyme (consulting fee, steering committee; advisory board; support for educational activities); Desitin (support of educational activities); Novartis, Roche (consulting fee, contracted research, speakers' bureau, steering committee; advisory board; support for educational activities; license fees for Neurostatus products; grants); Roche Research Foundation (grants); Sanofi (consulting fee, speakers' bureau, steering committee; advisory board; support for educational activities); Teva (consulting fee, steering committee; advisory board; support for educational activities; license fees for Neurostatus products).

Keywords: Disease-modifying treatments in MS, Immunology and MS, Injection-related reactions

(DXT62) Real-World Treatment Patterns in Patients with Multiple Sclerosis Using Disease-Modifying Therapies

Robert J. Fox,¹ Rina Mehta,² Tim Pham,² Julie Park,³ Kathleen Wilson,³ Machaon Bonafede³

¹Cleveland Clinic, Cleveland, OH; ²Bristol-Myers Squibb, Summit, NJ; ³IBM Watson Health, Cambridge, MA

Background: Numerous disease-modifying therapies (DMTs) have been approved for the treatment of multiple sclerosis (MS) in the past decade, and few studies have assessed patterns of use among all approved DMTs. **Objectives:** This study characterized patterns of DMT use in patients with newly diagnosed MS. **Methods:** Adults with newly diagnosed MS were identified from January 2007 to October 2017 using the IBM MarketScan Commercial and Medicare databases. Patients had at least 12 months of continuous enrollment prior to their initial MS diagnosis and 2 years of follow-up. Up to 3 DMT lines of therapy (LOTs) were reported during a follow-up of 2 to 10.5 years. Discontinuation or switch of therapy was assessed. **Results:** Of 29,647 patients with at least 2 years of follow-up from MS diagnosis, 14,627 were treated with DMTs. Of these, 49% had 1 DMT LOT during follow-up, 25% had 2 LOTs, and 27% had 3 LOTs. Injectable (subcutaneous or intramuscular) DMTs, comprising glatiramer acetate (GA), interferon beta-1a (IFNβ-1a) (subcutaneous and intramuscular), interferon beta-1b (IFNβ-1b), and peginterferon beta-1a (pegIFNβ-1a), were used by 87% of patients as first LOT, 68% as second LOT, and 67% as third LOT. Oral DMTs, including dimethyl fumarate, fingolimod, and teriflunomide, were used by 11% of patients as first LOT, but increased to 30% of second and 32% of third LOTs. Natalizumab, the only infusion DMT in this analysis, was used by less than 3% as first, second, and third LOTs. The most common pattern after ending the first LOT was discontinuation from all DMTs (51%), while 17% switched DMTs and 26% restarted the same DMT treatment later. Long-term discontinuation increased to 56% of patients with a second LOT and 58% with a third LOT. Patients on GA and IFNβ-1b had the highest rates of discontinuation (52% to 61% for each drug and each LOT) and restart (18% to 31% for each drug and each LOT). Patients on pegIFNβ-1a had the highest rates of switching to another DMT, ranging from 20% to 29%. Those taking dimethyl fumarate and fingolimod at first LOT were least likely to switch treatments throughout all LOTs. **Conclusions:** Over 2 to 10.5 years of follow-up, most patients with MS discontinued or switched from their first-line DMT, regardless of DMT type. Injectable DMTs were the most commonly used DMTs over this study period, although the rate of oral DMT use increased in later lines of therapy.

Supported by: None

Disclosure: Robert J. Fox: Actelion, Biogen, Celgene, EMD Serono, Genentech, Immunic, Novartis, Sanofi, Teva, TG Therapeutics (consulting fee); Actelion, Biogen, Immunic, Novartis (advisory committee); Biogen, Novartis (clinical trial contract and research grant funding). Rina Mehta, Tim Pham: Bristol-Myers Squibb (employment). Julie Park, Kathleen Wilson, Machaon Bonafede: IBM Watson Health (employment).

Keywords: Disease-modifying treatments in MS, Treatment patterns

(DXT63) Associations Between Treatment Satisfaction, Medication Beliefs, and Adherence to Disease-Modifying Therapies in Patients with Multiple Sclerosis Among Adult Saudis: A Tertiary Care Center Experience

Rola F. Alarief

King Fahad Medical City, Riyadh, Saudi Arabia

Background: Multiple sclerosis (MS) is considered one of the most common neuroimmune diseases that leads to major disabilities in an affected patient with a significant burden and consequences to patients and their families. Even though there is no available cure for MS, the past 2 decades witnessed a promising future for MS treatment drugs, specifically disease-modifying therapies (DMTs), to reduce MS relapse and delay disability. Adherence to DMTs has a significant impact on treatment outcomes and is considered a critical factor in therapeutic success. Accordingly, the need to examine this issue in Saudi Arabia stands. **Objectives:** To identify the factors associated with adherence to DMT medications among patients with MS in Saudi Arabia. To evaluate the relationship between treatment satisfaction, medication beliefs, and DMT adherence and other factors. **Methods:** A survey was conducted in 2019 in neurology clinics in King Fahad Medical City (KFMC) in Riyadh. Patients were sampled from the KFMC's Data Base with population size of 387 patients. The survey measured self-reported DMT adherence (doses taken divided by doses prescribed during previous 2-week period—adherence ≥ 0.80), DMT satisfaction using the Treatment Satisfaction Questionnaire for Medication version II, medication beliefs using the Beliefs About Medicines Questionnaire, and demographic and clinical covariates. Relationships between variables were examined using multivariate logistic regression. **Results:** Final analyses included 239 usable surveys. Mean \pm SD participant age was 35.07 ± 9.7 years. Most respondents were female (74.9%), taking an injectable DMT (49%), and adherent to DMT (64.4%). Significant predictors of DMT adherence were DMT experience (naive vs experienced [odds ratio (OR), 3.722; 95% CI, 1.487-9.316; $P = .005$]), DMT route (oral vs injectable; OR, 0.974; 95% CI, 0.952-0.995; $P = .017$), and global satisfaction (OR, 0.950; 95% CI, 0.926-0.975; $P < .001$). **Conclusions:** In patients with MS sampled from KFMC's Data Base, medication beliefs was not significantly associated with DMT adherence while global satisfaction was significantly associated with DMT adherence. Based on significant predictors, patients taking injectable DMTs and patients with previous experience with another DMT(s) are at higher risk for nonadherence. Future research is warranted to assess relationships between variables in more diverse MS populations.

Supported by: None

Disclosure: Rola F. Alarieb: King Fahad Medical City (ownership interest).

Keywords: Adherence to therapy, Disease-modifying treatments in MS

(DXT64) Evaluation of Rituximab Regimens and Outcomes in Neuromyelitis Optica Patients from a Single Academic Medical Center: A Retrospective Chart Review

Ryan Fuller,¹ Stephanie Witek,² Lauren Long¹

¹Neurology and ²Pharmacy, Hospital of the University of Pennsylvania, Philadelphia, PA

Background: Neuromyelitis optica (NMO) is a rare neurologic disease often mistaken for multiple sclerosis that has limited treatment options and significant morbidity and mortality. Rituximab has been used for several years with demonstrated efficacy, but the optimal use of this agent in this patient population remains unknown. This has led to varying practices even within a single center. The current study aims to identify current practices and associated outcomes at an academic medical center without a standardized treatment protocol. **Objectives:** The primary end point is time to relapse after first rituximab dose. Secondary end points include cumulative relapses, proportion of patients relapsing, and adverse event rate. These data will be used to work towards a standard protocol amongst NMO providers at the associated practice and will be hypothesis-generating for further research on the optimal use and place in therapy of rituximab for NMO/NMO spectrum disorder (NMOSD) in light of new and emerging treatment options. **Methods:** A retrospective chart review was conducted on approximately 60 patients at least 18 years old with NMO or NMOSD who received at least 2 doses of rituximab and were seen by a Penn Medicine neurology provider between February 1, 2013, and February 1, 2019. Patients were excluded due to a concurrent diagnosis of sarcoidosis, active cancer, or chronic infection (hepatitis B or C, TB, or HIV). Data collected include gender, age, weight/body mass index, race, diagnosis, NMO episode/relapse history, aquaporin-4 antibody serostatus, prior treatment, dates and doses of rituximab, CD19+ B-cell level and laboratory draw date, and adverse effects. **Results:** Data will be analyzed with descriptive statistics. **Conclusions:** It is anticipated that we will identify a superior dosing strategy that will edify the practice of rituximab dosing in patients with NMO at our institution.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Chart review, NMO, NMOSD

(DXT65) Longitudinal Disability Follow-up in Patients with 6-Month Confirmed Disability Improvement or Worsening in the CARE-MS and Extension Studies

Samuel F. Hunter,¹ Darren P. Baker,² Mark Ozog,² Elizabeth M. Poole,² Luke Chung,² Patrick Vermersch,³ Rany A. Aburashed,⁴ Raed Alroughani,⁵ Dominique Dive,⁶ Ho Jin Kim,⁷ Jan Lycke,⁸ Richard A.L. Macdonell,⁹ Carlo Pozzilli,¹⁰ Heinz Wiendl,¹¹ on behalf of the CARE-MS I, CARE-MS II, CAMMS03409, and TOPAZ investigators

¹Advanced Neurosciences Institute, Franklin, TN; ²Sanofi, Cambridge, MA; ³University of Lille, INSERM U995, CHU Lille, Lille, France; ⁴Institute for Neurosciences and Multiple Sclerosis, Owosso, MI; ⁵Amiri Hospital, Sharq, Kuwait; ⁶University Hospital Centre of Liège, Liège, Belgium; ⁷Research Institute and Hospital, National Cancer Center, Goyang, Korea, Republic of (South); ⁸Institute of Neuroscience and Physiology, University of Gothenburg, Gothenburg, Sweden; ⁹Austin Health and Florey Institute of Neuroscience and Mental Health, Melbourne, VIC, Australia; ¹⁰Sapienza University, Rome, Italy; ¹¹Department of Neurology, University of Münster, Münster, Germany

Background: In the 2-year CARE-MS trials (trial registration: NCT00530348; NCT00548405), alemtuzumab (12 mg/day; baseline: 5 days; 12 months later: 3 days) significantly improved clinical/magnetic resonance imaging outcomes vs subcutaneous interferon beta-1a in patients with relapsing-remitting multiple sclerosis (RRMS). Efficacy in alemtuzumab-treated patients was maintained through year 9 in 2 consecutive extension studies (NCT00930553; NCT02255656 [TOPAZ]). **Objectives:** To evaluate the status of disability over 9 years in pooled CARE-MS patients who had either 6-month confirmed disability improvement (CDI) or 6-month confirmed disability worsening (CDW) by years 2 or 9. **Methods:** Alemtuzumab-treated CARE-MS patients with baseline Expanded Disability Status Scale (EDSS) score ≥ 2 were stratified into 3 subgroups: with CDI, with CDW, or with stable disability. CDI and CDW were defined as ≥ 1 -point decrease and increase, respectively, in EDSS score from core study baseline confirmed over 6 months. Improved and stable EDSS scores each year were defined as ≥ 1 -point decrease and ≤ 0.5 -point change in either direction, respectively, from core study baseline in available patients. **Results:** 511/811 (63%) alemtuzumab-treated patients had baseline EDSS score ≥ 2 . Of these, 222 (43%) patients had 365 unique CDI events and 172 (34%) patients had 217 CDW events at any time during the 9-year study; 31 (6%) had both CDI and CDW. Few patients ($n = 12$) had a CDW event after CDI. Of patients with CDI at any time over 9 years, mean EDSS score change was -0.58 at year 9 vs core study baseline, and 51% had lower EDSS scores at year 9. Similar EDSS outcomes were observed at year 9 in the subset of patients who achieved CDI within the first 2 years of the study. However, patients with CDW any time over 9 years had worsened disability at year 9, with a $+1.71$ mean change in EDSS score from core study baseline; patients with CDW in the first 2 years of the study had a $+2.27$ EDSS score change by year 9. EDSS scores remained stable at 9 years (mean change, -0.10) in the 148 (29%) patients who had neither CDI nor CDW. Compared with previous years, no new safety signals were identified in year 9 in CARE-MS extension study patients. **Conclusions:** Achievement of CDI at any point in the CARE-MS studies was associated with improved disability at year 9 vs baseline. However, those with CDW had increased disability over 9 years regardless of when worsening occurred. These findings suggest that both CDI and CDW are meaningful end points for predicting long-term disability outcomes in patients with RRMS.

Supported by: Sanofi, Bayer HealthCare Pharmaceuticals

Disclosure: Samuel F. Hunter: AbbVie, Acorda, Avanir, Bayer, Biogen, Celgene, Genzyme, Mallinckrodt, Novartis, Osmotica, Roche, Teva (consulting agreements, speaker honoraria, grant/research support). Darren P. Baker, Mark Ozog, Elizabeth M. Poole, Luke Chung: Sanofi (salary). Patrick Vermersch: Almirall, Biogen, Celgene, Merck, Novartis, Roche, Sanofi, Servier, Teva (consulting and/or speaking fees and research support). Rany A. Aburashed: Bayer, Biogen, Genentech, Sanofi, Teva (consulting and/or speaker honoraria and scientific advisory boards); Novartis (consulting and/or speaker honoraria and scientific advisory boards, research grants). Raed Alroughani: Bayer, Biogen, Biologix, Genpharm, GlaxoSmithKline, Lundbeck, Merck Serono, Novartis, Sanofi (speaker honoraria, scientific advisory boards, and research grants). Dominique Dive: Bayer, Biogen, Merck Serono, Novartis, Sanofi, Teva (institutional honoraria for advisory board participation and travel grants). Ho Jin Kim: Alexion, Celltrion, Eisai, HanAll BioPharma, Merck Serono, Novartis, Sanofi, Teva-Handok, Viela Bio (consulting and speaking fees); Journal of Clinical Neurology, Multiple Sclerosis Journal (coeditor/associate editor); MedImmune/Viela Bio (steering committee member); National Research Foundation of Korea (research support). Jan Lycke: Acta Neurologica Scandinavica (editorial board); Almirall (scientific advisory boards); Biogen, Novartis, Teva (scientific advisory boards, travel support and/or lecture honoraria, unconditional research grants); Sanofi (scientific advisory boards, travel support and/or lecture honoraria). Richard A.L. Macdonell: Actelion, Biogen, Merck, Novartis, Sanofi, Teva (travel grants). Carlo Pozzilli: Actelion, Biogen,

Merck, Novartis, Sanofi, Teva (consulting and/or speaking fees, research, and travel grants). *Heinz Wiendl*: Bayer, Behring, Biogen, EMD Serono, Fresenius Medical Care, Merck Serono, Roche, Sanofi, Teva (consulting and/or speaking fees); Huber-Verlag (license fee payments); Neotope Biosciences, PML Consortium (grant/research support); Novartis (consulting and/or speaking fees, grant/research support)

Keywords: CNS repair, Disease-modifying treatments in MS

(DXT66) Clinical Benefits of Eculizumab Monotherapy in Neuromyelitis Optica Spectrum Disorder: Findings from the Phase 3 PREVENT Study

Sean J. Pittock,¹ Achim Berthele,² Ho Jin Kim,³ Kazuo Fujihara,^{4,5,6} Michael Levy,^{7,8} Ichiro Nakashima,⁹ Celia Oreja-Guevara,¹⁰ Jacqueline Palace,¹¹ Murat Terzi,¹² Natalia Totolyan,¹³ Shanthi Viswanathan,¹⁴ Kai-Chen Wang,^{15,16} Amy Pace,¹⁷ Marcus Yountz,¹⁷ Larisa Miller,¹⁷ Guido Sabatella,¹⁷ Roisin Armstrong,¹⁷ Dean Wingerchuk¹⁸

¹Mayo Clinic, Rochester, MN; ²Technical University of Munich, Munich, Germany; ³National Cancer Center, Goyang, Korea, Republic of (South); ⁴Southern TOHOKU Research Institute for Neuroscience (STRINS), Koriyama, Japan; ⁵Tohoku University, Sendai, Japan; ⁶Fukushima Medical University, Fukushima City, Japan; ⁷Massachusetts General Hospital and Harvard Medical School, Boston, MA; ⁸Johns Hopkins University, Baltimore, MD; ⁹Tohoku Medical and Pharmaceutical University, Sendai, Japan; ¹⁰Hospital Universitario Clinico San Carlos, Madrid, Spain; ¹¹John Radcliffe Hospital, Oxford, United Kingdom; ¹²Ondokuz Mayıs University, Samsun, Turkey; ¹³First Pavlov State Medical University of St. Petersburg, St. Petersburg, Russian Federation; ¹⁴Kuala Lumpur Hospital, Kuala Lumpur, Malaysia; ¹⁵National Yang Ming University, Taipei, Taiwan; ¹⁶Cheng-Hsin General Hospital, Taipei, Taiwan; ¹⁷Alexion Pharmaceuticals, Boston, MA; ¹⁸Mayo Clinic, Scottsdale, AZ

Background: In the PREVENT study in patients with aquaporin-4 immunoglobulin G-positive (AQP4-IgG+) neuromyelitis optica spectrum disorder (NMOSD), eculizumab was associated with a significant reduction in relapse risk vs placebo and was well tolerated. The time course of relapses in prespecified subgroups suggested a treatment effect consistent with that in the overall population, regardless of use of permitted concomitant immunosuppressive therapy (IST) (rituximab and mitoxantrone were excluded). **Objectives:** To examine the efficacy of eculizumab relative to placebo in patients with AQP4-IgG+ NMOSD who did not receive concomitant IST during the randomized, double-blind, placebo-controlled, phase 3 PREVENT trial (trial registration: NCT01892345). **Methods:** Adults with AQP4-IgG+ NMOSD received eculizumab (maintenance dose, 1200 mg/2 weeks, n = 96) or placebo (n = 47) with/without concomitant IST. A post hoc descriptive analysis of clinical outcomes was performed using data from patients who did not receive concomitant IST during PREVENT (ie eculizumab monotherapy or placebo without concomitant IST subgroup). Clinical outcomes comprised relapses, hospitalizations and acute treatment for relapses, and worsening of Expanded Disability Status Scale (EDSS) or Hauser Ambulation Index (HAI) scores. **Results:** Of 34 patients in the no-IST subgroup, 10 had never received IST for NMOSD and 14 had previously received rituximab. Adjudicated relapses occurred in 0/21 patients receiving eculizumab monotherapy and 7/13 (53.8%) receiving placebo ($P < .0001$; post hoc analysis). In the placebo group, 6/13 patients (46.2%) were hospitalized for adjudicated relapses, 3 (23.1%) received plasma exchange, 4 (30.8%) were treated with acute intravenous methylprednisolone, and 1 (7.7%) received high-dose oral corticosteroids. EDSS scores worsened in 1/21 patients (4.8%) receiving eculizumab monotherapy and 5/13 (38.5%) patients receiving placebo. HAI scores worsened in 1/21 patients (4.8%) receiving eculizumab monotherapy and 4/13 (30.8%) patients receiving placebo. **Conclusions:** These data further characterize the substantial efficacy of eculizumab monotherapy in reducing relapse risk in patients with AQP4-IgG+ NMOSD. Patients receiving eculizumab monotherapy were spared relapse-associated hospitalizations and acute treatments, and most (95%) did not experience disability worsening. Long-term results from PREVENT's open-label extension will be presented.

Supported by: None

Disclosure: *Sean J. Pittock*: Alexion Pharmaceuticals (all compensation paid directly to Mayo Clinic, consulting fee, contracted research); *Astellas* (all compensation paid directly to Mayo Clinic, consulting fee); *Autoimmune Encephalitis Alliance* (all compensation paid directly to Mayo Clinic, contracted research); *Grifols* (all compensation paid directly to Mayo Clinic, contracted research); *MedImmune* (all compensation paid directly to Mayo Clinic, consulting fee, contracted research); *UCB* (consulting fee, personal compensation for attending UCB advisory board meeting in Stockholm, Sweden, on Sept 10, 2019). *Achim Berthele*: Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent, speakers' bureau). *Ho Jin Kim*: Alexion Pharmaceuticals, Celltrion, Eisai, HanAll BioPharma, Merck Serono, Novartis, Sanofi Genzyme, Teva-Handok, Viela Bio (consulting fee); *Journal of Clinical Neurology* (associate editor); *MedImmune/Viela Bio* (steering committee); *Multiple Sclerosis Journal* (co-editor). *Kazuo Fujihara*:

Alexion Pharmaceuticals, Biogen, Chugai, Mitsubishi-Tanabi, Novartis (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent); *Asahi Medical, Bayer, Eisai, Roche, Teijin* (fees for non-CME/CE services received directly from commercial interest or its agent). *Michael Levy*: Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent); *Genentech, Quest Diagnostics, Viela Bio* (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent). *Ichiro Nakashima*: Alexion (consulting fee). *Celia Oreja-Guevara*: Alexion Pharmaceuticals, Merck, Novartis, Roche, Sanofi (consulting fee, contracted research). *Jacqueline Palace*: Biogen, Chugai (contracted research); *LEK, Viela Bio, Guidepoint* (consulting fee); *Merck Serono* (meeting/lecture/workshop participation); *Novartis, Roche, Argencx* (speakers' bureau); *UCB, Viela Bio, Roche* (conference/lecture participation). *Murat Terzi*, *Shanthi Viswanathan*, *Kai-Chen Wang*: Nothing to disclose. *Natalia Totolyan*: Alexion, Janssen, Novartis, Roche, Sanofi, Receptos Inc, Biocad (Russia) (contracted research); *Merck* (consulting fee); *Roche, Sanofi* (fees for lectures). *Amy Pace*, *Marcus Yountz*, *Guido Sabatella*, *Roisin Armstrong*: Alexion Pharmaceuticals (ownership interest, salary). *Larisa Miller*: Alexion Pharmaceuticals (RSU, salary). *Dean Wingerchuk*: *MedImmune, Novartis, Biogen, Celgene, Genentech, TG Therapeutics, Arcus Medica, Reistone* (consulting fee); *Terumo BCT, Alexion Pharmaceuticals* (fees for non-CME/CE services received directly from commercial interest or its agent).

Keywords: Eculizumab monotherapy in NMOSD

(DXT67) Cognitive Functions over the Course of 5 Years in Multiple Sclerosis Patients Treated with Disease-Modifying Therapies

Serkan Ozakbas,¹ Cavid Baba,² Taha Aslan,² Ozge Sagici,² Yasemin Simsek,² Pinar Yigit²

¹Department of Neurology, Dokuz Eylul University, Izmir, Turkey; ²Dokuz Eylul University, Izmir, Turkey

Background: Cognitive decline is common in multiple sclerosis (MS). Disease-modifying therapies (DMTs) are applied to delay or prevent disease progression in MS. While this has mostly been proven for physical status, comprehensive data on cognitive functions are not yet available. **Objectives:** We aimed to present 5 years of cognition data of patients treated with DMTs. **Methods:** In total, 756 patients with MS who were treated with interferon beta (1b or 1a, subcutaneous) (n = 342), glatiramer acetate (GA) (n = 188), or fingolimod (n = 226). Mean age was higher in the fingolimod group ($P < .05$). Physical disability was assessed with Expanded Disability Status Scale (EDSS), and cognitive status was assessed with Brief International Cognitive Assessment for MS (BICAMS) which included the Symbol Digit Modalities Test (SDMT), the California Verbal Learning Test-II (CVLT-II), and the Brief Visuospatial Memory Test-Revised (BVMTR). Assessments were performed at baseline and yearly until the end of fifth year of treatment. **Results:** 85% of patients treated with fingolimod were still having their medication at the end of fifth year (79% for GA and 78.9% for interferon beta, $P < .05$). Most of the patients treated with DMTs remained stable over the course of 5 years (fingolimod: 70.1%, interferons: 71.9%, GA: 75%, $P > .05$). Cognition improved in some patients (fingolimod: 11.2%, interferons: 9.6%, GA: 8%, $P > .05$). More than 80% of patients remained stable or improved. The most significant improvement was observed in SDMT, and it was significantly higher than CVLT-II and BVMTR (30.7%, 18.6%, and 17%, respectively, $P = .02$). **Conclusions:** In conclusion, cognitive functions remain stable under DMTs over 5 years. This condition has not shown a relationship with the type of medication. Furthermore, SDMT seems to be the best predictor for cognitive change over time.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Disease-modifying treatments in MS, Psychological issues and MS

(DXT69) One-Year Interim Analysis of Real-World Patient-Reported Outcomes in Relapsing-Remitting Multiple Sclerosis Patients Transitioning to Alemtuzumab (PRO-ACT Study)

Sibyl Wray,¹ Francois Jacques,² Jacqueline Nicholas,³ Susan Mozzicato,⁴ Benjamin Guikema,⁴ Elizabeth M. Poole,⁴ Tamara A. Miller⁵

¹Hope Neurology, Knoxville, TN; ²Clinique Neuro-Outaouais, Gatineau, QC, Canada; ³OhioHealth Multiple Sclerosis Center, Columbus, OH; ⁴Sanofi, Cambridge, MA; ⁵Advanced Neurology of Colorado, Fort Collins, CO

Background: Clinical trials of alemtuzumab have demonstrated its 9-year efficacy and safety, but real-world data are limited. PRO-ACT assesses patient-reported outcomes (PROs), safety, and treatment sequencing in adults with relapsing-remitting multiple sclerosis (RRMS) transitioning

from prior disease-modifying therapy to alemtuzumab in the United States and Canada. **Objectives:** To report the 1-year interim results of the real-world PRO-ACT study. **Methods:** PRO-ACT is an ongoing 24-month, prospective, multicenter, noninterventional, single-arm, observational study. The primary end point evaluates change in overall satisfaction on the Treatment Satisfaction Questionnaire for Medication v1.4 (TSQM; scale 0-100; higher scores indicate greater satisfaction), after transitioning to alemtuzumab. Other PROs evaluated (lower scores indicate improved outcomes) are the MS Impact Scale-29 (MSIS-29; scale 0-100), Modified Fatigue Impact Scale-5 (MFIS-5; scale 0-20), Patient-Determined Disease Steps (PDDS; scale 0-8), and Health-Related Productivity Questionnaire (HRPQ)-MS v2. **Results:** As of September 2019, enrollment was complete (N = 200 patients) and PRO data were evaluable in 170 patients. Patients transitioned from natalizumab (37%), dimethyl fumarate (14%), fingolimod (13%), teriflunomide (12%), or other therapies (24%) to alemtuzumab. Mean TSQM scores improved from baseline to year (Y) 1 for overall satisfaction (50.3 vs 66.5; $P < .0001$) and effectiveness (49.3 vs 60.7; $P < .0001$) domains; scores were unchanged for side effects (77.6 vs 76.5) and convenience (70.3 vs 70.7). Mean scores for other PROs showed improvement at Y1 vs baseline: MSIS-29 physical impact scale (52.4 vs 47.8; $P < .001$), MSIS-29 psychological impact scale (53.4 vs 47.9; $P < .001$), and MFIS-5 (12.8 vs 11.7; $P < .001$). Scores remained stable on the PDDS (3.1 vs 3.2). Mean hours of weekly employment productivity lost decreased from 11.4 at baseline to 7.4 at Y1 ($P < .05$). Incidence of adverse events was 92% and serious adverse events was 11%. One death occurred (suicide). **Conclusions:** PROs improved during the first year of alemtuzumab treatment after transitioning from another therapy. Alemtuzumab safety in Y1 was consistent with the pivotal studies.

Supported by: Sanofi

Disclosure: *Sibyl Wray:* Alkermes, Biogen, Celgene, Genentech, Novartis, Sanofi, TG Therapeutics (consulting, principal investigator, and/or speaking fees). *Francois Jacques:* Biogen, Merck Serono, Novartis, Roche, Sanofi (honoraria for presentations, advisory board participation, research funding, and for an infusion clinic). *Jacqueline Nicholas:* Biogen, Celgene, EMD Serono, Genentech, Multiple Sclerosis Association of America, National MS Society, Novartis, PCORI, Sanofi (consulting, research, and/or speaking). *Susan Mozzicato, Benjamin Guikema, Elizabeth M Poole:* Sanofi (salary). *Tamara A. Miller:* Acorda, Amgen, Mallinckrodt, Reven (speaking and/or consulting fees); Adamas, Elan, EMD Serono, Ipsen, Ono, Sun Pharma (research support); Allergan, Biogen, Genentech, Novartis, Sanofi, Teva (research support, speaking and/or consulting fees).

Keywords: Disease-modifying treatments in MS

(DXT70) Clinical Characteristics and Outcomes of Peginterferon Beta-1a Treatment by Age: A Subgroup Analysis of the Plegridy Observational Program

Sibyl Wray,¹ Marco Salvetti,² Gereon Nelles,³ Arman Altincatal,⁴ Megan C. Vignos,⁴ Achint Kumar,⁴ Maria L. Naylor⁴

¹Hope Neurology, Knoxville, TN; ²Sapienza University, S. Andrea Hospital, Rome, Italy; ³Neurology, NeuroMed Campus Hohenlind, Cologne, Germany; ⁴Biogen, Cambridge, MA

Background: Peginterferon beta-1a is approved to treat relapsing forms of multiple sclerosis (RMS) based on results from the ADVANCE trial, which had a mean patient age of 36 years. US claims data show a mean age in the peginterferon beta-1a patient postmarketing population of 45-50 years, yet data on disease-modifying therapies for patients aged ≥ 50 with RMS are limited. The 5-year, phase 4 Plegridy Observational Program (POP) study explores the real-world safety and effectiveness of peginterferon beta-1a in patients with RMS, a third of whom are aged ≥ 50 . **Objectives:** Report on baseline characteristics, safety outcomes, and clinical effectiveness of peginterferon beta-1a in patients ≥ 50 and < 50 years of age enrolled in POP. **Methods:** The safety analysis population included 1126 patients; 375 (33%) were ≥ 50 and 751 (67%) were < 50 years of age at baseline. The effectiveness analysis population included 1125 (≥ 50 , 375; < 50 , 750) patients. Data reflect the third interim data cut as of September 2018. Data from the fourth interim data cut (September 2019) will be presented. **Results:** The mean (SD) ages of ≥ 50 and < 50 patients were 58.0 (6.2) and 36.8 (8.2) years at baseline, respectively, and 45.2 (9.4) and 31.7 (7.6) years at multiple sclerosis (MS) diagnosis, respectively. The percentage of female patients was similar in the ≥ 50 (74.7%) and < 50 (77.1%) subgroups. At baseline, the mean time since MS symptom onset was higher for the ≥ 50 group than the < 50 group (17.1 vs 7.4 years). Patients ≥ 50 had fewer relapses in the year prior to baseline (mean: 0.3 vs 0.6) and in the 3 years prior to baseline (mean: 0.6 vs 1.0) than patients < 50 . At baseline, the mean Expanded Disability Status Scale score was higher in patients ≥ 50 (2.69) vs patients < 50 (1.45). The incidence of treatment-emergent adverse events (AEs) in POP was similar in the ≥ 50 (63.2%) and < 50 (62.7%)

subgroups. A higher incidence of serious AEs (SAEs) was reported in patients ≥ 50 (9.6%) vs < 50 (5.1%). The annualized relapse rate (ARR) was 0.08 in patients ≥ 50 and 0.15 in those < 50 . The percent of relapse-free patients over 3 years was 88.3% in patients ≥ 50 and 78.3% in patients < 50 . **Conclusions:** In both subgroups, the incidence of AEs was similar; the incidence of SAEs was higher in patients aged ≥ 50 . ARR was low and the percentage of relapse-free patients was high in both subgroups, indicating that peginterferon beta-1a has the potential to provide treatment benefits to patients with RMS, including those aged ≥ 50 .

Supported by: Biogen

Disclosure: *Sibyl Wray:* Bayer, Biogen, Celgene, EMD Serono, Genentech/Roche, Sanofi Genzyme, Novartis, Receptos, TG Therapeutics (paid consultant, speaker, and/or contract researcher). *Marco Salvetti:* Biogen, Merck, Novartis, Roche, Sanofi, Teva (grant support and/or speaker honoraria). *Gereon Nelles:* Bayer, Biogen, Celgene, Merck, Novartis, Roche (speaker honoraria). *Arman Altincatal, Megan C. Vignos, Achint Kumar, Maria L. Naylor:* Biogen (employee, may hold stock and/or stock options).

Keywords: Age, Disease-modifying treatments in MS

(DXT71) Efficacy and Safety of Teriflunomide in Patients with Relapsing-Remitting Multiple Sclerosis of Varying Disease Duration: Analysis of Pooled Clinical Trials

Stanley Cohan,¹ Albert Saiz,² David Rog,³ Mauro Zaffaroni,⁴ Sandra Vukusic,⁵ Steve Vucic,⁶ Jiwon Oh,⁷ Klaus Tiel-Wilck,⁸ Anat Achiron,⁹ Darren P. Baker,¹⁰ Janneke Wingerden,¹¹ Elizabeth M. Poole,¹⁰ Bhupendra O. Khatri¹²

¹Providence Brain and Spine Institute, Portland, OR; ²Department of Neurology, Hospital Clinic and Institut d'Investigacions Biomèdiques August Pi i Sunyer, University of Barcelona, Barcelona, Spain; ³Manchester Centre for Clinical Neurosciences, Salford, United Kingdom; ⁴MS Centre, Gallarate Hospital, Gallarate, Italy; ⁵University Hospital of Lyon, Lyon, France; ⁶Department of Medicine, University of Sydney, Sydney, NSW, Australia; ⁷University of Toronto, Toronto, ON, Canada; ⁸Neurologisches Facharztzentrum Berlin, Berlin, Germany; ⁹Sheba Medical Center, Ramat Gan, Israel; ¹⁰Sanofi, Cambridge, MA; ¹¹Sanofi, Amsterdam, Netherlands; ¹²Center for Neurological Disorders at Wheaton Franciscan Healthcare, Milwaukee, WI

Background: Teriflunomide is a once-daily oral immunomodulator approved for the treatment of relapsing forms of multiple sclerosis (MS) or relapsing-remitting MS (RRMS), depending on the local label. **Objectives:** To assess the long-term efficacy and safety of teriflunomide in patients with RRMS stratified by disease duration. **Methods:** This was a pooled efficacy and safety analysis using data from the phase 2 study (trial registration: NCT01487096) and the phase 3 TEMSO (NCT00134563, NCT00803049), TOWER (NCT00751881), and TENERE (NCT00883337) core and extension studies. Patients receiving placebo or teriflunomide 14 mg were stratified by disease duration at baseline (≤ 1 year, > 1 to 5 years, > 5 to 10 years, and > 10 years). Study end points included annualized relapse rate (ARR), Expanded Disability Status Scale (EDSS) score, 6-month confirmed disability worsening (CDW), and safety. **Results:** In the core period, ARR was lower in patients treated with teriflunomide 14 mg compared with placebo across disease duration subgroups: ≤ 1 year (0.33 [n = 272] vs 0.56 [n = 251], $P = .0013$), > 1 to 5 years (0.46 [n = 278] vs 0.70 [n = 268], $P = .0011$), > 5 to 10 years (0.39 [n = 191] vs 0.52 [n = 164], $P = .0571$), and > 10 years (0.33 [n = 154] vs 0.58 [n = 129], $P = .0005$). In the core+extension period (up to year 13), ARRs in teriflunomide-treated patients were similar regardless of disease duration: ≤ 1 year (0.19; n = 276), > 1 to 5 years (0.22; n = 699), > 5 to 10 years (0.26; n = 393), and > 10 years (0.25; n = 325). At year 13, 6-month CDW rates for each group were 48.3% (≤ 1 year), 37.1% (> 1 to 5 years), 52.6% (> 5 to 10 years), and 36.8% (> 10 years). From core study baseline to year 10 (the last time point at which all groups had at least 10 patients), EDSS scores were stable across teriflunomide-treated patients of different disease durations: ≤ 1 year, +0.27; > 1 to 5 years, +1.11; > 5 to 10 years, +0.05; and > 10 years, +0.7. Overall incidences of adverse events through year 13 were 93.1% (≤ 1 year), 87.2% (> 1 to 5 years), 88.0% (> 5 to 10 years), and 88.7% (> 10 years); incidences of serious adverse events during this period were 21.2% (≤ 1 year), 19.1% (> 1 to 5 years), 15.5% (> 5 to 10 years), and 18.0% (> 10 years). **Conclusions:** Teriflunomide 14 mg reduced relapses across all patients regardless of disease duration vs placebo in the core studies. Over 13 years, ARR remained low and EDSS score increased minimally. Safety outcomes from baseline to year 13 were consistent across disease duration subgroups.

Supported by: Sanofi

Disclosure: *Stanley Cohan:* AbbVie (research support); Biogen, Novartis, Sanofi (advisory boards, research steering committees, research support, and speaking honoraria, employment, consulting, scientific advisory board, speaking, or other activities); EMD Serono, Genentech, Pear Therapeutics (employment, consulting,

scientific advisory board, speaking, or other activities); Roche Genentech (advisory boards, research steering committees, research support, speaking honoraria). **Albert Saiz:** Bayer, Biogen, Merck-Serono, Novartis, Roche, Sanofi, Teva (remuneration for providing consulting services and lectures). **David Rog:** Actelion, Genzyme, GW Pharma, Mitsubishi, TG Therapeutics (research support); Bayer, MedDay, Roche, Sanofi (consulting fee); Biogen, Merck Serono, Novartis, Teva (consulting fee, research support). **Mauro Zaffaroni:** Biogen Italia, Merck Serono, Sanofi, Teva (refunds of travel expenses for attending congresses). **Sandra Yukusic:** Biogen, GeNeuro, MedDay, Merck Serono, Novartis, Roche, Sanofi (consultancy fees, speaking fees, or honoraria, research support); Celgene, Teva (consultancy fees, speaking fees, or honoraria). **Steve Vucic:** CSL Behring, Merck Serono (honoraria). **Juwon Oh:** Biogen, Roche (consulting or speaking fees, research support); Celgene, EMD Serono, Genzyme, Novartis (consulting or speaking fees); Sanofi (research support). **Klaus Tiel-Wilck:** Almirall, Bayer, Biogen, Genzyme, Ipsen, Merck Serono, Merz Pharma, Novartis, Roche, Sanofi, Teva (honoraria for lectures, studies, and consultancy). **Anat Achiron:** Bayer, Biogen, Genzyme, Merck Serono, Roche, Sanofi (honoraria for advisory boards, lectures, studies, and consultancy). **Darren P. Baker, Janneke Wingerden, Elizabeth M Poole:** Sanofi (employee with ownership interest). **Bhupendra O. Khatri:** Acorda, Celgene, Serono (consulting/honorarium [consulting work, speaker programs]); Alexion, Biogen, Genentech, Novartis, Sanofi (consulting/honorarium [consulting work, speaker programs], contracted research); Ra Pharmaceuticals, Teva (contracted research).

Keywords: Disease duration and age in MS, Disease-modifying treatments in MS, Natural history of MS

(DXT73) Updated Safety of Cladribine Tablets in the Treatment of Patients with Multiple Sclerosis: Integrated Safety Analysis and Postapproval Data

Stuart Cook,¹ Gavin Giovannoni,^{2,3} Thomas P. Leist,⁴ Giancarlo Comi,⁵ Sana Syed,⁶ Axel Nolting,⁷ Doris Damian,⁸ Regina Schick⁹

¹Rutgers, The State University of New Jersey, New Jersey Medical School, Newark, NJ; ²Blizard Institute of Cell and Molecular Science, London, United Kingdom; ³Queen Mary University of London, London, United Kingdom; ⁴Jefferson University Hospital, Philadelphia, PA; ⁵Department of Neurology and Institute of Experimental Neurology, Università Vita-Salute San Raffaele, Milan, Italy; ⁶EMD Serono Inc, Billerica, MA; ⁷Merck KGaA, Darmstadt, Germany

Background: Pooling of long-term safety data for integrated analysis from the clinical trial program allows comprehensive characterization of the cladribine tablets (CTs) 10 mg (3.5 mg/kg cumulative dose over 2 years [CT3.5]) safety profile in patients with relapsing multiple sclerosis (RMS). **Objectives:** To provide an update to the previously reported treatment-emergent adverse event (TEAE) profile for CTs using the latest integrated safety data from clinical studies, including final data from the PREMIERE registry, and report postapproval safety data from Europe. **Methods:** The monotherapy oral cohort (CT3.5, N = 923, patient-years [PYs]=3936.69; placebo, N = 641, PYs = 2421.47) was derived from the CLARITY, CLARITY Extension, and ORACLE MS trials and the PREMIERE registry. Incidences per 100 PYs were calculated for adverse events, cumulative to the end of PREMIERE. Adverse drug reactions (ADRs) including serious ADRs (SADRs; implied causality) from postapproval sources are summarized. **Results:** Demographics reported at first dosing date were balanced among treatment groups: mean age (CT3.5 = 37.8 years; placebo = 37.2 years), proportion of females (CT3.5 = 66.3%; placebo = 66.1%), and proportion of patients with prior disease-modifying drug experience (CT3.5 = 19.9%; placebo = 20.4%). Incidences per 100 PYs for having ≥1 serious TEAE were 3.80 (CT3.5) and 3.05 (placebo). Incidences per 100 PYs for serious lymphopenia (preferred term) were 0.10 for CT3.5 and 0 for placebo. For serious infections and infestations (system organ class), incidences per 100 PYs were 0.60 (CT3.5) and 0.42 (placebo); for serious herpes zoster (preferred term): 0.05 (CT3.5) and 0 (placebo). Incidences per 100 PYs for malignant tumors were 0.26 (CT 3.5) and 0.12 (placebo). Separately, 922 ADRs from postapproval sources were reported in the Periodic Benefit-Risk Evaluation Report, of which 136 were SADRs; none of which are new safety findings for CT3.5. **Conclusions:** This integrated analysis of trial data, exclusively focused on the frequency of serious TEAEs with CT3.5 in patients with RMS, further establishes the safety profile of this dose. This profile is consistent with the previously published integrated safety analysis profile. No new major safety findings were identified in this latest dataset which includes final data from the PREMIERE registry. The pattern of postapproval ADRs was consistent with the clinical safety profile for CT3.5.

Supported by: None

Disclosure: **Stuart Cook:** Actinobac Biomed Inc, Biogen, Merck KGaA, Teva Pharmaceuticals (advisory board); Bayer HealthCare (advisory board, grant support); Neurology Reviews, Sanofi Aventis (consulting fee). **Gavin Giovannoni:** AbbVie, Actelion, Almirall, Atara Bio, Bayer Schering Pharma, Five Prime,

GlaxoSmithKline, GW Pharma, Merck KGaA, Pfizer Inc, Protein Discovery Laboratories, Sanofi Genzyme, Teva Pharmaceuticals Industries Ltd, UCB, Vertex Pharmaceuticals (consulting fee); Biogen, Ironwood, Merck and Co, Novartis (consulting fee, unrelated research support). **Thomas P. Leist:** Acorda, Bayer, Biogen, Daiichi, EMD Serono, Novartis, Ono, Pfizer, Teva Neuroscience (consulting fee). **Giancarlo Comi:** Almirall SpA, Biogen, Biogen Italia Srl, Celgene Group, EXCEMED, F. Hoffman-La Roche, Forward Pharma, Genzyme Corporation, Genzyme Europe, MedDay, Merck KGaA, Merck Serono SpA, Novartis, Roche SpA, Sanofi Genzyme, Teva Italia Srl, Teva Pharmaceutical Industries Ltd (consulting fee). **Sana Syed, Doris Damian:** EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). **Axel Nolting, Regina Schick:** Merck KGaA (salary).

Keywords: Disease-modifying treatments in MS

(DXT74) An Analysis of the Relationship Between Cladribine Dose and Risk of Malignancies in Patients with Multiple Sclerosis

Stuart Cook,¹ Gavin Giovannoni,^{2,3} Thomas P. Leist,⁴ Giancarlo Comi,⁵ Axel Nolting,⁶ Elke Sylvester,⁶ Dominic Jack,⁶ Doris Damian,⁷ Andrew Galazka⁸

¹Rutgers, The State University of New Jersey, New Jersey Medical School, Newark, NJ; ²Blizard Institute of Cell and Molecular Science, London, United Kingdom; ³Queen Mary University of London, London, United Kingdom; ⁴Jefferson University Hospital, Philadelphia, PA; ⁵Department of Neurology and Institute of Experimental Neurology, Università Vita-Salute San Raffaele, Milan, Italy; ⁶Merck KGaA, Darmstadt, Germany; ⁷EMD Serono Inc, Billerica, MA; ⁸Merck, Aubonne, Switzerland

Background: Malignancy risk was previously characterized in a monotherapy oral cohort of patients with multiple sclerosis (MS) treated with cladribine tablets (CTs) 10 mg (3.5 mg/kg cumulative dose over 2 years; referred to as CT3.5) including cumulative data up to February 2015. In clinical studies, an imbalance in the number of malignancies with CT3.5 vs placebo was observed, suggesting malignancy risk may be increased. **Objectives:** To provide a more detailed assessment of malignancy using safety data integrated from clinical trials and a safety follow-up registry (up to May 2017), to further characterize the malignancy risk of CTs in patients with MS and investigate whether there is a dose-dependent risk. **Methods:** Cohorts were monotherapy oral: patients with MS receiving CTs at any dose as a monotherapy; all exposed: patients with MS receiving any formulation of cladribine to provide a larger cohort to identify rare events such as malignancies. **Results:** In the monotherapy oral cohort, patient numbers (patient-years [PYs]) were: placebo N = 641 (2275), CT3.5 N = 923 (3754), CT 5.25 mg/kg (CT5.25) N = 632 (2610). The incidence per 100 PYs for malignant tumors during the entire follow up was: placebo: 0.13, CT3.5: 0.27, and CT5.25: 0.23. The risk difference vs placebo was: CT3.5: 0.14 (95% CI -0.14 to 0.38) and CT5.25: 0.10 (-0.18 to 0.39). In CLARITY CT3.5 and CT5.25 patients randomized to CT3.5 in CLARITY Extension (N = 195 for each treatment group), incidence per 100 PYs by CLARITY CT dose was: 0.55 (CT3.5, 1301 PYs) and 0.31 (CT5.25, 1286 PYs) for the entire follow-up; 0.91 (CT3.5, 790 PYs) and 0.52 (CT5.25, 784 PYs) for the period following initiation of treatment in year 3. An analysis of the all exposed cohort (cladribine N = 1976; placebo N = 802) stratified by cumulative cladribine dose gave the number of patients with a malignant event (incidence per 100 PY) as: >0-3.5 mg/kg = 6 (0.37), >3.5-5.25 = 14 (0.40), >5.25-7.0 = 6 (0.29), >7.0-8.75 = 6 (0.46), >8.75 = 2 (0.21). No hematological malignancies were observed at any time in the pooled dataset. **Conclusions:** Overall, there was no clear evidence of a dose effect of cladribine on malignancy risk in patients with MS based on >9500 PYs of cladribine exposure.

Supported by: None

Disclosure: **Stuart Cook:** Actinobac Biomed Inc, Biogen, Merck KGaA, Teva Pharmaceuticals (advisory board); Bayer HealthCare (advisory board, grant support); Neurology Reviews, Sanofi Aventis (consulting fee). **Gavin Giovannoni:** AbbVie, Actelion, Almirall, Atara Bio, Bayer Schering Pharma, Five Prime, GlaxoSmithKline, GW Pharma, Merck KGaA, Pfizer Inc, Protein Discovery Laboratories, Sanofi Genzyme, Teva Pharmaceuticals Industries Ltd, UCB, Vertex Pharmaceuticals (consulting fee); Biogen, Ironwood, Merck and Co, Novartis (consulting fee, unrelated research support). **Thomas P. Leist:** Acorda, Bayer, Biogen, Daiichi, EMD Serono, Novartis, Ono, Pfizer, Teva Neuroscience (consulting fee). **Giancarlo Comi:** Almirall SpA, Biogen, Biogen Italia Srl, Celgene Group, EXCEMED, F. Hoffman-La Roche, Forward Pharma, Genzyme Corporation, Genzyme Europe, Merck KGaA, Merck Serono SpA, Novartis, Roche SpA, Sanofi Genzyme, Teva Italia Srl, Teva Pharmaceutical Industries Ltd (consulting fee). **Axel Nolting, Elke Sylvester, Dominic Jack:** Merck KGaA (salary). **Doris Damian:** EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). **Andrew Galazka:** Merck KGaA, Aubonne, Switzerland (a business of Merck KGaA, Darmstadt, Germany) (salary).

Keywords: Disease-modifying treatments in MS

(DXT75) Switches to Established and Recently Approved Oral Disease-Modifying Therapies: Comparison of Patient Clinical Profiles and Therapy Selection Drivers

Virginia R. Schobel,¹ Jennifer Robinson²

¹Neurology Market Intelligence, Spherix Global Insights, Exton, PA; ²President, Spherix Global Insights, Exton, PA

Background: With the approval of siponimod (BAF), cladribine (CdA), and diroximel fumarate (DRF) in the United States, the number of oral disease-modifying therapies (DMTs) for the treatment of multiple sclerosis (MS) has grown. **Objectives:** To compare clinical profile, treatment history, and switch drivers among patients with MS recently switched to an oral DMT. **Methods:** In February 2019, 209 US neurologists contributed to a cross-sectional, retrospective chart audit of patients with MS (n = 1003 total; n = 718 relapsing-remitting MS [RRMS]) switched to a new DMT no more than 3 months prior. Patients were characterized by likelihood of alternative switch to DMTs in development, if the therapies had been available at switch. Analyses will be updated with February 2020 data. **Results:** Most patients recently switched to an oral DMT were diagnosed with RRMS (87%-92%). Oral DMTs constituted 43% of RRMS switches, with 11% switched to teriflunomide (TLF), 16% to fingolimod (FTY), and 17% to dimethyl fumarate (DMF). Oral RRMS switches were predominantly first switches (85%-89%); such switches were frequently due to efficacy (39%-43%) or tolerability (25%-29%). Patients switched to FTY were more likely to have switched from another oral DMT compared to those switched to DMF (21% vs 7%, $P < .05$). Desire for a high-efficacy DMT drove more FTY switch selection (60% vs TLF: 32%, DMF: 35%; $P < .05$), whereas patient request (30% vs FTY: 15%; $P < .05$) and lack of monitoring (16% vs FTY: 2%; $P < .05$) were more influential in DMF switches and long-term safety in TLF switches (39% vs FTY: 20%; $P < .05$). Compared to the established oral DMTs, candidates for alternative BAF, CdA, or DRF switches were less likely to be diagnosed with RRMS and more likely to have had a second or later switch. Patients with RRMS considered CdA candidates were more likely to have switched from an oral DMT compared to noncandidates (22% vs 12%; $P < .05$); administration type preference (43% vs 26%; $P < .05$), good tolerability profile (49% vs 38%; $P < .05$), and long-term safety (34% vs 22%; $P < .05$) drove more of the switched-to DMT selections among CdA candidates. DRF candidates were more likely to have switched due to tolerability issues compared to noncandidates (34% vs 16%; $P < .05$). **Conclusions:** Switches to established oral DMTs are typically first switches among patients with RRMS, although FTY may be reserved as a high-efficacy option for patients in whom a prior oral DMT failed. Oral class impact will grow within the switch segment with the availability of new oral options; however, the recently approved therapies will initially be niched as later line options. 2020 chart audit data will assess early adoption patterns and selection drivers among patients switched to BAF, CdA, or DRF.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Disease-modifying treatments in MS

(DXT76) First-Line Ocrelizumab Use for Relapsing-Remitting Multiple Sclerosis in the United States: Trend and Comparison to Glatiramer Acetate and Dimethyl Fumarate

Virginia R. Schobel,¹ Jennifer Robinson²

¹Neurology Market Intelligence, Spherix Global Insights, Exton, PA; ²President, Spherix Global Insights, Exton, PA

Background: In March 2017, ocrelizumab (OCR) was approved for the treatment of relapsing-remitting and primary progressive multiple sclerosis (MS), irrespective of prior therapy exposure. The relapsing-remitting MS (RRMS) indication was updated in July 2019 to encompass all relapsing forms of MS, including clinically isolated syndrome, RRMS, and active secondary progressive MS. **Objectives:** To trend OCR uptake among previously treatment-naïve patients with RRMS and compare characteristics of patients with RRMS initiated on first-line OCR to those initiated on one of the more established platform disease-modifying therapies (DMTs), glatiramer acetate (GA) or dimethyl fumarate (DMF). **Methods:** A retrospective, cross-sectional chart audit of patients with MS who initiated their first DMT no more than 3 months prior (ie, new-start patients) has been conducted with US neurologists (2016 n = 242; 2017 n = 274; 2018 n = 213) each December. Data were based on contributed RRMS patient chart reviews (2016 n = 777; 2017 n = 801; 2018 n = 758). GA includes patients treated with either a branded or generic agent. December 2019 data will be included at presentation. **Results:** In 2018, OCR was prescribed to 5%, GA to 32%, and DMF to 19% of new-start

patients with RRMS—stable with prior years. Although neurologists are more likely to agree with being comfortable using OCR first line for patients with RMS (57% vs 38% in 2017; $P < .05$), statement agreement did not correlate significantly with OCR share. While OCR patients were more likely to be male (OCR: 43%, GA: 24%, DMF: 26%), mean age or age category allocation did not differ between subgroups. Patients with RRMS initiated on OCR were more likely to have a perceived unfavorable long-term prognosis (OCR: 49%, GA: 11%, DMF: 8%) and to be having disability progression (OCR: 24%, GA: 11%, DMF: 9%). In comparison, patients taking GA and DMF were more likely to have no symptoms per the modified Rankin Scale (OCR: 5%, GA: 29%, DMF: 33%). Patients taking OCR had a greater mean T2 lesion burden at the most recent magnetic resonance imaging scan (OCR: 6.9, GA: 4.3, DMF: 4.1) and, since first under the contributing neurologist's care, were more likely to have had a Gd-enhancing lesion(s) (OCR: 38%, GA: 11%, DMF: 7%), relapse(s) (OCR: 48%, GA: 22%, DMF: 21%), decreased walking speed (OCR: 38%, GA: 8%, DMF: 12%), and/or onset or progression of cognitive symptoms (OCR: 19%, GA: 5%, DMF: 2%). High efficacy desire, dosing schedule preference, peer recommendation, and patient nonadherence concern played significantly more of a role in the first-line selection of OCR compared to GA or DMF. **Conclusions:** Regardless of neurologists' perceived comfort with first-line OCR, use among new-start patients with RRMS has remained flat during the first 2 years of availability. Compared to platform DMTs, OCR is most likely to be selected when a high-efficacy agent is required due to risk factors suggesting a poor long-term prognosis at the time of DMT treatment initiation.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Disease-modifying treatments in MS

(DXT77) Alemtuzumab Maintains Efficacy on Clinical and Magnetic Resonance Imaging Lesion Outcomes, Including Slowing of Brain Volume Loss, Over 9 Years in Relapsing-Remitting Multiple Sclerosis Patients: CARE-MS II Follow-up (TOPAZ Study)

Barry A. Singer,¹ Raed Alroughani,² Ann D. Bass,³ Simon Broadley,⁴ Yang Mao-Draayer,⁵ Hans-Peter Hartung,⁶ Eva Kubala Havrdova,⁷ Ho Jin Kim,⁸ Kunio Nakamura,⁹ Carlos Navas,¹⁰ Alex Rovira,¹¹ Krzysztof W. Selmaj,¹² Patrick Vermersch,¹³ Sibyl Wray,¹⁴ Zia Choudhry,¹⁵ Nadia Daizadeh,¹⁵ Salman Afsar,¹⁵ Giancarlo Comi,¹⁶ on behalf of the CARE-MS II, CAMMS03409, and TOPAZ Investigators

¹The MS Center for Innovations in Care, Missouri Baptist Medical Center, St. Louis, MO; ²Amiri Hospital, Sharq, Kuwait; ³Neurology Center of San Antonio, San Antonio, TX; ⁴Griffith University School of Medicine, Gold Coast Campus, Southport, QLD, Australia; ⁵University of Michigan Medical School, Ann Arbor, MI; ⁶Department of Neurology, UKD, Heinrich-Heine-University Düsseldorf, Düsseldorf, Germany; ⁷First Medical Faculty, Charles University, Prague, Czech Republic; ⁸Research Institute and Hospital of National Cancer Center, Goyang, Korea, Republic of (South); ⁹Cleveland Clinic, Cleveland, OH; ¹⁰Clínica Universitaria Colombia, Bogotá, Colombia; ¹¹Vall d'Hebron University Hospital, Barcelona, Spain; ¹²University of Warmia and Mazury, Olsztyn, Poland; ¹³University of Lille, INSERM U995, CHU Lille, Lille, France; ¹⁴Hope Neurology, Knoxville, TN; ¹⁵Sanofi, Cambridge, MA; ¹⁶University Vita-Salute San Raffaele, Milan, Italy

Background: In CARE-MS II (trial registration: NCT00548405), alemtuzumab (12 mg/day; baseline: 5 days; 12 months later: 3 days) significantly improved clinical/magnetic resonance imaging (MRI) outcomes vs subcutaneous interferon beta-1a (SC IFN β -1a) over 2 years (y) in patients with RRMS with inadequate response to prior therapy. Efficacy was maintained in a 4-year extension study (NCT00930553), wherein patients could receive additional alemtuzumab courses (12 mg/day on 3 days, ≥ 12 months apart) as needed for disease activity, or receive other disease-modifying therapy (DMT) per investigator discretion. Further follow-up was available in an additional 5-year extension, TOPAZ (NCT02255656). **Objectives:** Evaluate the efficacy and safety of alemtuzumab in CARE-MS II patients over 9 years. **Methods:** At investigator discretion, patients in TOPAZ can receive additional as-needed alemtuzumab (≥ 12 months apart, no criteria) or receive other DMT (at any time). **Results:** From core study baseline through year 9, 288/435 (66%) CARE-MS II alemtuzumab-treated patients remained on study; 41% received neither additional alemtuzumab nor another DMT through year 9. Annualized relapse rate was 0.19 in years 3-9. From core study baseline through year 9, 68% of patients had stable/improved Expanded Disability Status Scale (EDSS) scores, and mean change in EDSS was +0.32. Over 9 years, 60% of patients were free of 6-month confirmed disability worsening and 49% achieved 6-month confirmed disability improvement. On average, 69% of patients were free of MRI disease activity, 89% were free of new gadolinium-enhancing lesions, and 69% were free of new/enlarging T2 hyperintense lesions, annually from year 3-9. From core study baseline through year 9, median percent cumulative

brain volume (BV) change was -1.22% ; median percent BV change was $\leq -0.19\%$ annually over years 3-9. Incidence of overall adverse events (AEs) and infections declined through year 9; cumulative thyroid AE incidence was 43.7% and immune thrombocytopenia (ITP) incidence was 3.7% (1 new case of ITP, 14 months after the fourth alemtuzumab course, was observed at year 9). No new cases of nephropathy were reported. Efficacy and safety in SC IFN β -1 α -treated patients from the core study who switched to alemtuzumab in the extension were consistent with those treated with alemtuzumab both in the core and extension. **Conclusions:** Efficacy of alemtuzumab on clinical, MRI, and BV loss outcomes was maintained over 9 years in CARE-MS II patients, with 41% receiving no further treatment through year 9. Safety in year 9 in this study was consistent with that of previous years.

Supported by: Sanofi, Bayer HealthCare Pharmaceuticals

Disclosure: Barry A. Singer: AbbVie, Biogen, Novartis, Roche, Sanofi (research support, speaking and/or consulting); Acorda, Alexion, Bayer, Celgene, EMD Serono, Genentech, Teva, TG Therapeutics (speaking and/or consulting); Alkermes, MedImmune (research support). Raed Alroughani: Bayer, Biogen, Biologix, Genpharm, GlaxoSmithKline, Lundbeck, Merck Serono, Novartis, Sanofi (speaker honoraria, scientific advisory boards, research grants). Ann D. Bass: Biogen, EMD Serono, Mallinckrodt, Novartis, Roche-Genentech, Sanofi, TG Therapeutics (consulting fees/fees for non-CME services from commercial interests or their agents/sponsor and research support). Simon Broadley: Bayer Schering, Merck Serono, Novartis, Sanofi (conference travel sponsorship, honoraria for advisory board participation); Biogen (conference travel sponsorship, honoraria for advisory board participation, speaker honoraria, unencumbered research grant); Genzyme (speaker honoraria). Yang Mao-Draayer: Acorda, Bayer, Biogen, Celgene, Chugai, EMD Serono, Genzyme, Novartis, Questor, Teva Neuroscience (consulting fees and/or grant support); NIH National Institute of Allergy and Infectious Diseases Autoimmune Center of Excellence: UM1-AI110557, NIH National Institute of Neurological Disorders and Stroke R01-NS080821 (grant support). Hans-Peter Hartung: Bayer, Biogen, CSL Behring, Grijfols, Merck Serono, Novartis, Roche, Sanofi (consulting and/or speaking fees). Eva Kubala Havrdova: Actelion, Biogen, Merck Serono, Novartis, Receptos, Roche, Sanofi, Teva (honoraria, grant support); Ministry of Education of Czech Republic [PROGRES Q27/LF1] (support). Ho Jin Kim: Bayer Schering, Biogen, Celltrion, Eisai, HanAll BioPharma, Novartis (consulting and speaking fees); Genzyme, Merck Serono, Teva-Handok, UCB (consulting and speaking fees, research support); Journal of Clinical Neurology, Multiple Sclerosis Journal - Experimental, Translational and Clinical (co-editor/associate editor); MedImmune (consulting and speaking fees, steering committee member); Ministry of Science and ICT (research support). Kunio Nakamura: Biogen (royalty fees for licenses, speaking fees, research support); Novartis, Sanofi (speaking fees, research support). Carlos Navas: Bayer Schering, Genzyme, Stendhal (consulting and speaking fees); Merck Serono, Novartis, Roche (consulting and speaking fees, research support). Alex Rovina: Bayer, Biogen, Bracco, icometrix, Novartis, OLEA Medical, Roche, Sanofi, SyntheticMR (consulting and/or speaking fees). Krzysztof W. Selma: Biogen, Merck, Novartis, Roche, Sanofi, Synthon (consulting and/or speaking fees). Patrick Vermersch: Almirall, Biogen, Celgene, Merck, Novartis, Roche, Sanofi, Servier, Teva (consulting and/or speaking fees, research support). Sibyl Wray: Alkermes, Bayer, Biogen, EMD Serono, Genentech/Roche, Novartis, Sanofi, Teva, TG Therapeutics (consulting, principal investigator, and/or speaking fees). Zia Choudhry, Nadia Daizadeh, Salman Afsar: Sanofi (salary). Giancarlo Comi: Actelion (consulting fee); Bayer Schering, Merck Serono, Novartis, Sanofi, Teva (consulting fee, lecture fees); Biogen Dompé, Serono Symposia International Foundation (lecture fees).

Keywords: Disease-modifying treatments in MS

(DXT78) The FLUENT Study: Changes in Immune Cell Profile, and in Clinical and Safety Outcomes, in Fingolimod-Treated Patients with Relapsing Multiple Sclerosis

Yang Mao-Draayer,¹ Jeffrey A. Cohen,² Amit Bar-Or,³ May H. Han,⁴ Barry A. Singer,⁵ Nina Jaitly,⁶ Scott Kolodny,⁶ Chelsea Elam,⁶ Xiangyi Meng,⁶ Marina Ziehn,⁶ Bruce A.C. Cree⁷

¹Autoimmunity Center of Excellence, Multiple Sclerosis Center, University of Michigan, Ann Arbor, MI; ²Mellen Center, Cleveland Clinic, Cleveland, OH; ³Center for Neuroinflammation and Experimental Therapeutics and Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ⁴Stanford University, Stanford, CA; ⁵Missouri Baptist Medical Center, St. Louis, MO; ⁶Novartis Pharmaceuticals Corporation, East Hanover, NJ; ⁷Department of Neurology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA

Background: Fingolimod inhibits sphingosine 1-phosphate receptor-mediated lymphocyte egress from lymph nodes, reducing lymphocytic infiltration of the central nervous system in multiple sclerosis (MS). However, characterization of immune cell changes in relation to patient outcomes in MS remains incomplete. **Objectives:** The FLUENT study

explored fingolimod-mediated immune-cell subset changes in the innate and adaptive immune systems and their association with efficacy and safety outcomes in relapsing MS. **Methods:** FLUENT was a 12-month, prospective, multicenter, nonrandomized phase 4 study in adults with relapsing MS receiving fingolimod 0.5 mg/day (trial registration: NCT03257358). Patients were stratified as fingolimod-naïve (cohort 1) or previously treated with fingolimod for ≥ 2 years (cohort 2). The primary outcome was change from baseline in immune cell subsets. Secondary outcomes included anti-John Cunningham virus (JCV) antibody status, neurofilament light-chain (NfL) concentrations, Patient-Determined Disease Steps (PDDS) score, and incidence of adverse events (AEs). Analyses were in patients completing month-12 follow-up. **Results:** At baseline, proportions of CD4+ T cells, CD8+ T cells, and B cells were higher in cohort 1 (n = 163) than in cohort 2 (n = 217). Cell counts in these immune subsets were substantially reduced at month 12 compared with baseline in cohort 1, with naïve and central memory T cells affected more than effector memory T cells and regulatory B cells; changes in these subsets were less pronounced in cohort 2 than in cohort 1. At baseline, 54% of cohort 1 and 64% of cohort 2 were anti-JCV positive, and anti-JCV antibody index was unchanged at month 12. Mean (SD) NfL concentration in cohort 1 was 12.16 (11.05) pg/mL at baseline and 8.57 (5.32) at month 12; NfL concentrations in cohort 2 were similar at baseline (9.59 [7.55]) and month 12 (9.78 [8.88]). Baseline PDDS scores were low (cohort 1, 1.7; cohort 2, 1.8) and unchanged at month 12. Respective rates of treatment-emergent AEs (54.6% vs 44.2%) and AEs leading to treatment discontinuation (12.3% vs 5.5%) were higher in cohort 1 than cohort 2; rates of serious AEs (5.5% vs 5.5%) were similar between cohorts. **Conclusions:** These findings expand understanding of temporal changes in immune cell subtype profiles and biomarkers in patients with relapsing MS receiving fingolimod.

Supported by: None

Disclosure: Yang Mao-Draayer: Acorda, Bayer, Biogen, Celgene, EMD Serono, Genentech, Teva (consulting fee); Chugai, Novartis, Sanofi Genzyme (consulting fee, contracted research); National Institute of Allergy and Infectious Diseases Autoimmune Center of Excellence, NIH National Institute of Neurological Disorders and Stroke (contracted research). Jeffrey A. Cohen: Convelo, Population Council (consulting fee); Multiple Sclerosis Journal (editor); Mylan (speakers' bureau). Amit Bar-Or: Atara Biotherapeutics, Bayer, Bayhill Therapeutics, Berlex, Biogen, BioMS, Celgene/Receptos, Diogenix, Eli Lilly, F. Hoffman-La Roche, Genentech, GlaxoSmithKline, Guthy-Jackson/GGF, Immunic, Janssen/Actelion, Mapi, MedImmune, Merck/EMD Serono, Novartis, Ono Pharmacia, Rochel Genentech, Sanofi Genzyme, Teva Neuroscience, Wyeth (consulting fee). May H. Han: Novartis (consulting fee). Barry A. Singer: AbbVie, Biogen, Novartis, Sanofi Genzyme (consulting fee, contracted research); Acorda, Alexion, Celgene, EMD Serono, Genentech, Teva, TG Therapeutics (consulting fee); Alkermes, MedImmune, Roche (contracted research). Nina Jaitly, Scott Kolodny, Chelsea Elam, Xiangyi Meng: Novartis Pharmaceuticals Corporation (salary). Marina Ziehn: Novartis Pharma AG (salary). Bruce A.C. Cree: Akili, Alexion, Atara, Biogen, EMD Serono, Novartis, TG Therapeutics (consulting fee).

Keywords: Disease-modifying treatments in MS

(DXT79) Efficacy of Ocrelizumab Treatment on Cognitive Functions in Persons with Multiple Sclerosis

Serkan Ozakbas,¹ Ozge Sagici,² Zuhal Abasiyanik,² Tuba Ozdogar,² Pinar Yigit¹

¹Department of Neurology, Dokuz Eylul University, Izmir, Turkey; ²Dokuz Eylul University, Izmir, Turkey

Background: Ocrelizumab is the first treatment which could be used for progressive forms of multiple sclerosis (MS). Generally ocrelizumab trials in MS investigate side effect and safety properties. There are limited studies examining effects of ocrelizumab on cognition in patients with MS (PwMS). **Objectives:** The aim of this study was to evaluate the efficacy of ocrelizumab treatment on cognitive functions in PwMS. **Methods:** In total, 35 PwMS were included in this study. The participants' clinical characteristics such as MS type, disease duration, and Expanded Disability Status Scale (EDSS) scores were recorded. Cognitive function was evaluated with The Brief International Cognitive Assessment for Multiple Sclerosis (BICAMS) battery, which consists of the Symbol Digit Modalities Test (SDMT), the California Verbal Learning Test-II (CVLT-II), and the Brief Visuospatial Memory Test-Revised (BVMTR). The assessment was done at baseline and 6 months after the treatment. **Results:** The average disease duration was 16.84 \pm 8.19 and EDSS score was 6.23 \pm 1.43. Participants' clinical characteristics of the disease were 11.4% relapsing-remitting (RRMS), 57.1% (n = 20) secondary progressive (SPMS), and 31.4% (n = 11) primary progressive MS (PPMS). BVMTR and CVLT-II scores were significantly increased at month 6 according to baseline assessment (20.71 \pm 7.85 vs 23.68 \pm 7.05, 45.03 \pm 11.97 vs 49.43 \pm 12.45,

respectively) ($P < .05$). No significant differences were observed between baseline and month 6 in terms of SDMT scores (34.37 ± 14.95 vs 34.51 ± 15.67) ($P > .05$). **Conclusions:** This study has suggested that ocrelizumab treatment could be affected positively on verbal and visual learning/memory. On the other hand, there was no positive or negative effect on information processing speed. In view of most of our patients being in progressive form, the protective or positive effect of ocrelizumab on cognitive functions is clinically important. Additionally, to better understand the effects of ocrelizumab treatment on cognitive functions, it is necessary to design studies with longer follow-up periods.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Disease-modifying treatments in MS, Psychological issues and MS

EPIDEMIOLOGY AND GENETICS

(EPI01) Determining the Effect of Early Versus Later Diagnosis of Multiple Sclerosis on Long-Term Prognosis in a Real-World Setting

Jikku Zachariah,¹ Rebecca Schorr,² Tim Quezada,¹ Thomas F. Scott^{1,3}

¹Department of Neurology and Neuroscience Institute, Allegheny General Hospital, Allegheny Health Network, Pittsburgh, PA; ²Highmark Health Data Science R&D, Highmark Health, Pittsburgh, PA; ³Department of Neurology, Drexel University College of Medicine, Philadelphia, PA

Background: Early treatment of multiple sclerosis (MS) is recommended based on studies involving patients identified early with clinically isolated syndromes. Although early diagnosis and subsequent early treatment might be expected to lead to a better outcome (lower Expanded Disability Status Scale score), in a real-world setting outcomes may be different concerning patients presenting (and treated) early after onset of symptoms (EP group) vs later (LP group). Other risk factors for worsening MS (poor recovery from first attack, motor onset, etc) may be more predictive than EP vs LP status. **Objectives:** To compare characteristics of EP patients (with less than 1 year of symptoms at presentation) vs LP groups. **Methods:** Newly diagnosed patients with MS seen up to 15 years divided as EP or LP were studied for attack type, frequency, and recovery, and group statistics were applied. **Results:** There were 121 patients in the EP group, and 86 patients in the LP group. More patients with a high attack frequency in years 0-2 were seen in the EP group vs the LP group (45% vs 28%, $P = .014$). The median time to treatment was shorter in the EP group vs LP group (by 32 months [8.91]), and we found no significant difference in disability long-term outcomes. Other clinical risk factors were evenly divided between EP and LP groups. **Conclusions:** These EP and LP groups were the same in terms of long-term outcome. Earlier treatment and more active disease were seen in the EP group. We suggest a nuanced approach to interpretation of the "early diagnosis and treatment equals better outcome" rule.

Supported by: None

Disclosure: Jikku Zachariah, Rebecca Schorr, Tim Quezada: Nothing to disclose. Thomas F. Scott: Genentech, Biogen, Novartis, Genzyme (consulting fee, contracted research, speakers' bureau).

Keywords: Disease-modifying treatments in MS, Epidemiology of MS, Natural history of MS

(EPI02) Motor Impairment in Multiple Sclerosis: Analysis from the North American Registry for Care and Research in Multiple Sclerosis (NARCRMS)

Kottil Rammohan,¹ David Li,² June Halper,³ Sara McCurdy Murphy,⁴ Trent Wallace,⁵ Lisa Patton,⁴ Sarah Chrisant,⁵ Chao Zheng,⁶ Seema Khurana⁶

¹Department of Neurology, Multiple Sclerosis Center, University of Miami, Miami, FL; ²The University of British Columbia, Vancouver, BC, Canada; ³CEO-CMCS, Hackensack, NJ; ⁴CRB, Social & Scientific Systems, Inc, Silver Spring, MD; ⁵CRB, Social & Scientific Systems, Inc, Durham, NC; ⁶University of Miami, Miami, FL

Background: The North American Registry for Care and Research in Multiple Sclerosis (NARCRMS) is a longitudinal registry studying the course of multiple sclerosis (MS) in the disease-modifying era. **Objectives:** To examine motor performance metrics of upper and lower extremity function in NARCRMS patients at enrollment. **Methods:** Recruitment began in 2016, and by December 31, 2019, 662 patients were enrolled at 23 MS sites across the United States and Canada. People with any subtype of MS within 15 years of disease onset and an Expanded Disability Status Scale (EDSS) score of up to 6.5 are eligible for enrollment. Various clinical metrics are collected including motor performance for upper and lower extremities. Our initial observations about EDSS, 25-foot

timed walk, and the Nine-Hole Peg Test (NHPT) are reported below. **Results:** EDSS scores and 25-foot walk times were available in 579 patients and upper extremity function in 571 patients. A mean walking speed of 4.9 seconds was recorded in patients with an EDSS score of 0 ($n = 100$). 5.0 seconds remained the mean speed until an EDSS score of 3.0 ($n = 37$), where a mean speed of 5.6 seconds was recorded. Walking truly became affected at an EDSS score of 3.5 ($n = 25$), where a mean speed of 6.1 seconds was recorded. Thereafter, mean speed progressively declined at every EDSS score increase. For an EDSS score of 4.0 ($n = 25$), mean speed was 7.9 seconds; for an EDSS score of 4.5 ($n = 6$), mean speed was 9.1 seconds and continued to increase until an EDSS score of 6.5 ($n = 10$), where mean speed was 16.8 seconds. For the NHPT, patients with an EDSS score of 0 ($n = 96$) had a mean speed of 19.4 seconds in the dominant and 20.7 seconds in the nondominant hand. Hand function remained unimpaired until an EDSS score of 2.0, and significant slowing occurred in patients with EDSS scores ranging from 2.5 to 6.5. For an EDSS score of 2.5 ($n = 40$), mean speed was 24.7 seconds in the dominant and 24 seconds in the nondominant hand. For an EDSS score of 4.0 ($n = 26$), mean speed was 26.1 seconds in the dominant and 26.6 seconds in the nondominant hand. For an EDSS score of 6.5 ($n = 15$), hand function had declined to a mean speed of 39.1 seconds for the dominant and 49.8 seconds for the nondominant hand.

Conclusions: A linear correlation of the 25-foot walk speed to EDSS score increases was remarkable, reiterating the commonly held belief that the EDSS is a "walking scale." Decline in hand function at an EDSS score of 2.5 was unexpected, because hands are often perceived to be unaffected early in MS and seldom observed as impaired by patients. Progressive decline of hand function at every EDSS score increase would suggest that the NHPT test is a good marker of declining hand function and should be included in clinical monitoring of patients.

Supported by: None

Disclosure: Kottil Rammohan: Alexion, Biogen, EMD Serono, Genentech, Novartis (consulting fee, grant); Department of Defense, TG Therapeutics (grant); Genzyme (consulting fee). David Li: Roche, Sanofi Genzyme (contracted research); UBC MSIMRI Research Group (salary); Vertex Pharmaceuticals (consulting fee). June Halper, Sara McCurdy Murphy, Trent Wallace, Lisa Patton, Sarah Chrisant, Chao Zheng, Seema Khurana: Nothing to disclose.

Keywords: Epidemiology of MS

(EPI03) Increase in Family Recurrence in Patients Diagnosed with Multiple Sclerosis in the Years 2017-2019 in Hispanic Population of Puerto Rico

Viviana Martinez, Astrid Diaz, Cristina M. Rubi, Ivonne Vicente, Angel Chinea
San Juan Multiple Sclerosis Center, Guaynabo, PR

Background: Multiple sclerosis (MS) is a neurodegenerative disease in which the immune system damages the central nervous system. The cause of MS is not known, but several studies look at environmental, immunologic, geographic, and genetic factors. Thus, MS is not considered hereditary, but rather polygenic. However, there are several cases of patients with family history of MS in several relative degrees. **Objectives:** Evaluate family recurrence of MS among patients diagnosed with MS in recent years in Puerto Rico (PR), including any type of degree relative. **Methods:** For this, data from the Puerto Rican MS Registry (PRMS Registry) of all patients diagnosed in the years 2017, 2018, and 2019 registered at present in PR were analyzed. **Results:** Overall 11.4% (45/396) of patients had family recurrence of MS. For 2017, 9.7% (14/143) of patients had family history of MS of at least 1 family member. For 2018, 8.3% (14/167) of patients presented family history of MS. Lastly, for 2019 a total of 19.7% (17/86) of patients reported having family history of MS. **Conclusions:** A slight increase in recurrence was observed when compared to the previous study from 2013 to 2016 (10.2%). Further investigations need to be done to elucidate the genetic aspects of family recurrence of MS among the Puerto Rican population. The genetic mix of Caucasian, African, and Taino races could have an influence on genetic risk among this population. Also, it is important to keep this study ongoing to analyze familial risk in the Hispanic population and compare it to other ethnic groups.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Epidemiology of MS, Genetics and MS

(EPI04) Diet Quality and Nutritional Adequacy of Micronutrients Among People with Relapsing-Remitting Multiple Sclerosis: An Analysis of Weighed Food Records

Tyler J. Titcomb,¹ Linda G. Snetselaar,² Terry L. Wahls¹

¹Department of Internal Medicine and ²Department of Epidemiology, University of Iowa, Iowa City, IA

Background: Multiple sclerosis (MS) is a neurodegenerative disease that affects nearly 1 million in the United States. Poor diet quality and micronutrient deficiencies have been reported in several studies and are associated with more severe disease. However, previous work has relied on diet screeners and questionnaires for data collection. Thus, these findings need to be repeated using more precise data collection instruments such as weighed food records. **Objectives:** Weighed food records were used to evaluate diet quality and usual intake of micronutrients from people with diagnosed relapsing-remitting MS (RRMS). **Methods:** As part of a dietary intervention study comparing the Wahls and Swank diets, 3 weighed food records were collected on 2 weekdays and 1 weekend day at a prerandomization run-in visit from (n = 95) participants and again at a baseline visit from the (n = 87) nonexcluded participants. Food records obtained from all participants were analyzed at the University of Minnesota Nutrition Coordinating Center. Diet quality was assessed using the Healthy Eating Index (HEI), which compares food groups and selected nutrient intakes to the Dietary Guidelines for Americans. Mean intake of each micronutrient was calculated for each individual and adjusted using the National Cancer Institute method to estimate usual intake. Usual intake of each micronutrient was then evaluated with the estimated average requirement cutpoint method for each life stage group and combined by weighted means to assess the overall nutritional adequacy of each micronutrient for the group. **Results:** Preliminary analyses indicate that this cohort has an HEI score of 61 ± 12 , which suggests that diet quality needs improvement. Furthermore, this cohort has high prevalence of inadequate intake for vitamins D 92.9%, E 61.4%, C 50.8%, A 35.3%, folate 31.0%, and B₆ 22.8%, and minerals including calcium 49.8%, magnesium 45.8%, and zinc 19.5%. However, low prevalence of inadequate intake was observed for niacin 0.2%, thiamin 7.5%, riboflavin 3.0%, B₁₂ 5.4%, phosphorus 2.0%, copper 6.6%, and selenium 1.0%. **Conclusions:** Diet quality is low and intake of several micronutrients is inadequate in this cohort with RRMS. These findings confirm observations from previous studies that poor diet quality and inadequate intake of micronutrients is common among those with RRMS. These findings may lead to new dietary strategies to manage symptoms and improve quality of life among those with MS.

Supported by: None

Disclosure: Tyler J. Titcomb, Linda G. Snetselaar: Nothing to disclose. Terry L. Wahls: BioCeuticals, Genova Diagnostics, Institute for Functional Medicine, MCG Health LLC (consulting fee); Dr Terry Wahls LLC, FBB Biomed Inc, The Wahls Institute PLC, TZ Press LLC (ownership interest); Penguin Random House (royalty).

Keywords: Complementary/alternative therapies in MS, Comprehensive care and MS, Epidemiology of MS

FAMILY AND CAREGIVERS

(FAM01) Characterizing Predictors of Resilience Among Family Caregivers of People with Advanced Multiple Sclerosis Disability: Work in Progress

Afolasade Fakolade,¹ Odessa McKenna,¹ Mark S. Freedman,² Marcia Finlayson,³ Amy E. Latimer-Cheung,⁴ Lara A. Pilutti¹

¹Interdisciplinary School of Health Sciences, University of Ottawa, Ottawa, ON, Canada;

²University of Ottawa and the Ottawa Hospital Research Institute, Ottawa, ON, Canada;

³School of Rehabilitation Therapy, Queen's University, Kingston, ON, Canada; ⁴Kinesiology and Health Studies, Queen's University, Kingston, ON, Canada

Background: Providing ongoing support can adversely affect physical, psychological, and social health of care partners of people with advanced multiple sclerosis (MS). However, some care partners have also reported positive experiences and benefits of MS caregiving (eg, greater sense of commitment and pride in the role). This variability highlights the importance of understanding protective factors that can buffer against the adverse effects of caregiving. Psychological resilience describes positive adaptations to stressful situations and includes individual, community, and societal level factors. Research in other caregiving contexts has shown that resilience is associated with improved health outcomes and lower levels of care partner burden. Unfortunately, there is limited research on resilience in the context of MS caregiving. Generating this knowledge is important to guide the development of interventions to enhance resilience and to identify individuals most likely to benefit from future intervention. **Objectives:** To examine the relationships between resilience and a comprehensive set of individual, community, and societal level factors in the context of MS caregiving. **Methods:** A cross-sectional survey study. We

are collecting data as part of a pilot randomized controlled trial evaluating the feasibility and efficacy of a dyadic physical activity program for people with advanced MS and their care partners. Eligibility criteria include care partners who: 1) are ≥ 18 years old, 2) provide ≥ 1 hour/day of care, 3) are inactive, 4) are asymptomatic, and 5) are able to communicate in English. Care partners will complete demographics and general health questionnaires. The following scales will also be administered: 1) Caregiving Tasks in MS Scale, 2) Connor-Davison Resilience Scale, 3) Interpersonal Support Evaluation List-12, 4) Coping with MS Caregiving Inventory, 5) Measure of Experiential Aspects of Participation, and 6) Godin Leisure-Time Exercise Questionnaire. We will conduct regression modelling to identify predictors of resilience among care partners. **Results:** Data collection is ongoing. Anticipated completion date is March 2020. We will present findings on resilience and associated factors among MS care partners. **Conclusions:** Resilience may be an important protective factor against the adverse effects of MS caregiving. We anticipate that the findings from this study may have implications for interventions designed to enhance and sustain resilience among MS care partners.

Supported by: None

Disclosure: Nothing to disclose

Keywords: MS and the caregiver/family

(FAM02) Understanding Units of Energy: Key to Communicating Multiple Sclerosis Fatigue

Angela Jones,¹ Anne L. Diaz²

¹Health Services, Clark County School District, Las Vegas, NV; ²Kent, WA

Background: Fatigue, defined as the difficulty or inability to perform tasks due to a lack of sufficient energy, is one of the key symptoms of multiple sclerosis (MS). The level of fatigue is difficult to quantify and explain to a person without the condition. Interactions between persons with MS (PsWMS) and those without (PsWO) with whom they have interpersonal relationships can be damaged by this lack of understanding. The Roy Adaptation Model (RAM) (Roy C Sr, Andrews HA. The Roy Adaptation Model. 2nd ed. Appleton and Lange; 1999) provides a nursing framework through which to assess, intervene, and evaluate the effective communication between PsWMS and PsWO. The RAM views persons as adaptive entities in constant interaction with their environment. One portion of the theory, the interdependence adaptive mode, focuses on the communication between a person and his or her significant other/support group. Clear understanding of the message sent and received and the ensuing healthy adaptations to their interactive communication resulting in behavioral change are the goals. **Objectives:** By quantifying the amount of energy each activity takes, PsWMS will be able to communicate more clearly their level of energy and, inversely, their level of fatigue. Relational stress will decrease and communication between PsWMS and PsWO will improve. **Methods:** A MS support group composed of 23 persons, some with MS and some without, participated in a 2-hour educational, interactive session, which focused on defining units of energy and describing how many may be needed for both activities of daily living and special events. The PsWO "tried on" different symptoms of MS throughout the session to gain a better understanding of the challenges facing the PsWMS. At the same time, the PsWMS focused on self-assessment and became more aware of the frustrations expressed by the PsWO. Both subgroups practiced active listening techniques. **Results:** After 2 months, all 23 persons reported improved relationships due to decreased stress and a better understanding of the effects of MS. **Conclusions:** Teaching a simple method of communicating MS fatigue greatly improved the quality of life of PsWMS and those with whom they interact. Using units of energy within conversations improved communication and enabled more positive interdependent interactions with caregivers, family members, and their colleagues in the workplace.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Fatigue, Management of activities of daily living in MS, MS and the caregiver/family

INTERNET AND INFORMATION SERVICES

(IIS01) North American Registry for Care and Research in Multiple Sclerosis (NARCRMS) Model for Implementing OpenClinica Insight for Data Sharing and Visualization

Peter Gilbertson,¹ Trent Wallace,² Lydia Rogers,¹ Sara McCurdy Murphy,³ Lisa Patton,³ Sarah Chrisant,² David Li,⁴ June Halper,⁵ Kotil Rammohan⁶

¹Social & Scientific Systems, Inc, Silver Spring, MD; ²CRB, Social & Scientific Systems, Inc, Durham, NC; ³CRB, Social & Scientific Systems, Inc, Silver Spring, MD; ⁴University of British Columbia, Vancouver, BC, Canada; ⁵CEO-CMSC, Hackensack, NJ; ⁶Department of Neurology, Multiple Sclerosis Center, University of Miami, Miami, FL

Background: The North American Registry for Care and Research in Multiple Sclerosis (NARCRMS) is a clinic-based longitudinal registry of sites located in the United States and Canada. Active since 2015, the registry is a public-private partnership aiming to improve multiple sclerosis (MS) care and understanding as a database of clinical records and patient-centered outcomes. With enrollment, yearly follow-up and exacerbation-based visits, patients provide demographic, medical history, attack history, and health productivity information supplemented with physician-collected physical and cognitive assessments. In just a few years the registry has generated several hundred thousand points of data on a wide variety of health-related topics from the current 22 participating sites. The registry includes data from standardized interviews and physician evaluations and is adding self-administered patient-reported outcomes in the coming months. **Objectives:** To develop a model to share aggregated data from individual sites to enable participating sites and industry partners the ability to view and analyze larger, multisite datasets for research and exploration. **Methods:** OpenClinica is an open-source and software-as-a-service electronic data collection (EDC) system most often used for data collection and data management in a clinical setting. OpenClinica Insight, built on the open-source Metabase platform, is a data reporting and sharing tool available as part of OpenClinica's Enterprise system that connects directly to the EDC database and allows real-time data access, visualization, and downloading. To implement Insight, the team coordinated with project leadership, industry, and cooperating investigators to develop a process for defining roles and relationships, defining appropriate summary charts and graphs to summarize collected data, and defining data access parameters and restrictions. **Results:** OpenClinica Insight provides a platform to leverage limited standardized patient-derived data on a prospective basis. The platform is useful and end-user friendly and allows for efficient information sharing across the pool of geographically diverse clinical research sites to provide insights into local, regional, and continental patterns and standards of MS care. **Conclusions:** OpenClinica Insight is a powerful tool to report information from OpenClinica Enterprise, and the model developed from NARCRMS should serve as an example for integrating informatics from large databases developed to study natural history of various chronic disorders such as MS.

Supported by: None

Disclosure: *Peter Gilbertson: Sandford Health (salary). Trent Wallace, Lydia Rogers, Sara McCurdy Murphy, Lisa Patton, Sarah Chrisant, June Halper: Nothing to disclose. David Li: Roche, Sanofi Genzyme (contracted research); UBC MS/MRI Research Group (salary); Vertex Pharmaceuticals (consulting fee). Kottil Rammohan: EMD Serono, Biogen, Sanofi-Aventis, Genzyme, Novartis, Teva Neurosciences, Acorda, Roche/Genentech (consulting fee, speakers' bureau).*

Keywords: Epidemiology of MS

IMAGING

(IMG01) Conformance to CMSC Magnetic Resonance Imaging (MRI) Guidelines in a Real-World Multicenter MRI Dataset

Sophie Vercrayssen,¹ Arne Brys,¹ Melissa Verheijen,¹ Brandon Steach,² Eline Van Vlierberghe,² Diana M. Sima,¹ Dirk Smeets¹

¹R&D, icometrix, Leuven, Belgium; ²icometrix, Chicago, IL

Background: Acquiring magnetic resonance images (MRIs) in a standardized way allows early and accurate diagnosis of multiple sclerosis (MS) and patient follow-up. The Consortium of Multiple Sclerosis Centers (CMSC) guidelines for magnetic resonance (MR) imaging suggest a standardized protocol to improve the diagnosis and routine follow-up of MS. **Objectives:** This study evaluates how many MRI acquisitions from a real-world dataset of patients with MS satisfy the CMSC brain MRI guidelines (2018 revision). In particular, for every individual MR sequence, it is assessed how well the guidelines are met. **Methods:** CMSC brain MRI guidelines of 2018 impose that 4 different scans should be acquired: a 2D or 3D (sagittal and axial) fluid-attenuated inversion recovery (FLAIR), a 2D axial or 3D T2-weighted scan, an axial 2D diffusion-weighted image (DWI), and a 3D inversion-recovery prepared (IR-prep) gradient-echo T1. For every scan, an in-slice pixel resolution $\leq 1 \text{ mm} \times 1 \text{ mm}$, slice thickness $\leq 3 \text{ mm}$ with no slice gap, as well as whole-brain coverage are required. These requirements are checked on a multicenter MRI dataset from the United States, consisting of 1233 sessions, acquired in 581 different

centers/scanners from 2018 onwards. **Results:** None of all 1233 MRI sessions fully complied with the guidelines. For the T1 sequence, only 8% satisfied the criteria. For the other data, 23% did not have a T1 sequence, 73% had a too-large slice thickness, 71% had a too-large pixel size, 56% had a slice gap, and 48% did not use an IR-prep gradient echo. For FLAIR, only 18% satisfied the requirements. For the other data, 8% and 21% missed the axial and sagittal FLAIR acquisition, respectively, and 82% had too-large slice thickness, slice gap, or too-large pixel size. For the T2 sequence, only 7% satisfied the criteria. The most important reason for failing was a too-large pixel size (92% of unsatisfying images). 17% of the scans had no DWI, 1% of all scans had a good DWI, the other 81% had a too-large pixel size. If a post-gad T1 was provided, 29% satisfied the guidelines. Of all post-gad T1 sequences that did not comply, 70% had a too-large pixel size, mostly in combination with suboptimal slice thickness and/or slice gap. **Conclusions:** In a real-world MRI dataset of patients with MS, the conformance to the CMSC brain MRI guidelines was extremely low. The main reason was the use of a too-large pixel size, mostly in combination with a too-large slice thickness and a slice gap, which could be due to speeding up the protocol.

Supported by: None

Disclosure: *Sophie Vercrayssen, Arne Brys, Melissa Verheijen, Brandon Steach, Eline Van Vlierberghe, Diana M. Sima, Dirk Smeets: icometrix (salary).*

Keywords: Imaging and MS

(IMG02) Optic Nerve Head Volume Is Significantly Decreased in Pediatric Multiple Sclerosis and Not Pediatric Myelin Oligodendrocyte Glycoprotein (MOG)-Related Disorders

Giulia Longoni,¹ Tara Berenbaum,² Sunil K. Yadav,^{3,4} Ella M. Kadas,^{3,4} Michael J. Wan,⁵ Arun Reginald,⁵ Donald Mabbott,⁶ Alexander Brandt,³ E. Ann Yeh¹

¹Department of Paediatrics, Division of Neurology, The Hospital for Sick Children, Toronto, ON, Canada; ²Department of Neurosciences and Mental Health, The Hospital for Sick Children, Toronto, ON, Canada; ³Charité – Universitätsmedizin, Berlin, Germany; ⁴Nocturne GmbH, Berlin, Germany; ⁵Ophthalmology and Vision Sciences, The Hospital for Sick Children, Toronto, ON, Canada; ⁶Research Institute, Neuroscience and Mental Health Program, The Hospital for Sick Children, Toronto, ON, Canada

Background: Youth with multiple sclerosis (MS) or myelin oligodendrocyte glycoprotein (MOG)-associated disorders exhibit changes in retinal nerve fiber layer (RNFL) and ganglion cell inner plexiform layer (GCIPL) thickness, but knowledge about abnormalities in optic nerve head (ONH) morphology in these groups is lacking. **Objectives:** To compare ONH characteristics of youth with MS or MOG-associated disorders with a healthy control population. **Methods:** Standardized OCT data (Cirrus HD-OCT [Carl Zeiss Meditec, Dublin, CA]) were collected from healthy controls ($n = 27$ [16 female, median age 15 years (IQR 1.3)]), youth with MS ($n = 17$ [11 female, 16.9 years (1.5), disease duration 2.6 years (2.8)]), and youth with MOG-related disorders (MOG, $n = 13$ [11 female, 11.5 years (4.2), 1.7 years (1.3)]). ONH parameters (total ONH volume, Bruch membrane opening [BMO] region volume, and BMO minimal rim width) were computed by triangulated 3D surface reconstruction. Mean peripapillary RNFL and macular GCIPL were derived from manufacturer's fully-automated segmentation software, and results were manually reviewed where necessary. Multivariable mixed effects models were used to model the 5 ONH parameters between groups, accounting for age at OCT, eye-specific number of optic neuritis episodes, and a subject-specific random intercept. Results were Bonferroni adjusted for multiple comparisons ($P = .01$). **Results:** Participants with MS showed lower ONH total volume (-0.24 mm^3 , SE 0.073, $P = .002$), BMO region volume (-0.22 mm^3 , SE 0.083, $P = .012$), and BMO minimal rim width (-0.42 dMRW , SE 0.014, $P = .004$) compared to controls. ONH parameters were not different in MOG patients vs controls. Neither RNFL nor GCIPL were significantly different between controls and MS. GCIPL was lower in MOG than controls ($-7.9 \text{ } \mu\text{m}$, SE 3.05, $P = .012$). **Conclusions:** Our analysis showed ONH volume loss in pediatric MS despite no significant differences in RNFL or GCIPL in comparison to controls. These differences, except for a lower GCIPL, were not present in MOG vs controls. ONH analysis may therefore be superior to RNFL or GCIPL in detecting anterior visual pathway injury in MS early in the disease course. Larger studies are needed to confirm these findings.

Supported by: None

Disclosure: *Giulia Longoni, Tara Berenbaum, Michael J. Wan, Arun Reginald, Donald Mabbott, E. Ann Yeh: Nothing to disclose. Sunil K. Yadav, Ella M. Kadas: Nocturne GmbH (co-founder, shareholder). Alexander Brandt: Nocturne GmbH, Motognosis GmbH (co-founder, shareholder).*

Keywords: Imaging and MS

(IMG03) Cerebellar Connectivity Is Associated with Verbal Memory Impairment in Multiple Sclerosis

Mark D. Zuppichini,¹ Kathryn West,² Dinesh Sivakolundu,³ Darin T. Okuda,⁴ Bart Rypma³

¹School of Behavioral and Brain Sciences, University of Texas at Dallas, Richardson, TX; ²Center for BrainHealth, Dallas, TX; ³School of Behavioral and Brain Sciences, University of Texas at Dallas, Richardson, TX; ⁴UT Southwestern Medical Center, Dallas, TX

Background: Network connectivity is disrupted in multiple sclerosis (MS) and is related to cognitive function. Verbal memory impairment is common in MS yet the underlying neuropathology is still unknown. Recent research suggests cerebellar involvement in verbal memory and the cerebellum is adversely affected by MS. Thus, investigating the role cerebellar dysfunction plays in MS-related abnormal connectivity and verbal memory impairment may aid in understanding MS-related verbal memory impairment. **Objectives:** Investigate the relationship between cerebellar-cortical connectivity and verbal memory impairment in MS. **Methods:** 45 patients with MS and 23 healthy controls completed a verbal memory task (Selective Reminding Task [SRT]) and underwent magnetic resonance imaging. Resting-state (RS) functional connectivity analysis and diffusion kurtosis imaging were used to assess functional and structural connectivity, respectively. **Results:** The MS group performed significantly worse on SRT trial 1 ($t[66] = 2.18, P = .033$), SRT trial 3 ($t[66] = 2.20, P = .031$), SRT trial 6 ($t[66] = 2.44, P = .017$), and SRT Delayed Recall ($t[66] = 2.27, P = .026$). Resting-state analysis of cerebellar-cortical connectivity revealed significant differences between the cerebellum and several cortical areas. In MS, higher connectivity was observed between the cerebellum and superior frontal gyrus, precuneus, supramarginal gyrus, medial frontal gyrus, inferior parietal lobe, cingulate, and parahippocampal gyri ($P = .05$, FWE-corrected). Correlation analysis within the MS group revealed significant correlations between SRT Delayed Recall scores and connectivity values between the cerebellum and parahippocampal gyrus, fusiform gyrus, insula, cingulate, inferior frontal gyrus, uncus, middle temporal gyrus, and angular gyrus ($P = .05$, FWE-corrected). We observed higher connectivity in memory-impaired patients with MS between the cerebellum and the left parahippocampal gyrus compared to memory-preserved patients with MS. Diffusion analysis showed that axonal volume of the middle cerebellar peduncle significantly explained variability in SRT Delayed Recall scores in patients with MS over and above age and education ($F_{1,32} = 4.62, P = .039, R^2 = 0.17, \Delta R^2 = 0.12$). **Conclusions:** Abnormal resting-state and structural connectivity between the cerebellum and cortical areas, specifically the left parahippocampal gyrus, may contribute to verbal memory impairment observed in MS.

Supported by: None

Disclosure: *Mark D. Zuppichini, Kathryn West, Dinesh Sivakolundu, Bart Rypma:* Nothing to disclose. *Darin T. Okuda:* Acorda (speaking fees); Biogen (contracted research); EMD Serono, Genentech, Novartis, Teva Neuroscience (consulting fee); Genzyme (consulting fee, speaker fees).

Keywords: *Imaging and MS, Memory, Psychological issues and MS*

(IMG04) The Association Between Magnetic Resonance Imaging Brain Volumes and Computerized Cognitive Scores of People with Multiple Sclerosis

Daniel Golan,¹ Jared Srinivasan,² Olivia Kaczmarek,² Jonathan Bautista,² Myassar Zarif,² Barbara Bumstead,² Marijean Buhse,² Lori Fafard,² Jeffrey Wilken,³ Cynthia Sullivan,⁴ Timothy Fratto,⁵ Eline Van Vlierberghes,⁵ Diana Sima,⁶ Wim Van Hecke,⁶ Mark Gudesblatt²

¹Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel; ²South Shore Neurologic Associates, Patchogue, NY; ³Washington Neuropsychology Research Group, Washington, DC; ⁴Nursing, State University of New York at Stony Brook, Stony Brook, NY; ⁵icomatrix, Chicago, IL; ⁶icomatrix, Leuven, Belgium

Background: Cognitive impairment is common and disabling among people with multiple sclerosis (PwMS) but is not often monitored or only partially screened due to complexity of evaluation. Computerized cognitive assessment facilitates the incorporation of multidomain cognitive monitoring into routine clinical follow-up. **Objectives:** To explore the associations between brain volumes and cognitive scores of a computerized cognitive assessment battery (CAB, NeuroTrax) among PwMS. **Methods:** Participants were tested with the CAB and underwent brain magnetic resonance imaging (MRI) within specified time intervals. The global cognitive score (GCS) is the average of age- and education-adjusted scores of the various cognitive domains (memory, information processing speed, attention, executive function, motor and verbal). Whole brain volume (WBV), gray matter volume (GMV), white matter volume (WMV), thalamic volume, hippocampal volume, white matter lesion volume, and lateral ventricles volume were assessed by IcoMetric, a fully automated tissue and lesion segmentation and quantification software, that uses 3D T1-weighted and fluid-attenuated inversion recovery (FLAIR)

MRI. **Results:** 201 PwMS were tested with both CAB and MRI within 180 days (age: 52.3 ± 11.1 , 143 [71%] female). Significant correlations were found between the GCS and WBV, WMV, GMV, thalamic volume, and FLAIR lesion volume (Spearman rhos: 0.33, 0.3, 0.43, 0.4, -0.26, $P < .01$, respectively). Correlation coefficients remained significant but decreased as the time between MRI and CAB increased. The number of impaired cognitive domains was also associated with both lesion volume and GMV ($\rho = 0.25, -0.44; P < .05, <.01$, respectively). The only cognitive domain score that was associated with hippocampal volume was memory ($\rho = 0.27, P < .05$). **Conclusions:** Computerized cognitive scores are significantly associated with quantified MRI. These findings demonstrate the added information that can be derived from integrating digital assessment tools into the routine clinical assessment of PwMS.

Supported by: None

Disclosure: *Daniel Golan, Jared Srinivasan, Olivia Kaczmarek, Jonathan Bautista, Marijean Buhse, Lori Fafard, Timothy Fratto:* Nothing to disclose. *Myassar Zarif:* Acorda, Biogen, Genzyme, Teva (speakers' bureau). *Barbara Bumstead:* Biogen, Genzyme (speakers' bureau). *Jeffrey Wilken:* Biogen (contracted research); EMD Serono (speakers' bureau); Genzyme (contracted research, speakers' bureau). *Cynthia Sullivan:* Roche (contracted research). *Eline Van Vlierberghes, Diana Sima, Wim Van Hecke:* icometrix (salary). *Mark Gudesblatt:* Acorda, Amgen, Medtronic, Saol Therapeutics (speakers' bureau); Biogen, EMD Serono, Novartis, Sanofi, Teva (contracted research).

Keywords: *Equipment in MS, Imaging and MS, Natural history of MS*

(IMG05) Analysis of Demyelinating Injuries in Magnetic Resonance Imaging in People with Multiple Sclerosis

Dyana Fernandes Sr., Ana Canzonieri, Juliana Rhein, Liliana Russo

Associação Brasileira de Esclerose Múltipla, São Paulo/SP, Brazil

Background: Multiple sclerosis (MS) is a demyelinating autoimmune disease in which the immune system affects the myelin sheath of neurons, resulting in several clinical manifestations. Magnetic resonance imaging (MRI) is essential for understanding MS, as it allows objective visualization of both acute and chronic lesions. Signs and symptoms of MS depend on the location of the lesions and often affect people's quality of life. **Objectives:** To analyze MRI lesions in a group of individuals with MS. **Methods:** Ten individuals with MS participated. Retrospective MRI analyses were performed in the database over a period of 1 year with the Doctor Neurologist of the Civil Social Institution for MS, in the city of São Paulo, in 2019, where the individuals present MRI at the consultation. **Results:** Data from both sexes were used, being 60% women and 40% men, minimum age of 29 and maximum of 59. Of these individuals, 50% had the recurrent remitting clinical subtype, 30% secondary progressive, and 20% primary progressive; 40% had been diagnosed for over 10 years. Regarding Expanded Disability Status Scale score, most (70%) were between 4.5 and 6.5 and 50% had outbreaks for more than 3 years. The images revealed white matter lesions, T2 and fluid-attenuated inversion recovery hyperintensities, with contrast uptake in T1-weighted images, with predominantly juxtacortical involvement with 12%; 11% in the corpus callosum; followed by 8% in the temporal, periventricular regions; 6% in the cervical and thoracic regions; 4% in the subcortical, frontal, cerebellar, and parietal regions; and finally 2% in the bridge, bulb, and thalamic regions. The most frequent types of injuries are atrophy 12% and Black roller 4% with less incidence in the bridge and bulb. **Conclusions:** This paper suggests some areas that are most affected by the disease through MRI, and in our study population, the greatest involvement was brain and a small percentage was at the medullary level, generating motor, sensory, and cognitive impairment. Studies of this kind favor health professionals in understanding the evolution of the disease and the development of neurorehabilitation techniques. Further studies in this line should be conducted with a larger number of participants, so that we can have a better idea about the incidence of the affected areas.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Injuries, Imaging and MS*

MULTIDISCIPLINARY CARE

(MDC01) Social Assistance Intervention in Multiple Sclerosis

Alice Estevo Dias,¹ Priscila S. Santos²

¹Scientific Research and ²Social Service, Brazilian Association of Multiple Sclerosis, Sao Paulo, Brazil

Background: People with multiple sclerosis (PwMS) have complex symptoms and different types of needs. These demands include how to manage the burden of physical disability as well as how to organize daily life and restructure social roles in the family and at work. **Objectives:** To identify difficulties and obstacles experienced by PwMS, highlighting the work of social service in promoting physical, psychological, and social well-being. **Methods:** The sample involved 113 PwMS, 82 women and 29 men, aged 17 to 77 years (mean [SD] = 41 [11.38]). All answered the semistructured sociodemographic questionnaire containing 30 questions, developed specifically for this population. **Results:** Despite the highlighted needs, lack of knowledge about treatment, rehabilitation, and maintenance of quality of life was widely identified among the participants. In this sense, specific referrals and specializes guidance pertinent to our country were carried out as follows: n = 117/100% health (high-cost medicines), n = 35/30% social security (retirement), n = 23/20% education (educational institutions), n = 59/50% judicialization (medicines demand in lawsuits). **Conclusions:** The individual reception procedure was necessary and sensitive for understanding the demands through the questionnaire. The difficulties identified in this study determined social assistance actions directed to the development of joint actions with multidisciplinary and interdisciplinary teams, which directly affected the quality of life of patients with all types of MS.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Management of activities of daily living in MS, MS and the caregiver/family

(MDC02) Is a 2-Week Intensive Day Program an Effective Approach to Provide Outpatient Services for People with Multiple Sclerosis?

Clare T. Hartigan,¹ Tracy Walker²

¹Multiple Sclerosis Outpatient Rehabilitation and ²Virginia C. Crawford Research Institute, Shepherd Center, Atlanta, GA

Background: A unique 1-week intensive MS day program rehabilitation model was initiated in 2012 and continues to evolve. The goal of the program is to educate and provide patients, families, and caregivers with a structured plan to improve overall health and wellness. Our interdisciplinary team would like to share "lessons learned the hard way" over the past 7 years and to report the most recent patient outcomes. **Objectives:** 1) Identify key factors that may indicate appropriateness for patient enrollment in an intensive day program (DP). 2) Identify "lessons learned" over the past 7 years. 3) Present results of objective patient outcome measures.

Methods: Patients are evaluated to determine appropriateness for participation in a 2-week intensive DP. To qualify for DP, they must require skilled services for physical therapy (PT), occupational therapy (OT), speech therapy (ST), and wellness. Additionally, they may also receive nursing, counseling, and vocational rehabilitation as appropriate. It is mandatory that a caregiver/family member be present for all sessions. Pre and post outcome measures used to assess change include Modified Fatigue Impact Scale (MFIS), Fatigue Severity Scale (FSS), Symbol Digit Modalities Test (SDMT), Nine-Hole Peg Test (NHPT), and Timed 25 Foot Walk (T25ft). **Results:** Expanded Disability Status Scale (EDSS) scores ranged from 2.5 to 9.0 with 67% of patients having EDSS score 5.0 or greater. Patient and family feedback regarding DP has generally been positive. The most frequent negative comments are that too much information is presented and becomes overwhelming, not enough rest breaks are given, and the days are too long. Despite reports of being too intensive, a large majority of participants have shown progress on outcome measures. Percentage of patients whose scores improved are as follows: MFIS 70%, FSS 90%, SDMT 70%, NHPT 67% (right) 87% (left), and T25ft 75% (note 3 patients went from being nonambulatory to walking with a rollator). A more detailed report of the data will be ready at the time of presentation. **Conclusions:** Multiple factors need to be considered prior to recommending DP to persons with MS. To achieve success, patients/families must be willing to modify old behaviors and staff must adapt session intensity so as not to overwhelm or overfatigue patients. An interdisciplinary community outing at the end of the first week has proven valuable to patients/families. Changes to the structure of the DP have been incorporated based on patient feedback. Re-evaluations 2 months post-DP are encouraged.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Day program outcomes

(MDC03) The Waiting Room: A Successful Experience in the Multiple Sclerosis Care and Treatment Center (CATEM)

Ivone R. Fernandes, Ana C. Sena, Maria L. Santos, Rodrigo N. Fonseca, Vanessa D. Vale, Natali M. Marques, Elisa Melo, Charles P. Tilbery

Santa Casa de Misericórdia, São Paulo, Brazil

Background: The constant waiting of patients for the call for medical consultation is a reality in the Brazilian public service. The National Policy for Humanization (NPH) of the Brazilian Ministry of Health (MH) advocates providing welcoming, strengthening, and wholesome care, with the adoption of measures and communication among multiprofessional teams. One of the practices used by the health team to share experiences, feelings, and knowledge between patients of professionals is the Waiting Room Group. The Multiple Sclerosis Care and Treatment Center (MSCTC), established in 1997 at the Neurology Clinic of Santa Casa de Sao Paulo, attends through the Unified Health System (UHS) every Friday morning and has about 500 registered patients who make use of several therapies for treatment of multiple sclerosis (MS). To minimize the waiting period, the "Waiting Room" project was created in 2013, consisting of patients, family members, caregivers, social worker, nurse, physiotherapist, psychologist, and neuropsychologist, among others, to provide a welcoming space to minimize anxiety and fear, advise on their rights, inform about the disease and its complications, types of treatment, importance of adoption and adherence of exercise routines, and hospital policy and routines, and providing patients, family members, and caregivers with a space of power and affection in order to have a more active and participative role in their treatment. **Objectives:** To describe the experience of the Waiting Room Group as part of humanized care for patients with MS, their family members, and caregivers. **Methods:** Welcoming, integration, and interaction are the key words: the earlier patients talk about and share their experiences with those who are starting their treatment, their anxiety and doubts are minimized. The doctor, psychologist, social worker, nurse, and physiotherapist participate in all meetings, as well as other invited professionals. During the meetings, the doubts presented by the participants are clarified and provide topics for discussion in the next meetings. **Results:** Not applicable. **Conclusions:** This project has been developed for 6 years and has been successful, with an average participation of 500 patients per year.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Interdisciplinary team, MS and the caregiver/family

(MDC05) Implementation of a Pharmacist-Led Immunization Program in a Center for Comprehensive Multiple Sclerosis (MS) Care

Lauren Long, Ryan Fuller

Neurology, Hospital of the University of Pennsylvania, Philadelphia, PA

Background: In 2019, the American Academy of Neurology issued a practice guideline update for vaccinations in patients with multiple sclerosis (MS). In this guideline, they recommend providers "assess and reassess vaccination status of patients with MS before prescribing immunosuppressive or immunomodulating (ISIM) therapy and should vaccinate these patients." Also in 2019, eculizumab was the first US Food and Drug Administration (FDA) treatment approved for neuromyelitis optica spectrum disorder (NMOSD). Prior to starting, the FDA requires patients receive 2 types of meningococcal vaccines (MenACWY plus MenB-4C or MenB-FHbp) at least 2 weeks prior to starting eculizumab. MS centers need to create an efficient process to ensure patients receive vaccines in a timely fashion to reduce harm from vaccine-preventable diseases and not delay start of patient's disease-modifying therapy (DMT). In 2002, Pennsylvania allowed pharmacists to provide immunizations to patients under a collaborative practice agreement or a direct order from a provider. At the Hospital of the University of Pennsylvania, there are 2 clinical pharmacists dedicated full-time to the MS Center. Little information in the literature addresses providing immunizations to patients with MS and prior to starting DMTs, and the utilization of MS clinical pharmacists and a health-system based specialty pharmacy in providing this unmet need. **Objectives:** The purpose of this quantitative pilot study will be to review the results of a pharmacist-led immunization program imbedded within a 16-provider MS clinic. **Methods:** Patients will be identified by provider referrals and pharmacist comprehensive chart reviews of newly diagnosed patients and patients starting ISIM and eculizumab therapy. **Results:** Data will be analyzed with descriptive statistics. **Conclusions:** It is anticipated this pilot study will increase rates of vaccination of patients with MS, reduce time to start DMT, and increase awareness of vaccine-preventable diseases.

Supported by: None

Disclosure: *Lauren Long:* Celgene, Novartis (consulting fee). *Ryan Fuller:* Nothing to disclose.

Keywords: *Comprehensive care and MS, Disease-modifying treatments in MS, Preventative health*

(MDC06) The African American Experience and Multiple Sclerosis

Marie LeGrand

Multiple Sclerosis Association of America, Chicago, IL

Background: Incidence rates of multiple sclerosis (MS) have been found to be higher in both black males and black females vs white males and white females. In particular, black females have a 47% higher risk for MS when compared to white females. Disease progression is significantly faster in black patients with MS in both brain and retinal measures. Magnetic resonance imaging scans show whole brain and gray and white matter atrophy to occur twice as fast in African Americans compared to Caucasian Americans. African American patients also show quicker atrophy of the thalamus, a possible link to cognitive impairment. Furthermore, African Americans are dying from MS at an earlier age, suggesting that MS burden weighs disproportionately across race demographics. To fully understand, the Multiple Sclerosis Association of America (MSAA) launched the African American MS Advisory Board. **Objectives:** The African American Experience & Multiple Sclerosis initiative included 11 MS clinicians and 16 African Americans affected by MS. The meeting's objectives were to 1) create a dialogue allowing both groups to share their views on the problems that African Americans with MS are facing; 2) evaluate programmatic initiatives that address the unmet needs; 3) gather from the meeting to aid in the development of an actionable plan, tailored educational offerings, and services provided; and 4) identify next steps to continue building on the work of the African American committees. **Methods:** Participants attended a half-day meeting, sharing their views on problems African Americans with MS are facing. Sessions included moderated discussions and a brief presentation, emphasizing the need for research and the importance in developing programmatic initiatives. **Results:** Key findings from the meeting in Atlanta elucidated the aforementioned performance gaps experienced by MS clinicians responsible for treating African Americans with MS. Characteristics of patients with MS less likely to see neurologists include the following: 1) lack of health insurance, 2) lower income, 3) African American, 4) living in rural areas, and 5) illness longer than 15 years. Conversely, patients cared for by a neurologist are more likely to 1) undergo diagnostic tests, 2) undergo treatment-related tests, 3) be treated with disease-modifying therapies, 4) receive medication for symptoms, and 5) report their providers had a treatment plan. The presence of these gaps requires behavioral change on the behalf of MS clinicians that will appropriately address barriers resulting in initial treatment delays, and a much faster disease progression in black patients with MS. **Conclusions:** Data from this comprehensive initiative will drive the materials and information for a set of multifaceted interventions to improve the knowledge, competence, and/or performance of MS clinicians who are currently treating, or have the potential to treat, black patients with MS. The anticipated result is to better identify and detect early signs of disease progression and timely therapeutic intervention.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *African American and MS, Comprehensive care and MS*

(MDC07) The Impact of the Nurse Practitioner Model of Care within Multiple Sclerosis

Meaghan Osborne, Suyinn Chong

Neurology, Royal Brisbane and Women's Hospital, Brisbane, Australia

Background: The nurse practitioner (NP) within Australia and within multiple sclerosis (MS) is designed to improve outcomes for people with MS through the adoption of strategies that optimize therapeutic decision-making and improve communication around benefit and risk. The NP model is a safe and effective use of resources that enhances patient-related outcomes. **Objectives:** There is much evidence in the literature to support the NP internationally, however within Australia the model has been limited with numerous barriers to fulfill the role to its full potential. The objective of this data set is to compare and contrast the effectiveness and safety of the NP role in the MS setting. **Methods:** A cohort analysis of patients with a primary diagnosis of MS was conducted over a period of 12 months. This analysis assessed a diverse range of end points; including time to treatment, shared decision-making practices, benefit and risk discussion, and access to care. A comparison was made between the

NP and specialist neurologists in a large neurology department of a quaternary hospital within Brisbane, Queensland, Australia. A simple random sampling process was used to generate 168 patients from each group. Further statistical analysis will be completed upon this data set. **Results:** Of 168 patients, a greater proportion were seen within 30 days of referral to an NP (59%) contrasted with a neurologist (11%). It was found that of patients who saw an NP vs a neurologist, there was increased discussion of potential benefit and risk (40% and 11%), increased initiation of disease-modifying therapy (50% compared with 21%), more frequent identification of relapses (23% vs 4%), and a greater appointment attendance in the NP cohort (99% and 82%). When examining the access to treatment between the 2 cohorts, it was found that 75% of the NP cohort received treatment within 0-60 days, as opposed to 65% of the neurologist cohort. Both cohorts were comparable in terms of hospital readmissions and emergency department presentations within 90 days of clinician consultation. **Conclusions:** The results highlight the benefit of the NP role within a neurology practice in an Australian setting; it was evident that the NP provided timely access to care and treatment. NPs remain a safe, effective, and valuable role within the MS community. Further resources as well as research should be implemented to support and evaluate this role within hospitals and communities across Australia.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Comprehensive care and MS, Disease-modifying treatments in MS, Nursing management in MS*

(MDC08) Late-Onset Multiple Sclerosis: Comorbidity and Disease Progression

Smathorn Thakolwiboon,¹ Mirla Avila,¹ Pavidia Pachariyanon,² Jie Pan,¹ Amputch Karukote,¹ Gyeongmo Sohn¹

¹Neurology and ²Internal Medicine, Texas Tech University Health Sciences Center, Lubbock, TX

Background: Late-onset multiple sclerosis (LOMS) defined as multiple sclerosis (MS) with clinical onset after the age of 50. Previous studies have demonstrated that late onset is a poor prognostic factor for MS. Moreover, several comorbidities such as hypertension, diabetes, and dyslipidemia were reported in association with poor clinical outcome in patients with MS. Although the prevalence of comorbidity is increased in the aging population, the roles of comorbidity in LOMS has not been explored. **Objectives:** To evaluate the influence of comorbidities in LOMS. **Methods:** This retrospective study included 38 patients with MS with clinical onset after the age of 50. Demographic, clinical, radiologic, and laboratory findings were collected. The survival analysis was performed to identify the comorbidities that associated with losing of no evidence of disease activity (NEDA) status. Relapse, Expanded Disability Status Scale worsening, or new T2 or T1 gadolinium-enhancing lesions on magnetic resonance imaging resulted in loss of NEDA status. **Results:** The median follow-up was 26 (IQR 12-45.75) months. Forty-five percent of participants remained on NEDA status until the last follow-up. Hypertension (HTN) significantly increased the risk of disease progression (HR 4.36, 95% CI 1.39-13.66, $P = .012$). Moreover, the risk of losing NEDA in patients with poor-controlled HTN is higher than in patients with well-controlled HTN (HR 7.67, 95% CI 1.01-58.12, $P = .049$). Interestingly, there was no significant difference in disease progression risk between well-controlled HTN and normotensive patients with LOMS (HR 1.58, 95% CI 0.28-9.01, $P = .608$). Diabetes, dyslipidemia, coronary artery disease, stroke, obesity, hypothyroidism, depression, and anxiety were not significantly associated with MS progression in the late-onset population. **Conclusions:** Our cohort study suggests that HTN is a modifiable risk factor of disease progression in LOMS. Previous studies have demonstrated that HTN can potentiate neurodegenerative process. Larger prospective studies are needed to further explore the interaction between HTN and disease-modifying therapy and the effect of antihypertensive agents.

Supported by: None

Disclosure: *Smathorn Thakolwiboon, Pavidia Pachariyanon, Jie Pan, Amputch Karukote, Gyeongmo Sohn:* Nothing to disclose. *Mirla Avila:* Biogen, Genentech, Genzyme (education); Celgene (education).

Keywords: *Comorbidity, Comprehensive care and MS*

(MDC09) Comparing Patient Perceptions on Multiple Sclerosis Management and Care: A Subanalysis of Geographic Differences

Sarah A. Morrow,^{1,2} Alexey Boyko,³ Heidi Thompson,⁴ Maija Pontaga,⁵ Nektaria Alexandri⁶

¹Western University, London, ON, Canada; ²Clinical Neurological Sciences, London Health

Sciences Centre, London, ON, Canada; ³Pirogov Russian National Research Medical University, Moscow, Russia; ⁴Neurology, Southern Health & Social Care Trust, Portadown, United Kingdom; ⁵MS in the 21st Century, Riga, Latvia; ⁶Neurology and Immunology, Merck KGaA, Darmstadt, Germany

Background: The MS in the 21st Century initiative is led by a steering group of international multiple sclerosis (MS) specialists and patient advocates with a current focus of improving education and communication between health care professionals and people with MS (PwMS). **Objectives:** To compare the perceptions of PwMS on MS management and care across 2 geographical regions: Europe and North America (United States and Canada). Particular emphasis was on patient support at diagnosis, treatment decisions, and communication. **Methods:** An electronic survey was developed to gain insight into patients' opinions on unmet needs in MS management. The surveys were conducted at 2 international patient meetings in 2017 and 2018. Multiple answers were solicited in response to 10 questions. **Results:** A total of 55 PwMS from Europe and 46 from North America completed the survey. PwMS in Europe listed a lack of time in medical appointments as their biggest challenge at diagnosis (60.0%), whereas PwMS in North America reported understanding disease progression to be their biggest challenge (57.8%). European PwMS reported greater levels of additional patient support available in their clinics (ie, MS nurse [67.3%], information about employment [21.2%], or psychological support [25.0%]), whereas 26.1% of North American PwMS reported no additional support. PwMS in Europe reported being less involved in treatment decisions, with 20.5% saying they were not involved, compared with 2.4% in North America. European PwMS placed more importance on the safety of their treatment (57.7%), whereas PwMS in North America placed more importance on the efficacy of their treatment (71.7%). **Conclusions:** There were distinct geographical variations between PwMS's perceptions and priorities relating to MS care, education, and treatment decisions. European PwMS reported less time in appointments and lower involvement in treatment decisions than North American PwMS, however they also reported greater levels of support and education outside of their neurologist appointments including greater access to specialist MS nurses.

Supported by: None

Disclosure: Sarah A. Morrow: Biogen, Novartis (contracted research); Celgene (consulting fee); EMD Serono, Roche, Sanofi Genzyme (speakers' bureau). Alexey Boyko: Biogen, Schering, Merck, Teva, Novartis, Sanofi Genzyme, Actelion, Biocad, Generium (consulting fee, participated in clinical trials supported by). Heidi Thompson: Biogen, Merck KGaA (consulting fee). Maija Pontaga: Merck KGaA (consulting fee). Nektaria Alexandri: Merck KGaA (salary).

Keywords: Comprehensive care and MS, Shared decision making

METHODS OF CARE

(MOC01) Understanding the Health Care Provider-Patient Relationship in Treating Multiple Sclerosis

Beth Schneider

MyHealthTeams, San Francisco, CA

Background: Health care providers (HCPs) play a critical role in treating patients with multiple sclerosis (MS), especially in helping patients get on treatment to help slow progression. An online survey of patients with MS as well as natural language analysis of organic interactions on a social network provided insight into the relationship between HCPs and their patients with MS. Understanding the HCP-patient dynamic is crucial to improving these interactions as well as identifying tools and educational materials to help patients better manage MS. **Objectives:** Leverage MS patient social network to understand the HCP-patient dynamics, including what is working and any opportunities to improve HCP-patient interactions. **Methods:** A 2-pronged approach was undertaken. First, an online survey was completed by 658 US members of MyMSTeam in November 2018. Research was also conducted on de-identified organic discussions within MyMSTeam.com, a social network of >127,000 people diagnosed with MS (approximately 1:9 patients with MS in United States). A natural language processing tool (NLP) analyzed 178,884 verbatim April to September 2019. Survey results showed 71% of patients indicated they rely on HCP for information about disease-modifying treatments (DMTs), 69% indicated HCP's recommendation to get on a specific DMT is most important factor, and 20% indicated decision to get on a DMT would be easier had HCP provided stronger opinion rather than relying on the patient to research and decide. **Results:** Analysis of doctor-patient relationship found 15% of conversations on MyMSTeam, n = 37,978 discussions focused on HCP-patient relationship. 57% of conversations were generally negative in tone. NLP analysis revealed the

dichotomy of the HCP-patient relationship. Sentiment was very positive when HCP addressed MS holistically as well as when patient felt HCP was truly listening to their needs and concerns. Conversations negative in sentiment were often the result of feeling rushed, not being listened to, or when the patient felt only MS progression was being addressed by HCP and not other symptoms also experienced. **Conclusions:** Understanding the needs of patients with MS provides significant opportunities for HCPs to better support and educate their patients, including through consultation that includes a stronger recommendation on treatment path as well as listening to patient concerns and addressing the sum total impact of MS including pain, depression, and fatigue and not just disease progression.

Supported by: None

Disclosure: Beth Schneider: MyHealthTeams (contracted research).

Keywords: Comprehensive care and MS, Management of activities of daily living in MS, Psychological issues and MS

(MOC02) Cancelling Clinic Appointments: What Factors Are Associated with Higher Rates of Cancellations in Patients with Multiple Sclerosis?

Elizabeth S. Gromisch,^{1,2,3,4} Aaron P. Turner,^{5,6,7} Steven L. Leipertz,⁷ John Beauvais,^{2,8} Jodie K. Haselkorn^{5,6,7,9}

¹Mandell Center for Multiple Sclerosis, Mount Sinai Rehabilitation Hospital, Trinity Health Of New England, Hartford, CT; ²Psychology Service, VA Connecticut Healthcare System, West Haven, CT; ³Department of Neurology, University of Connecticut School of Medicine, Farmington, CT; ⁴Departments of Rehabilitative Medicine and Medical Sciences, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT; ⁵Department of Rehabilitation Medicine, University of Washington, Seattle, WA; ⁶Rehabilitation Care Service, VA Puget Sound Health Care System, Seattle, WA; ⁷Multiple Sclerosis Center of Excellence West, Veterans Affairs, Seattle, WA; ⁸Department of Psychiatry, Yale University School of Medicine, New Haven, CT; ⁹Department of Epidemiology, University of Washington, Seattle, WA

Background: It is important that persons with multiple sclerosis (MS) attend their scheduled appointments to maintain continuity of care and promote successful self-management. A recent study examined missed appointments in veterans with MS and developed a model that included 7 predictor variables (suboptimal disease-modifying therapy [DMT] adherence (>80%), emergency visits, age, distance, and histories of post-traumatic stress disorder, chronic obstructive pulmonary disease, and congestive heart failure). However, to date, there is limited information on appointment cancellations, which is a different appointment attendance behavior but can also disrupt care. **Objectives:** 1) To identify the rate of cancelled appointments in a large national sample of persons with MS and 2) to examine the demographic and clinical factors associated with high levels of cancelled appointments (defined as $\geq 50^{\text{th}}$ percentile). **Methods:** Administrative data between January 1, 2013, and December 31, 2015, were extracted from the VA MS Center of Excellence Data Repository, an electronic health record-based dataset composed of US veterans receiving services at any Veterans Affairs (VA) medical center. The cancellation rate was calculated by dividing the number of cancelled appointments (excluding "no shows") by the total number of scheduled appointments during this 2-year timeframe. Bivariate analyses were conducted to examine demographic and clinical characteristic differences between individuals with and without high rates of cancellations, and variables with a *P* value of <.10 were entered into a logistic regression. **Results:** Over 96% (n = 3623) had at least 1 cancelled appointment, with a median cancellation rate of 25%. Flags for high rates of cancellations included 1 or more inpatient hospitalization (odds ratio [OR]: 1.78), wheelchair issuance (OR: 1.48), distance (≥ 24.1 miles away; OR: 1.47), gender (male; OR: 1.28), suboptimal DMT adherence (OR: 1.26), and history of a mood disorder (OR: 1.28). **Conclusions:** Cancelled appointments are prevalent among veterans with MS. The similarities and differences in the variables included in the cancelled and missed appointment models highlight both malleable and nonmalleable factors associated with each type of appointment attendance behavior. While further information is needed to elucidate the reasons behind these cancelled appointments, these results may help clinicians identify individuals at risk for higher rates of cancellations and to plan targeted interventions.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Appointment attendance, Comprehensive care and MS

(MOC03) From Therapy Enrollment to First Dose: A Quality Improvement Initiative for Multiple Sclerosis Care

Jamie Bolling,¹ Ryan McNiff,² Tirisham V. Gyang,¹ Carlos Vervloet Solleró,¹ Aaron Carlson¹

¹Neurology, University of Florida, Gainesville, FL; ²UF Health, University of Florida, Gainesville, FL

Background: Timely treatment is important for reducing relapses and risk of disability in people with multiple sclerosis (MS). However, disease-modifying medications entail a complex enrollment process that can delay treatment initiation. **Objectives:** This pilot study tracked individuals initiating the enrollment process for ocrelizumab and natalizumab at the University of Florida (UF) MS Clinic. **Methods:** This quality improvement initiative captured all relevant documents, including enrollment forms, insurance communications, referrals, and other documentation. We monitored the dates of completion, signature, fax, and local medical record upload for all forms. **Results:** Preliminary data from January 1, 2019, to October 1, 2019, captured 19 patients enrolled in either medication. Of these 19 patients, 6 received treatment as of October 2019. On average, enrollment submission to treatment initiation took 56 (range, 5-135) days. Of the remaining 13 awaiting treatment, the average interval from enrollment to October 2019 was 83 (range, 4-175) days. Overall, 11 quantifiable delays were identified, resulting in 17 recorded contacts. Of the 11 delays, 45.5% were insurance related, 36.4% were clerical delays, and 18.1% were patient related. Delays took an average 24.6 days to resolve (30.4, 9.25, and 41 days, respectively). For the 17-recorded contacts, 41% were insurance-related, 47% were clerical errors, and 12% were related to patient compliance. Most delays occurred in patients referred to outside infusion centers for treatment. Five of 13 ocrelizumab patients initiated treatment as of October 2019. Four of the 5 patients were infused at UF. The 8 patients awaiting treatment were referred to outside infusion centers and averaged 101 days without treatment as of October 2019. Two of 4 natalizumab patients have received treatment, both infused at UF. For the 2 awaiting treatment, both were referred to outside infusion centers and have waited an average of 71 days for treatment as of October 2019. **Conclusions:** Preliminary analysis suggest that insurance-related delays were the largest barrier to treatment initiation. Referring patients to outside centers further impeded the process. The results indicate that revisions to standard operating procedures for insurance inquiries and referrals may be beneficial. Moreover, while clerical errors were common, they were quickly resolved. Patient compliance issues, though rare, had the most enduring effect on treatment initiation.

Supported by: None

Disclosure: *Jamie Bolling, Ryan McNiff, Tirisham V. Gyang, Carlos Vervloet Soller:* Nothing to disclose. *Aaron Carlson:* Novartis Pharmaceutical (contracted research); Sanofi Genzyme (consulting fee).

Keywords: Comprehensive care and MS, Quality improvement

(MOC04) Nurse Telephone Encounters in a Multiple Sclerosis Clinic in 2020

Janice Lake, Colleen Harris, Sharon Peters, Jackie Gaythorpe

University of Calgary MS Clinic, Calgary, AB, Canada

Background: With the growing treatment landscape in multiple sclerosis (MS), nurses are being challenged with a constantly increasing work load, and more of their time is being spent on telephone encounters. Previous data collected from January to April 2001 at the University of Calgary MS Clinic showed that 50% of nursing time was devoted to telephone encounters, and 30% of those encounters dealt with issues around disease-modifying therapies (DMTs). In 2001 there were 3 Health Canada-approved DMT options as compared to 12 approved DMTs at this time in 2020. Increased patient therapy choice has required more nursing time to educate patients on treatment expectations, potential adverse effects, and adherence to more complex medication protocols. There is a need to understand the type, frequency, and time spent on telephone encounters, which will assist MS Nurses to develop efficient and effective management protocols. **Objectives:** To determine the frequency, type, and duration of nurse telephone encounters. To compare work previously done in the MS Clinic by Harris et al in 2001. **Methods:** Patient telephone encounters will be analyzed from January 2, 2020, to the end of March 2020. A telephone call log designed based on previous workload analysis at the University of Calgary MS Clinic will be used to capture the type, duration, and frequency of calls. **Results:** Telephone encounters are ongoing, with data analysis completed April 1, 2020. **Conclusions:** Analyzing telephone encounters will provide information to assist with the development of work load management strategies.

Supported by: None

Disclosure: *Janice Lake:* Sanofi Genzyme (consulting fee). *Colleen Harris:* Biogen, Merck Serono, Novartis, Roche, Sanofi Genzyme (consulting fee). *Sharon Peters, Jackie Gaythorpe:* Nothing to disclose.

Keywords: Comprehensive care and MS

(MOC05) Pioneering Multiple Sclerosis Center Program with MSHA Certification to Improve Patient Care and Experience

Jennifer R. Hillner, Jennifer L. Rusk

Neurology/MS, OSF HealthCare Illinois Neurological Institute, Peoria, IL

Background: The OSF HealthCare Illinois Neurological Institute's Multiple Sclerosis Center is a Certified Comprehensive MS Center located in Peoria, Illinois, serving over 2100 patients and is focused on providing excellence in multiple sclerosis (MS) care for patients and families in need through diagnosis, treatment, education, and research. Like many MS Centers we are fortunate to have physicians, nurses, physical therapists, occupational therapists, and a registered dietician who are specialized in MS. However, we came to realize there were opportunities in our level of patient care and experience with the potential to be transformed if we invested in our medical office assistants and our MS patient navigators through MSHA Certification. Certified employees are able to deliver a higher level of care, better connect patients to resources, and also better support our comprehensive care team, which allows our center to ensure all members are working at the height of their licensure. We set a goal in May 2019 to have a mission partner in every job role of our MS Center to be certified in MS. We propose that achieving this level of certification in each and every MS Center around the world is a goal worth aspiring to in our combined fight against MS. **Objectives:** MSHA certification within our medical office assistant and MS patient navigator job roles in 2019. In addition, lower risks associated with delay of treatment, removing barriers to care, increasing clinical competency, and employee efficiency in tasks such as prior authorizations for disease-modifying therapies. **Methods:** Investing in team study and elevating employees through MSHA certification. We also brought our team to Consortium of Multiple Sclerosis Centers (CMSC) for education, networking, and patient resource opportunities. **Results:** The OSF HealthCare Illinois Neurological Institute's MS Center has an employee in every job role who is MS Certified. MSHA certification has given our team members greater context and compassion through a deeper understanding of MS at a professional level. We increased efficiency and decreased delay in care and also the time it took to obtain prior authorizations. Our certified employees are able to communicate with an increased level of empathy and confidence with patients. MSHA certification has fostered career dedication and driven passion while motivating our MS care team as a whole. **Conclusions:** MS is a complex and lifelong neurologic disease that requires all individuals involved with patient care to have a basic level of knowledge of the disease. The MSHA certification of our employees is a pioneering strategy that has elevated the level of care and experience we provide to our patients with MS by empowering our employees through the education and understanding of MS. MSHA certification is an effective tool for efficient and empathetic health care delivery to those individuals living with MS.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, MSHA certification

(MOC06) Conceptualizing Access Through the Perspectives of Canadians with Multiple Sclerosis

Julie Petrin,¹ Mary Ann McCall,¹ Catherine Donnelly,¹ Marcia Finlayson²

¹Rehabilitation Sciences, Queen's University, Kingston, ON, Canada; ²School of Rehabilitation Therapy, Queen's University, Kingston, ON, Canada

Background: Access to health care is vital to the health and well-being of people with chronic conditions like multiple sclerosis (MS). Access is often measured using service utilization as a proxy. Utilization measures may fail to capture the complexities of the experiences of accessing care that populations with chronic illness face. The Candidacy Framework offers an alternative to utilization measures, by examining the dynamic process a patient must engage in, with the health care system and all components of it, to negotiate their eligibility for care, which is described as one's candidacy. The process of accessing care is examined from the perspective of vulnerable populations and considers the impact of social patterning and health system environments on this process. **Objectives:** To investigate access to health care for the management of MS, in a Canadian context, from the perspective of persons with MS. **Methods:** The study design was informed by an interpretive descriptive methodology. Forty-eight individuals with MS living across 7 communities in Ontario were recruited primarily through the MS Society of Canada to participate in 1 of 5 focus groups or 10 individual telephone interviews. The sessions were digitally recorded and transcribed. The transcriptions were then analyzed using constant comparative methods. Additionally, the data was re-interpreted in light of the Candidacy Framework to determine its align-

ment to this framework. **Results:** All dimensions of the Candidacy Framework were relevant to the experiences of persons with MS. However, the framework failed to account for important aspects of the participants' lifelong experience of accessing care. Importantly, participants discussed the process of engaging in help-seeking which was informed by past experiences of seeking and receiving health care services, as well as the accumulated knowledge of living with MS. The most commonly reported past experiences were those regarding instances of patient-centered care, where negative experiences were described when this approach to care was not taken. Past negative experiences oftentimes made persons with MS hesitant to seek further care. **Conclusions:** The Candidacy Framework alone does not account for the lifelong interaction with health care that individuals with chronic illnesses such as MS face. To capture the full experience of access the framework should consider 3 main extensions: 1) The addition of recursivity, which captures the reciprocal nature of interacting with the health care system; 2) inclusion of help-seeking behavior and related decision making; and 3) inclusion of the concepts of patient-centered care.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Access to health care, Comprehensive care and MS

(MOC07) Access to Health Care for Canadians with Multiple Sclerosis: Prioritizing Concerns

Julie Petrin,¹ Mary Ann McColl,² Simon French,³ Catherine Donnelly,² Marcia Finlayson¹

¹School of Rehabilitation Therapy, Queen's University, Kingston, ON, Canada;

²Rehabilitation Sciences, Queen's University, Kingston, ON, Canada; ³Chiropractic, MacQuarie University, Sydney, Australia

Background: Current Canadian literature demonstrates that persons with multiple sclerosis (MS) are high users of health care services, yet still have multiple unmet needs and low satisfaction with health care services. International studies showing similar results suggest this may be related to issues in access to health care. **Objectives:** We aimed to 1) describe the health care service use of Canadians with MS in managing their condition; 2) identify the most pressing concerns Canadians with MS have in relation to accessing care to manage their MS. **Methods:** The aims were addressed using an online cross-sectional survey guided by Concerns Report Methodology. Inclusion criteria were: older than 18 years, a Canadian citizen, and self-reported diagnosis of MS. Data were collected about health care service use and the importance and satisfaction with access to health care service in the community. Data were analyzed using descriptive statistics. Access concerns were prioritized by calculating a Needs Index (NI). **Results:** To date, 211 persons with MS have completed the survey. Participants were predominantly female (86%), with a mean (SD) age of 46.4 (11.7), living with relapsing remitting MS (77%) for a mean (SD) of 10.4 (8.9) years. Just over half of the participants were still working (57%), with varying levels of disability ranging from 0-7 on the Patient-Determined Disease Steps (median: 2). Preliminary findings indicate that nearly all participants had a regular neurologist (97%), many of whom practice in an MS clinic (83%). Most individuals also had a general practitioner (96%). Participants on average visited their neurologist 1.5 times a year (SD: 0.7) and their general practitioner 4.6 times (SD: 2.3) a year. The highest concerns among participants regarding their access to care were: 1) affordability of complementary care (example: massage therapy, yoga, naturopathic care) (NI: 33.8) and physiotherapy and occupational therapy (NI: 29.7), both aimed at improving wellness; 2) availability of health care providers with MS-related knowledge in their communities to guide their care plan (NI: 33.7); and 3) communication between health care providers to ensure coordination of care (NI: 29.2). **Conclusions:** Preliminary findings suggest that for persons with MS, merely having regular neurologists and general practitioners is not considered satisfactory access to care. Persons with MS identified concerns regarding the availability of affordable health care services aimed at maintaining wellness. They also had concerns regarding the availability of community providers with sufficient MS-related knowledge to guide their referrals and care plans. Targeting policy reform promoting the coverage of health care services aimed at preventative and maintenance care may be a critical step in improving care for this population.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Access to health care, Comprehensive care and MS

(MOC08) Use of a Clinical Decision Support Tool to Support Monitoring and Care of Patients with Multiple Sclerosis Receiving Disease-Modifying Therapy

Lisa M. Mitchell,¹ Mitchell T. Wallin,² Natasha M. Antonovich³

¹VA Maryland Health Care System, Multiple Sclerosis Center of Excellence – East, Baltimore, MD; ²Washington, DC VAMC, MS Center of Excellence, Washington, DC;

³Orlando VA Medical Center, Multiple Sclerosis Center of Excellence – East, Orlando, FL

Background: As the number of US Food and Drug Administration–approved disease-modifying therapies (DMTs) used to treat various forms of multiple sclerosis (MS) increases, the monitoring requirements for DMTs also increases. The use of the TheraDoc software program, a clinical decision support tool (DST), has been shown to improve patient care, comply with guidelines of care, optimize cost of care, create efficiencies of care, and minimize adverse drug events. **Objectives:** To compare the delay to treatment and monitoring using manual vs automated surveillance. To compare the time to manually enter data vs automated data for case management. To highlight medication adherence. **Methods:** TheraDoc has been used for case management in a variety of medical settings in the VA health care system; it has not been used for the care of patients with MS. A dashboard of patients with MS who are prescribed a DMT was developed with consideration of the VA criteria for use guidelines. Features include automation of laboratory results, flag reports, real time alerts, team communication, ability to export report, and various views of the dashboard. TheraDoc is linked to the VA's computerized patient record system and is currently being piloted at 3 sites. **Results:** The features of the TheraDoc MS DMT software program will be reviewed in a series of interactive figures. We assessed the experience of 120 patients (at present) with MS utilizing various DMT modalities. Demographic and clinical characteristics of all patients will be summarized. Outcomes including patient adherence, time to assess and order laboratory assessments, and follow-up with patients will be contrasted between the manual and automated surveillance groups. Errors in making assessment will be noted. Data will be analyzed, and we anticipate a decrease in delay to treatment, improved compliance with monitoring as established by the criteria for use guidelines with the use of TheraDoc clinical DST, and case management. Overall time efficiency will be contrasted between the manual and automated surveillance approaches. **Conclusions:** Using a clinical DST linked with an electronic health record providing the MS clinical team with required data improves MS DMT monitoring, decreases risk for adverse drug events of patients who receive DMT, and improves MS team communication.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Disease-modifying treatments in MS, Nursing management in MS

(MOC09) Multiple Sclerosis Disease Impact Monitoring: Longitudinal Exploration of the Relationship of Ocular Coherence Tomography to Computerized Cognitive Testing

Mark Gudesblatt,¹ Jared Srinivasan,¹ Olivia Kaczmarek,¹ Glen Doniger,² Daniel Golan,³ Jeffrey Wilken,⁴ Timothy Fratto,⁴ Robert C. Sergoff⁵

¹South Shore Neurologic Associates, Patchogue, NY; ²NeuroTrax Corporation, Modiin, Israel; ³Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel; ⁴Washington Neuropsychology Research Group, Washington, DC; ⁵Wills Eye Hospital, Philadelphia, PA

Background: Disease impact and change in routine care for people with multiple sclerosis (PwMS) is typically measured by combining Expanded Disability Status Scale (EDSS) score, magnetic resonance imaging (MRI) change, and reported relapse. Cognitive impact can be objectively tracked with digital cognitive assessment batteries (CABs) and retinal nerve fiber layer (RNFL) density (ocular coherence tomography [OCT]). Cognitive and RNFL impact are not quantified by traditional care approach (EDSS, MRI, relapse). Enhancing shared decision making to optimize PwMS's treatment selection could be accomplished by incorporating quantitative measures that provide objective examiner-independent information reflecting disease impact and possibly differentiating relapse from progression. Utilization of the NeuroTrax CAB to measure a global cognitive summary score (GCS) based on 7 tested domains (memory, executive function, visual spatial, verbal function, attention, information processing, motor skills) combined with OCT RNFL measures could lead to earlier detection of disease activity as well as assisting in optimal treatment selection. **Objectives:** To explore the relationship between rate of change in PwMS seen in both a global CAB score and various OCT measurements. **Methods:** Retrospective chart review of CAB and OCT scores collected in the process of routine care. Paired sample T-tests were done between percent change of 2 visits, 1 year apart, with GCS and the following OCT measurements (right OD and left OS eyes): RNFL (sublayers: global [G], nasal/temporal ratio [N/T]), papillomacular bundle (PMB), and macular volume (MV). **Results:** N = 103 (75% female, average age at first visit 51 ± 10). All regressions run between CAB-GCS and OCT

measures yielded P values $> .05$ at both visit 1 and visit 2, save for the CAB-GCS G-OS relationship. A significant difference was not observed ($P > .05$) in the percent change between visit 1 and 2 when comparing the following OCT measurements with the CAB GCS: G-OD&OS, N/T-OS, PMB-OS, and MV-OD&OS. Significance was observed ($P < .05$) of N/T-OD and PMB-OD. **Conclusions:** The relationships of global RNFL densities to global CAB scores remained the same after a year, which suggests that both measures identify disease change in a synchronous manner in monitoring disease progression of PwMS. Nonsignificance between percent changes suggests that OCT and cognitive scores change at similar rates at least within a year's period. A larger longitudinal study is suggested to further determine the relationship between OCT and cognitive changes over greater lengths of time.

Supported by: None

Disclosure: *Mark Gudesblatt:* Acorda, Amgen, Biogen, EMD Serono, Medtronic, Novartis, Sanofi, Saol Therapeutics, Teva (consulting fee); *Biogen, EMD Serono, Novartis, Sanofi, Teva (contracted research); Jared Srinivasan, Olivia Kaczmarek, Daniel Golan, Timothy Fratto:* Nothing to disclose. *Glen Doniger:* NeuroTrax (salary). *Jeffrey Wilken:* Biogen (contracted research); *EMD Serono (speakers' bureau); Genzyme (contracted research, speakers' bureau). Robert C. Sergott:* Biogen, Clene Nanomedicine, Heidelberg Engineering GmbH, Janssen Global Services, LLC, Medtronic, Merck & Co Inc (consulting fee); *Biogen, Clene Nanomedicine, Janssen Global Services, LLC, Medtronic, Nightstar, ThromboGenics NV (contracted research); Biogen, Genzyme Corporation, Novartis Pharmaceuticals Corporation, Teva Pharmaceutical Industries, Ltd (speakers' bureau).*

Keywords: Comprehensive care and MS, Equipment in MS, Natural history of MS

(MOC11) Improving Understanding of Clinical Phenotype for Patients with Multiple Sclerosis: Design and Implementation of Smarttools in Electronic Health Record Systems

Thomas Grader-Beck,¹ Yujie Wang,² Kathryn C. Fitzgerald,² Peter A. Calabresi,² Ellen M. Mowry²

¹School of Medicine, Johns Hopkins University, Baltimore, MD; ²Neurology, Johns Hopkins University School of Medicine, Baltimore, MD

Background: Patients with multiple sclerosis (MS) frequently require complex clinical care and decision making, including monitoring for relapses, chronic immunosuppressive therapy, and extensive serological and imaging work-up. Modern electronic health record (EHR) systems offer the opportunity to facilitate understanding of each patient's clinical phenotype by allowing discrete data collection, as well as summarization and presentation of critical information to providers. However, such tools often require custom design and build, for which many institutions do not have assigned resources. **Objectives:** To implement a comprehensive set of EHR-based Smarttools for patients with MS that can be shared among institutions to improve data quality and understanding of patient phenotype. **Methods:** The Epic EHR system was used to develop several Smarttools. A Smartform was designed to allow discrete data collection on items considered critical by MS experts. Content included date of diagnosis, documentation of relapse characteristics, such as date, duration and therapies, current and past immunosuppressive therapy, critical imaging and results of CSF studies. Longitudinal caption of disease impact was incorporated, including number of falls and ability to walk since last visit. Logic was applied to conditionally display the SmartForm for patients with a diagnosis of MS. A synopsis was designed to visualize longitudinal data, and a SmartPhrase was implemented to automate documentation of discrete data in provider notes. **Results:** Version 1 of the MS Smartform was implemented in August 2017. Over the course of 18 months, data on approximately 1000 unique patients were collected. The SmartForm was found to be easily accessible and easy to navigate by providers, and the completion rate was high. Based on the initial experience, version 2 of the Smartform was designed and was recently implemented with minor modifications to minimize erroneous data collection, and a predefined list of immunomodulatory therapies and more detailed information about reasons for starting and stopping therapies was included. Data entry for the full cohort of ~3000 patients is ongoing. With the support of the Epic Neurology Steering Board, all discrete data elements were incorporated into the Epic Foundation system to facilitate implementation of the Smartform at other institutions. **Conclusions:** EHR systems provide opportunities to improve understanding of complex clinical phenotype. We built a comprehensive Smartform that facilitates discrete data collection and review for patients with MS. This tool is being made available in the Epic Community library and can be implemented without charge at other institutions. Besides offering a better understanding for individual patients, it is our hope that this SmartForm will contribute to better streamlining of data

collection across institutions and ultimately, better outcomes for patient care and collaborative research.

Supported by: None

Disclosure: *Thomas Grader-Beck:* AbbVie, Celgene (contracted research); *Eli Lilly (consulting fee). Yujie Wang, Kathryn C. Fitzgerald:* Nothing to disclose. *Peter A. Calabresi:* Biogen, Disarm Therapeutics (consulting fee). *Ellen M. Mowry:* Biogen, Sanofi, Genzyme (contracted research); *Biogen, Sun Pharma (site PI); Teva (PI clinical trial); UpToDate (royalty).*

Keywords: Comprehensive care and MS, Electronic health record

(MOC12) Changing Language to Acknowledging Patients' Perceptions of Treatment in Multiple Sclerosis Care

Tim O'Maley

Nursing, MS Queensland, Brisbane, QLD, Australia

Background: The language we use in supporting our patients is crucial in fostering a long-term relationship based on trust and understanding. However, it is important to remember that information does not equal education, and the ability to make the complexities of medicine comprehensible for patients is an important skill. Two recently published papers of almost opposing direction have highlighted the message that "I" can do better. Yeandle et al strongly emphasized the increased role patients and their families have in shared decision making, noting "its success is reliant on effective patient-physician communication." Burke et al in their paper on management of surplus suffering discuss how health care providers can negatively impact the perception of the disease and recognizing our control of information sharing may "go a long way to improving clinical encounters with patients and ultimately lead to greater satisfaction in care and shared decision making." **Objectives:** This presentation will briefly explore the core concepts of recent literature that have resulted in a shift in language used by the author in delivery of both clinical patient care and group educational opportunities, and the qualitative patient responses that emphasize the value of patient centricity in multiple sclerosis care. **Methods:** None. **Results:** None. **Conclusions:** None.

Supported by: None

Disclosure: *Tim O'Maley:* Biogen, Novartis, Roche (consulting fee).

Keywords: Comprehensive care and MS, Disease-modifying treatments in MS, Nursing management in MS

NEUROIMMUNOLOGY AND DISEASE MODELS

(NDM01) Efficacy of the Influenza Vaccine in Multiple Sclerosis Patients: A Systematic Review and Meta-analysis

Jackie T. Nguyen,¹ Patrick Hardigan,¹ Michelle Demory Beckler,¹ Marc Kesselman²

¹Dr. Kiran C. Patel College of Allopathic Medicine, Nova Southeastern University, Davie, FL;

²Dr. Kiran C. Patel College of Osteopathic Medicine, Nova Southeastern University, Davie, FL

Background: Multiple sclerosis (MS) is a neurodegenerative disease thought to be of autoimmune origin. It leads to the development of neurologic symptoms and increases the risk of infection from communicable diseases. Thus, vaccines are endorsed to mitigate this risk. However, it has not yet been confirmed whether these patients' dysfunctional immune system combined with taking immunosuppressants can lead to a dampened immune response against the influenza vaccine. Infection with the influenza virus is a concern for patients with MS. Previous research on patients with MS who have received the influenza vaccine focuses on safety and relapse rates. Studies that focus on the immune response mounted against the vaccine in these patients are scant. **Objectives:** This study serves to compile this previous research to provide a comprehensive picture of the efficacy of the influenza vaccine in patients with MS. **Methods:** This was done through a systematic review and meta-analysis. **Results:** The results of this study suggest that patients with MS can mount an adequate immune response to the influenza vaccine when compared to healthy controls. Most of the immunotherapies these patients are on do not appear to affect this immune response. **Conclusions:** Therefore, the influenza vaccine should continue to be recommended to patients with MS.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Immunology and MS, Vaccination in MS

NONIMAGING BIOMARKERS

(NIB01) Higher Sensitivity of Quantitative Reverse Transcriptase–Polymerase Chain Reaction Compared with Flow Cytometry for Quantification of B Cells After Anti-CD20 Monoclonal Antibody TherapyIsmahane Touil Allaoui,¹ Marija Colic,¹ David Leppert,² Friedrich Raulf,¹ Gisbert Weckbecker¹¹Novartis Pharma AG, Basel, Switzerland; ²Neurologic Clinic and Policlinic, University Hospital Basel, University of Basel, Basel, Switzerland

Background: Preclinical and clinical evidence have shown promising results for CD20+ B-cell-target-based therapies in multiple sclerosis (MS). Ofatumumab, a fully human anti-CD20 monoclonal antibody (mAb), depletes circulating peripheral B cells and is in phase 3 development for relapsing MS. **Objectives:** Compare sensitivity of reverse transcriptase–polymerase chain reaction (RT-PCR) and current standard fluorescence-activated cell sorting (FACS) for quantification of CD20+ B-cell depletion. **Methods:** Raji B cells, stained with anti-CD19-FITC (H1B19) and anti-CD20-PE (2H7), were spiked at 12 dilutions (0.3-100 000 cells) into 100 000 THP-1 cells (CD19-/20-); each sample was split for FACS and RT-PCR. Absolute counting beads (Invitrogen) were used to define minimal FACS detection level of CD19+/20+ cells by BD Fortessa (stopping gate fixed at 2000 beads). Limit of detection and limit of blank were determined as per international guidelines. For RT-PCR, total RNA was extracted (RNeasy Mini kit). Duplex TaqMan assay with CD19-VIC (Hs00174333m1) and CD20-FAM (Hs00544818m1) probes was run in triplicate, after reverse transcription (SuperScript III). Copy numbers were determined via a standard curve from serial dilutions of linearized quantified plasmid of CD19/CD20 target sequences. PCR reactions were set up by an Echo 525 acoustic liquid handler with a total volume of 2.5 μ L. **Results:** A limit of blank of 70 cells and limit of detection of 90 Raji cells spiked into 100 000 THP1 were observed by FACS with no difference between CD19+ and CD20+ cells. A reliable correlation between spiked and bead-extrapolated counts of approximately 300 cells was observed. Sensitivity of RT-PCR was assessed similarly; efficiency for both TaqMan assays was >98% with limit of detection of 2 copies of mRNA. On average, Raji cells expressed 15-20 copies of CD19 and 60-100 copies of CD20 transcripts per cell. A reliable correlation was seen for CD19 and CD20 down to 10 spiked cells. **Conclusions:** Quantification of total B cells in blood and tissue after anti-CD20 mAb treatment by sensitive and specific RT-PCR seems feasible. In addition to simpler sample logistics, this method can measure CD20 gene expression directly.

Supported by: None**Disclosure:** *Ismahane Touil Allaoui, Marija Colic, Friedrich Raulf, Gisbert Weckbecker:* Novartis Pharma AG (salary). *David Leppert:* Novartis Pharma AG (former employee).**Keywords:** Disease-modifying treatments in MS, Immunology and MS**(NIB02) Quantification of Smooth Pursuit Dysfunction in Multiple Sclerosis**Neda Dastgheyb,¹ Miryam Palomino,² Annalise Miner,² Revere P. Kinkel,² Dorion Liston,³ Jennifer S. Graves⁴¹University of California San Diego (UCSD), La Jolla, CA; ²UCSD, San Diego, CA; ³NeuroFit, Mountain View, CA; ⁴Neurosciences, UCSD, San Diego, CA

Background: A significant proportion of patients with multiple sclerosis (MS) have efferent visual system dysfunction that may impair ability to read or work or lead to dizziness, gait impairment, or falls. Quantifying oculomotor deficits also offers promise as a biomarker of MS disease burden. **Objectives:** The purpose of this study is to use a novel noninvasive NASA-developed eye tracker (neuroFit ONE) to detect and quantify abnormalities in oculomotor function in patients with MS vs healthy controls. **Methods:** We recruited consecutive subjects from the UCSD MS clinic who met the 2017 published MS criteria and healthy controls without any history of neurologic disease, head trauma, or other source of afferent/efferent visual deficit. The neuroFit ONE eye tracker was used to measure latencies, acceleration, gain, saccadic correction amplitudes, proportion of smooth vs saccadic eye movements, direction tuning, and speed tuning in participants with MS vs healthy controls. Each test was repeated 3 times to assess reproducibility and to generate a mean value. A composite of the individual tests was generated (nfit). Comparisons of the oculometrics with case status were assessed by regression methods. **Results:** Ten patients with MS and 17 healthy controls have been enrolled to date. The mean overall composite score of oculomotor function (nfit score) was worse in patients with MS vs controls (-1.6, 95% CI -2.9 to -0.20, $P = .026$). Mean latency for initiating smooth pursuit was 6

ms longer for MS participants (95% CI 0.65-12.6, $P = .031$). There was decreased mean initial acceleration of eye movements (-37.2, 95% CI -67.6 to -6.9, $P = .018$) in the MS eyes. The proportion of smooth vs saccadic eye movement was lower in MS eyes (-16%, 95% CI -30% to -23%, $P = .024$). The additional metrics measured showed similar patterns. **Conclusions:** We were able to comprehensively capture and quantify oculomotor dysfunction in participants with MS compared to controls, including aspects of smooth pursuit that have not previously been well quantified in MS.

Supported by: None**Disclosure:** *Neda Dastgheyb, Miryam Palomino, Annalise Miner:* Nothing to disclose. *Revere P. Kinkel:* Biogen (speakers' bureau). *Dorion Liston:* neuroFit (founder). *Jennifer S. Graves:* Biogen, Octave, Genentech (research support); Novartis, Genentech, Alexion, Celgene (consulting fee).**Keywords:** Neuro-ophthalmology

NEUROPHYSIOLOGY, NEUROPSYCHOLOGY, AND NEUROPSYCHIATRY

(NNN02) Education as a Moderating Variable in the Relationship Between Patient Self-Perception of Cognitive Impairment and Symbol Digit Modalities Test Performance in Multiple Sclerosis

Elizabeth Kera, William A. Tsang, Nina A. Curko, Lee S. Iffhar, Florian Thomas, Krupa Pandey

Neurology, Hackensack University Medical Center, Hackensack, NJ

Background: Approximately 40%-60% of patients with multiple sclerosis (MS) experience some degree of cognitive impairment. Available evidence suggests greater education level serves as a protective factor and is related to cognitive reserve and premorbid intelligence. Individuals with higher levels of education may be differentially affected by reduced processing speed, when compared to those with lower levels of education. Alternatively, there may be differences in degree of awareness of cognitive deficits as a result of educational attainment. Numerous studies signal the importance of early detection in cognitive impairment in overall MS disease burden and treatment outcomes. However, many clinicians rely on patient self-report in determining whether to refer a patient for in-depth cognitive testing. It can be difficult for clinicians to accurately gauge cognitive impairments during brief clinical visits, and certain factors may affect how patient self-report is interpreted (ie, education level, cognitive reserve). Recent standard-of-care guidelines have been published which outline the utility of the Symbol Digit Modalities Test (SDMT) as an early screening measure to be used to establish baseline cognitive functioning. **Objectives:** The current study ($N = 75$) evaluated whether education level moderated the association between SDMT scores and patient self-report of cognitive dysfunction. **Methods:** Patient self-report of cognitive dysfunction was evaluated via EMR review of neurology consult notes. All patients were administered the SDMT after their neurology consult visit, and highest level of educational attainment was obtained via self-report. **Results:** Linear regression modeling compared SDMT scores of those who endorsed cognitive symptoms ($n = 22$; $M = -1.09$, $SD = 1.1$) to those who did not ($n = 53$; $M = -0.19$, $SD = 1.16$). Results showed a significant difference in model 1, $F_{1,72} = 9.4$, $P = .003$, indicating that those who endorsed cognitive deficits yielded lower SDMT scores. Model 2 did not support education in moderating this effect. **Conclusions:** While patient self-perception did correlate with SDMT performance, level of education of the patient did not affect this relationship.

Supported by: None**Disclosure:** *Elizabeth Kera, William A. Tsang, Nina A. Curko, Lee S. Iffhar:* Nothing to disclose. *Florian Thomas:* Genentech, Novartis, Sanofi (speakers' bureau). *Krupa Pandey:* Alexion, Biogen, Genentech, Novartis, Sanofi (speaking/consulting).**Keywords:** Comprehensive care and MS, Management of activities of daily living in MS, Neuropsychology**(NNN03) Preliminary Cognitive Outcomes Following Mesenchymal Stem Cell Therapy in Multiple Sclerosis**Lisa A.S. Walker,^{1,2,3} Jason A. Berard,^{4,5} Maha Abu-AlHawa,² Ruth Ann Marrie,⁶ James Marriott,⁷ Harold Atkins,⁸ David Courtman,⁹ Mark S. Freedman¹⁰¹Psychology, University of Ottawa Brain and Mind Research Institute, Ottawa, ON, Canada; ²Neuroscience, Ottawa Hospital Research Institute, Ottawa, ON, Canada;³Psychology, Carleton University, Ottawa, ON, Canada; ⁴Neuroscience, Ottawa Hospital Research Institute, Ottawa, ON, Canada; ⁵University of Ottawa, Ottawa, ON, Canada;

⁴Department of Medicine, University of Manitoba, Winnipeg, MB, Canada; ⁷Internal Medicine, University of Manitoba, Winnipeg, MB, Canada; ⁸Cancer Therapeutics, Ottawa Hospital Research Institute, Ottawa, ON, Canada; ⁹Ottawa Hospital Research Institute, Ottawa, ON, Canada; ¹⁰University of Ottawa and the Ottawa Hospital Research Institute, Ottawa, ON, Canada

Background: Mesenchymal stem cells (MSCs) are being investigated as an alternative disease-modifying therapy for multiple sclerosis (MS) given their immunomodulatory and tissue repair properties. MSCs are multipotent progenitor cells that can differentiate into mesodermal cells with neuroprotective and pro-oligodendrogenic properties. Little is known about potential effects on cognition. **Objectives:** To evaluate cognition following MSC therapy over 48 weeks. **Methods:** 28 individuals with inflammatory MS (17 relapsing-remitting MS, 7 secondary progressive MS, 4 primary progressive MS) were enrolled in a randomized, double-blind, sham-controlled cross-over study of autologous MSC with the primary outcome determined at 24 weeks. Participants were randomized to receive either a single intravenous infusion of MSC or a sham infusion at week 0, then crossed over at 24 weeks to the alternate treatment arm for a further 24 weeks of observation. Participants underwent a comprehensive neuropsychological battery at weeks 0, 24, and 48. Cognitive domains assessed included attention/information processing speed, language, visual perception, learning, memory, and executive functioning. To account for potential practice effects associated with serial testing, data were analyzed using reliable change analyses at the individual level. Performance on any given cognitive task was considered improved or declined if most of those who demonstrated change (at least 3 or more) obtained significant RCI values (± 1.64). **Results:** Participants were 15 females/13 males [Expanded Disability Status Scale score 4.27 [1.25], age 37.36 [5.21] years, education 13.64 [1.61] years]. Immediately after treatment, relative stability was noted for most cognitive tasks. Nonetheless, some change was detected. Decline was observed in some aspects of attention/information processing speed, visual learning and memory, as well as language. Improvement was noted in verbal learning and memory, as well as visual perception. In the early-treatment group, where longer-term follow-up was possible, there was a trend for performance to return to pretreatment baseline, with the exception of visual learning and memory, which remained below baseline levels. **Conclusions:** Except for visual learning and memory, there appears to be little detrimental effect of MSC therapy on cognition. While some changes may occur in the initial period following treatment, these appear to be transient and, in general, return to baseline over time.

Acknowledgments: Funded in part by the MS Scientific Research Foundation and Research Manitoba

Disclosure: Nothing to disclose

Keywords: CNS repair, Cognition in MS, Disease-modifying treatments in MS

(NNN04) Relationship Between Expanded Disability Status Scale Scoring and Attention Performance in People with Multiple Sclerosis

Mauricio O. Bando,^{1,2} Ana Maria Canzonieri,² Alice E. Dias,¹ Roger P. Silva,^{1,2} Giuliana M. Ferrero,² Ibis Ariana P. De Moraes,² Talita D. Da Silva,² Marcelo Massa²

¹Neurorehabilitation, ABEM-Brazilian Multiple Sclerosis Association, Sao Paulo, Brazil;

²EACH, USP-Sao Paulo University, Sao Paulo, Brazil

Background: Multiple sclerosis (MS) is a degenerative, autoimmune, and chronic neurologic pathology. In addition to the symptoms of spasticity, fatigue, muscle weakness, numbness, and urinary incontinence, among others, report of attention performance difficulty is very common. The Expanded Disability Status Scale (EDSS) is a method of quantifying disability in people with MS, which is scored from 0 to 10. The higher the scoring, the greater the person's functional disability. **Objectives:** To verify and analyze the relationship between EDSS scoring and attention performance in people with MS. **Methods:** A quantitative study was performed with 41 people diagnosed with relapsing-remitting MS (RRMS), aged 23-58 (mean [SD] = 42.70 [10.62]) years, 14 men (34.1%) and 27 women (65.9%), with EDSS score from 0 to 6.5 and time of diagnosis between 1 and 26 (mean [SD] = 10.09 [6.67]) years. For evaluation, an interview was conducted to collect data and a battery of neuropsychological attention tests was applied to each patient. The SPSS software was used for data analysis. **Results:** It was observed that 20 patients (48.8%) presented alteration of sustained attention, 29 patients (70.7%) presented alteration of alternating attention, and 27 patients (65.9%) presented alteration of divided attention. There was a negative association between EDSS scoring and sustained attention performance ($P < .0001$) and alternating attention performance ($P = .037$). That is, the higher the EDSS scoring, the worse the performance of sustained and alternating attention. There was no significant association between EDSS scoring and divided attention performance ($P = .094$). **Conclusions:** It is suggested

from the results of this study that the level of disability status may affect sustained and alternating attention performance of people with MS.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Multiple sclerosis, EDSS, Neurological disease, Neuropsychology, Psychological issues and MS

(NNN05) Objective Measurement of Cognitive Impairment in Multiple Sclerosis Patients Using Novel Computerized Testing

Roberto Bomprezzi,¹ Kerime Ararat,² Kara Smith,¹ Reina Benabou³

¹Neurology, University of Massachusetts Medical School, Worcester, MA; ²Neurology, University of Massachusetts Memorial Medical Center, Worcester, MA; ³Cognivue Inc, Victor, NY

Background: Cognivue is a US Food and Drug Administration-cleared computerized testing tool rooted in adaptive psychophysics and designed to assess early signs of cognitive impairment. Cognitive impairment has a substantial impact on productivity and quality of life in patients with multiple sclerosis (MS), but testing has been limited. A brief, easy-to-administer neuropsychological test could increase the frequency of routine assessment of cognitive impairment among patients with MS, leading to a positive impact on MS management. **Objectives:** At the completion of this presentation, participants should be able to assess the reliability of Cognivue as a cognitive assessment tool in MS. **Methods:** The study was conducted at the University of Massachusetts Medical School between June 2016 and May 2017 and enrolled consecutive patients who consented to testing. Study participants completed the Expanded Disability Status Scale (EDSS), Symbol Digit Modality Test (SDMT), Nine-Hole Peg Test, timed 25-foot walk, and 10-minute Cognivue testing (basic motor and visual ability, perceptual processing, and memory processing). Statistical analyses using a 1-way analysis of variance were performed to determine differences between neuropsychological testing methods. **Results:** Thirty-six patients (mean age 48.6 [range 20-74] years, 78% female [n = 28/36]), completed the various tests. Based on Cognivue scores, 50% of patients were categorized as having normal cognitive function (mean 84.7; EDSS score 2.64), 33.3% as having low to moderate cognitive impairment (mean 66.0; EDSS score 3.38), and 16.7% as having severe cognitive impairment (mean 39.2; EDSS score 5.17). Overall Cognivue scores demonstrated statistically significant correlations with EDSS (Pearson correlation coefficient -0.54), SDMT (0.67), and timed 25-foot walk (-0.56). No relationship was seen between patient age and Cognivue scores. All key cognitive domains were equally affected. **Conclusions:** Cognivue is beneficial in detecting early stages of multidomain cognitive impairment in patients with MS, providing a potential opportunity for early intervention strategies to improve patient outcomes.

Supported by: None

Disclosure: Roberto Bomprezzi, Kerime Ararat: Nothing to disclose. Kara Smith: Acorda (served on expert panel). Reina Benabou: Cognivue, Inc (CMO).

Keywords: Neuropsychology, Cognition

PROGRAMS

(PGM01) The Use of a Multiple Sclerosis Documentary Film Screening Program as an Educational Intervention to Increase Knowledge and Awareness About MS and Support Resources

A. Suzanne Boyd,¹ Shelby Veri²

¹The University of North Carolina at Charlotte School of Social Work, Charlotte, NC;

²Department of Public Health Sciences, University of North Carolina at Charlotte, Charlotte, NC

Background: Typical multiple sclerosis (MS) educational methods include brochures, handouts, community presentations, or online resources. First-person accounts of experiences living with illness are common ways to learn about health conditions. Film is a nonthreatening modality to increase knowledge and awareness about MS and its impact. **Objectives:** Increase MS knowledge and awareness via documentary film screening *When I Walk*; Increase awareness of local MS resources. **Methods:** Study approved by UNC Charlotte institutional review board. This post-test only study included 68 participants who attended an evening MS film documentary screening event in November 2019. *When I Walk* is the first film in a trilogy of films about MS. The second film, *When We Walk*, premiered in 2019, and the third film, *When They Walk*, is in production. 45 participants completed an online event survey. Participants

ranged in age from 18 to 74 years and most were female (79.1%; n = 35). Just over half of the participants were Caucasian (n = 27); 7 were black/African American and were well-educated with university degrees. The film screening was delivered by a social work researcher and health services doctoral student and shown in a 600-person auditorium. A slide-show with information about the film, panelist biographies, vendors, and follow-up MS film and educational events scrolled before and immediately after the film screening. A panel discussion including 5 participants immediately followed the film screening: local National MS Society representative, university ADA director, a physician's assistant specializing in neurology, 1 person with MS, and 1 person with MS who is also a health care professional and support group facilitator. **Results:** Results suggest an overwhelmingly positive and enthusiastic outcome and impact on participants as a way to increase knowledge and awareness about MS and available resources. For example, 35 participants rated the MS film screening experience as "excellent," 41 participants "strongly agreed" or "agreed" that participation in the MS documentary screening increased their knowledge of MS and its related symptomatology, 36 participants "strongly agreed" or "agreed" that participation in the MS film screening increased their knowledge of available resources at UNC Charlotte and the surrounding area, and 41 participants (95%) stated that the film screening helped them to better understand the social and cultural views of others who have had different life experiences. **Conclusions:** Participant responses support using film documentary as an effective, creative, and friendly intervention to increase knowledge and awareness about MS and available resources and to increase collaborative partnerships between the university and community partners. Few examples exist in the literature about using film as an educational learning tool to educate persons about physical illness. There are several study limitations to consider in future events.

Supported by: None

Disclosure: Nothing to disclose

Keywords: MS and the caregiver/family, Psychological issues and MS

(PGM03) Dance for MS: A Structured Dance Program Targeted for Multiple Sclerosis Patients

Carlos Eduardo Vervloet Sollero,¹ Keely Mason,² Brooke Borgert,³ Elizabeth Johnson,⁴ Jill Sonke,² Whitney Wilson,³ Tirisham V. Gyang¹

¹Neurology, University of Florida, Gainesville, FL; ²Center for Arts in Medicine, University of Florida, Gainesville, FL; ³University of Florida, Gainesville, FL; ⁴School of Theatre + Dance, University of Florida, Gainesville, FL

Background: Multiple sclerosis (MS) is a demyelinating disease of the central nervous system and a leading cause of disability. It presents multiple symptoms such as ataxia, weakness, and fatigue, impairing independence and quality of life. Although advances have been made in preventing disability, pharmacologic approaches to reverse it are not available. The best tool for functional recovery in MS is rehabilitation, typically physical and occupational therapies. Exercise therapies provide symptomatic benefit and are widely used in rehabilitation protocols. In Parkinson disease, different dance regimens have been shown to improve functional outcomes and to be superior to traditional exercise programs. Encouraged by this, we developed a structured dance class for MS, with specific targets, such as balance and exercise tolerance. The protocol was a collaboration of dance faculty, neurologists, and physical and occupational therapists. **Objectives:** To present the University of Florida's (UF's) "Dance for MS"—a dance program for symptomatic improvement and quality of life. **Methods:** Classes occur weekly, with 75-minute duration. They are taught by faculty from UF Center for Arts in Medicine and UF Health Shands Arts in Medicine, with an artist in residence. They start with a 15-minute seated warm-up, 15-minute barre exercise, 5-minute break, 15-minute center or across-the-floor section, 20-minute improvisation/dance composition, and a 5-minute cool down. Classes combine elements of modern dance, ballet, jazz, and social dance. **Results:** Classes launched in August 2018. Three to 6 people with MS attended each class, as well as 1 to 3 caregivers. In the past 18 months, the class performed interactive dances in community events such as the National MS Society Walk, HealthStreet's Night of Dance, and the Harn Museum's Museum Nights. While participant population is small, retention rate is high. Participants have reported improved balance, body awareness, and confidence in their movement. They have also appreciated the accessible approach to dance, and their enjoyment of the class. **Conclusions:** The Dance for MS program presents a feasible rehabilitation strategy for patients with MS, with a targeted approach to common symptoms in this population. It is presented in a social and ludic format, which may be beneficial for affective symptoms. Similar dance programs can be implemented as complementary rehabilitation strategies. Formal trials to measure the impact of the dance program are needed.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS, Comprehensive care and MS

(PGM04) Development of an Effective Age-Span Program for Women with Multiple Sclerosis: A Patient Perspective

Dina Jacobs,¹ Sona Narula,² Vanessa Zimmerman¹

¹Neurology, University of Pennsylvania, Philadelphia, PA; ²Neurology, Children's Hospital of Philadelphia, Philadelphia, PA

Background: Multiple sclerosis (MS) is a chronic, lifelong, unpredictable, and potentially highly debilitating neurologic condition that occurs more commonly in women than in men. It strikes most often in young adulthood, but can even start in childhood and adolescence. Young girls and adolescents with pediatric-onset MS, as well as adult women, often find it intimidating and overwhelming to navigate the complexities of the health care system. Typical MS programs may not focus on managing the impact of MS on puberty, fertility, pregnancy, the postpartum period, breastfeeding, and menopause. **Objectives:** To develop a Comprehensive Age-Span Program for Women with MS at the Children's Hospital of Philadelphia (CHOP) and the Hospital of the University of Pennsylvania (HUP), health care providers must identify the health care needs from the patients' perspective. The population includes women with MS at every age, from teenager to older adult. The purpose of this qualitative study will be to identify personal and health care needs of women with MS of all ages. The ultimate goal will be to provide patients and their families with high-quality education about their condition and establish a multidisciplinary team approach that will engage physicians, nurses, pharmacists, and social workers to provide optimal care and support that will help women with MS have better outcomes at every stage of life. **Methods:** Female patients with MS from both CHOP and HUP will be invited to participate in a single focus group. It is anticipated there will be 4 groups. In an open forum style, the group will be asked 10 open-ended questions to identify what services they would want to be available in a comprehensive age-span program. **Results:** The data will be analyzed using thematic analysis. **Conclusions:** It is anticipated that this study will reveal the needs of these women. The ultimate goal will be to develop an Age-Span Program that will meet the needs of this patient population.

Supported by: None

Disclosure: Dina Jacobs: Biogen, Genentech (consulting fee, contracted research); Celgene, EMD Serono, Sanofi Genzyme (consulting fee); MedImmune (contracted research). Sona Narula, Vanessa Zimmerman: Nothing to disclose.

Keywords: Age-Span Women's MS Program, Comprehensive care and MS

(PGM05) National MS Society Pathways to a Cure: An In-Person Educational Program for People Affected by Multiple Sclerosis

Elaine Liserio,¹ Eric Szafran,¹ Eric Garmon,¹ Kathleen Costello²

¹Advocacy, Services and Research, National MS Society, Denver, CO; ²National MS Society, New York, NY

Background: Important strategic goals for the National MS Society are to deliver breakthroughs to a cure and expand resources and reach to all those who are affected by multiple sclerosis (MS). The Society strives to empower people affected by MS to solve everyday challenges by informing them and connecting them to their communities, and the Society, so they can be more powerful than the challenges of MS. The Pathways to a Cure in-person program, conducted in cities across the country, is a key initiative in achieving Society goals. The program presented current research findings and strategies to help people meet the everyday challenges that MS imposes on them. **Objectives:** The objectives of the program were to 1) increase participant knowledge on the latest research breakthroughs, 2) increase awareness about wellness strategies and services resources that they can act on now to positively impact their health and quality of life, and 3) create connections among participants and those within the MS community. **Methods:** During 2019, 101 in-person, 3-hour Pathways to a Cure programs were held throughout the United States. The program consisted of didactic presentations on current research and wellness and lifestyle strategies. Presentations were followed by a facilitated Q-and-A session. Participants were requested to complete a postprogram survey to assess the impact of the program. **Results:** A total of 3519 individuals participated in the programs. Of those, 71% identified as living with MS, 72% were women, 77% identified as Caucasian, 15% black or African American, 6% Hispanic or Latino, and 2% Asian. Surveys were completed by 2216 (63%) participants. Survey results demonstrated that 95% of participants agreed or strongly agreed

that the Society is a source of support where they can find solutions; 91% of participants agreed or strongly agreed that they made connections to information, resources, people, or other sources of support; 87% of participants agreed or strongly agreed they had increased confidence and support to cope with the challenges of MS following the event; and 87% of participants indicated they planned to take action on information they learned. **Conclusions:** The National MS Society Pathways to a Cure in-person program reached a large number of people affected by MS throughout the United States. Survey results indicate that the objectives of the program were met. This was a successful method to reach people affected by MS with research breakthroughs and practical strategies to help them live their best lives. A next step that is being explored to extend the reach is offering the program virtually through a streaming platform.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Comprehensive care and MS, Management of activities of daily living in MS, Wellness*

(PGM06) How Well Do Junior Neurology Residents Recognize Multiple Sclerosis? Analysis of the "Close the Loop" Clinical Acumen Project

Emily M. Schorr, Jamie Nichols, Rachel Brandstadter, Stephen Krieger

Neurology Department, Icahn School of Medicine at Mount Sinai, New York, NY

Background: Multiple sclerosis (MS) misdiagnosis is an important issue with major potential consequences. Residency training is where future neurologists develop their clinical acumen and diagnostic ability. We describe MS/demyelinating disease cases presented by junior neurology residents. **Objectives:** To evaluate initial diagnostic accuracy and identify educational needs to prevent MS misdiagnosis and enhance quality of care. **Methods:** From July 2010 to June 2016 all patients independently assessed and presented by on-call junior neurology residents during daily morning report were logged, including case demographics and the initial diagnostic impression. Cases were subsequently revisited to "close the loop" with a final diagnosis. Cases were retrospectively categorized as neurological (subdivided by localization and etiology [eg, MS/demyelinating, stroke]) or "non-neurological" (eg, medical, psychiatric). Accuracy of the initial diagnosis was determined and errors were fully characterized. **Results:** Of the total 1301 cases, 4.4% carried a final diagnosis of MS/demyelinating disease (n = 57). Most of these patients were evaluated in the emergency department (80.7%), and most were admitted to the neurology service. Resident accuracy for MS/demyelinating disease cases was slightly higher than the overall case accuracy (66.7% vs 64.0%, respectively). There were 11 cases of MS/demyelinating disease that were initially mistakenly diagnosed as other neurologic conditions. Only 1 MS case was missed at the neurologic/non-neurologic decision point, while this type of error represented a large proportion of errors in the entire database (49.1%). Residents were more likely to miss true MS/demyelinating disease in men (6/11 [54.5%]). Of the 22 cases incorrectly deemed to be MS/demyelinating, 40.9% of errors (9/22) were at the neurologic/non-neurologic level, including 2 psychiatric and 6 medical cases. Diseases mistaken for MS/demyelinating disease include CNS neoplasm (n = 3) and ischemic stroke (n = 3). **Conclusions:** "Close the Loop" represents an educational initiative to provide feedback to neurology residents for improvement in clinical acumen. Despite the relatively small number of MS cases presenting in the acute hospital setting, resident diagnostic accuracy for MS/demyelinating disease was similar to overall accuracy. Analysis of errors made represent an important opportunity to improve recognition and hopefully enhance quality of inpatient care of MS.

Supported by: None

Disclosure: *Emily M. Schorr, Jamie Nichols, Rachel Brandstadter: Nothing to disclose. Stephen Krieger: Biogen, EMD Serono, Genentech, Genzyme, Mallinckrodt, MedDay, Novartis, Teva, TG Therapeutics (consulting fee, advisory work); Biogen, EMD Serono, Genentech, Novartis (speakers' bureau, nonpromotional speaking).*

Keywords: *Imaging and MS, Medical education*

(PGM07) Time to Adult: Transitioning from Pediatric to Adult Health Care in Demyelinating Disorders

Katherine Chapman,¹ Denise Maddox,¹ Lana Harder,^{2,3} Patricia Plumb,^{4,5} Benjamin Greenberg,⁶ Morgan McCreary²

¹Neurology, UT Southwestern Medical Center and Children's Health, Dallas, TX;

²Neurology, UT Southwestern Medical Center, Dallas, TX; ³Neuropsychology, Children's Health Dallas, Dallas, TX; ⁴UT Southwestern Medical Center, Dallas, TX; ⁵Neurology, Children's Health, Dallas, TX; ⁶Department of Neurology, UT Southwestern, Dallas, TX

Background: Transition to adult health care is a challenge in pediatric populations with chronic medical conditions, such as multiple sclerosis (MS). Patients require knowledge and proficiency in managing their health, engaging in wellness behaviors, and understanding health insurance and community resources. We developed a formal transition program to support our adolescent patients. Beginning at age 14, patients diagnosed with MS or related disorders and their parents/caregivers complete a questionnaire during annual clinic visits. Questions relate to their readiness to transition, knowledge of their condition, post-high school plans, stress and anxiety levels relative to transition, emotional support, quality of life, sleep, and priorities regarding information they would like on future aspects of care. Based on results, tailored education plans are used to improve patient knowledge and proficiency. Patients are tracked over time relative to their successful transition into adult care settings. **Objectives:** Examine the results from patient- and parent-completed questionnaires at baseline and 1-year postbaseline. Assess improvement in knowledge, readiness to transition, and priorities regarding information they would like on future aspects of care. **Methods:** Exploratory analysis of changes from baseline were analyzed using the Wilcoxon signed-rank test when numeric variables were at least ordinal. Binary variables were analyzed using Fisher exact test. Multiple testing adjustments were not performed. **Results:** To date, 67 patients were seen for baseline visits and 21 returned for a 1-year follow-up visit. For those patients who were seen 1-year postbaseline, there was an increase in the patient- and parent-reported Readiness scores ($P = .02$ and $P = .001$, respectively). Additionally, there was a decrease in the parent-reported support rating ($P = .004$). Lastly, 33 patients and 18 parents ranked a series of future aspects of their care in which they want information with managing their condition as the most important for most participants, followed by medication knowledge. **Conclusions:** Preparing adolescents to manage their own health care is critical, especially when faced with a chronic neurologic illness like MS. We noted an increase in patient and parent readiness for transition in our program across time. Future research will seek to identify factors that affect patient ability to successfully transition to adult health care.

Supported by: None

Disclosure: *Katherine Chapman: Genentech (speakers' bureau). Denise Maddox, Lana Harder, Patricia Plumb, Morgan McCreary: Nothing to disclose. Benjamin Greenberg: Alexion, EMD Serono, Novartis, Roche (consulting fee).*

Keywords: *Comprehensive care and MS, Psychological issues and MS, Transition and pediatric health care*

(PGM08) Successful Pilot of MS VA-ECHO Tele-education Program for Rural Providers

Lynda R. Hillman,¹ Jodie K. Haselkorn²

¹MS Center of Excellence, West, VA Puget Sound Healthcare, Seattle, WA; ²Department of Epidemiology, University of Washington, Seattle, WA

Background: Comprehensive care for persons with multiple sclerosis (MS) requires expert interdisciplinary teams. While such care is readily available at MS specialty centers in academic and urban environments, health care providers who live in smaller cities and in rural areas often do not have MS specialty expertise, and persons with MS living in these rural areas find it difficult to access appropriate care. Solutions are needed for this gap in care. The Veterans Affairs (VA) Healthcare System's Extension for Community Healthcare Outcomes (ECHO) allows specialists to share expertise with rural providers, enabling these clinicians to provide care not previously available in their communities and to save veterans from traveling long distances to access specialty care. Although VA-ECHO offers programs in many specialty areas, it did not yet have an MS program. As nearly 29,000 veterans in the VA Healthcare System live with MS, such a program is needed. **Objectives:** 1) To develop a pilot MS VA-ECHO program introducing the basic concepts of MS care most relevant to rural providers, including those which may be managed locally and which may be managed by or in collaboration with specialty MS providers. 2) To analyze audience evaluation of the ECHO presentation in relation to their educational needs and by provider discipline. 3) To ascertain focus for future program development and content. **Methods:** We used the traditional ECHO 3-part format: 1) didactic material; 2) case study; 3) audience questions and discussion. The pilot session was 75 minutes total, with ~45 minutes didactic, ~15 minutes case study, and ~10-15 minutes of question-and-answer and discussion. Presenters were a doctorally prepared nurse practitioner (NP) with MS certification and an MS specialist physiatrist, both with extensive practice in MS specialty centers. Content covered an overview of demographics, neuroimmunology and neuropathology, disease-modifying therapies (DMTs), symptom management, and patient and provider resources with special attention to rural applications. Audience metrics were collected. **Results:** The

audience total of 119 clinicians included physicians; advanced-practice providers; registered nurses; physical, occupational, and speech therapists; pharmacists; and social workers, among other disciplines. 96% of the audience found the material relevant to their practice, 45% indicated their practice would change to incorporate information learned, and 95% would like to hear the presenters again. A local focus group of the target audience found that nurses were strongly interested in MS certification. End-of-quarter regional ECHO all-program evaluations included over 400 requests for more MS material. **Conclusions:** This pilot MS VA-ECHO session was highly successful. There is demand for an expanded program and for material on MS nursing certification. Program development is underway for the expanded MS VA-ECHO series.

Supported by: None

Disclosure: *Lynda R. Hillman: Celgene (advisory).* *Jodie K. Haselkorn: Nothing to disclose.*

Keywords: *Comprehensive care and MS, Telehealth*

(PGM09) Current Topics in MS Webinar Series: A Professional Education Collaboration Between the National MS Society, Consortium of Multiple Sclerosis Centers, and the VA MS Centers of Excellence

Lynette Coleman,¹ Coleen Friedman,² Jaimie Henry,³ Doris Lill,⁴ Heidi W. Maloni,⁵ Angela Young,⁶ Kathleen Costello⁷

¹Advocacy, Services and Research, National Multiple Sclerosis Society, New York, NY;

²National MS Society, Bradenton, FL; ³Specialty Services, VA MS Center of Excellence West, Seattle, WA; ⁴Advocacy, Services and Research, National Multiple Sclerosis Society, Denver, CO; ⁵Washington, DC VAMC, MS Center of Excellence East, Washington, DC; ⁶Education, VA MS Center of Excellence – East, Baltimore, MD; ⁷National MS Society, New York, NY

Background: Multiple sclerosis (MS) is a complex disease that requires a highly educated workforce. To help meet the educational needs of MS health care professionals, the National MS Society (NMSS), Consortium of Multiple Sclerosis Centers (CMSC), and the Veteran Health Administration MS Centers of Excellence (VAMSCoE) collaborated to develop a professional educational program to provide evidence-based information on MS diagnosis and management. **Objectives:** Participants will 1) have easy access to evidence-based content with relevance to a variety of disciplines involved in MS care, 2) gain improved knowledge about MS and MS management, and 3) consider a change in their practice as a result of the information and resources presented. **Methods:** An educational program was developed entitled Current Topics in MS and it consisted of six 1-hour webinars on topics identified by health care providers including MS diagnosis, spasticity in MS, rehabilitation/tele-rehabilitation, reproductive care, MS in the African American population, and depression. Each webinar consisted of a 50-minute didactic presentation followed by a 10-minute facilitated question-and-answer session. Participants had the option to attend a live webinar or view a recorded presentation. Both were accredited for CME and CE. Following the live webinars, all registrants were sent a link to the webinar recording, and all participants were provided access to a program evaluation survey and a portal to complete a post-test and claim their free educational credits or a certificate of participation. **Results:** As of December 1, 2019, 866 health care providers (261 VA) attended the live or recorded webinars and 629 (73%) claimed continuing education credit or certificates. Survey results indicated 95% of respondents agreed or strongly agreed that the content was relevant to their current practice, 83% agreed or strongly agreed that participation improved knowledge, and 94% agreed or strongly agreed that participation encouraged them to consider a change to their practice. **Conclusions:** The Current Topics in MS webinar series is an important collaborative effort between the NMSS, CMSC, and MSVACoE. The series reached VA and non-VA health care providers with free and easily accessible professional education. Survey results indicate that participants found the programs useful, and a large percentage planned a change in their practice based on what they learned. Six new webinars are planned for 2020.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Comprehensive care and MS, Diagnosis, Psychological issues and MS*

(PGM10) Multiple Sclerosis Nurse Fellowship Pilot: A 6-Month Immersion

Marijean Buhse,¹ Patricia Melville,² June Halper,³ IOMSN Research Committee

¹South Shore Neurologic Associates, Patchogue, NY; ²MS Comprehensive Care Center, SUNY at Stony Brook, Stony Brook, NY; ³CEO-CMSC, Hackensack, NJ

Background: The current shortage of registered nurses (RNs) is expected to intensify as “baby boomers” age and retire from nursing. The retire-

ment wave of these nurses will create a drain on clinical expertise that is critical to quality patient care, especially in multiple sclerosis (MS). It is more difficult to quantify the loss of knowledge and understand its impact than it is to measure projected workforce demands due to retirement. Although nursing schools try to admit more students, they focus on preparing generalist nurses in acute care settings. This has resulted in unpreparedness of the nurse graduate for an independent role focusing on outpatient specialty practices such as MS. New RNs who enter MS practice will need much mentoring to learn to care for patients. This knowledge gap was identified by the International Organization of Multiple Sclerosis Nurses (IOMSN) in collaboration with the School of Nursing at the State University at Stony Brook. **Objectives:** A pilot program was developed to train a registered nurse enrolled in an RN to Bachelors (RNBS) degree nursing program. The student will complete a 6-month clinical fellowship in the care of patients with MS. RNs in the RNBS program were invited to apply by completing the application, submitting a 1-page essay describing their interest and experience in MS, and a reference from a professor. The call for applications described the plan for training and the 6-month clinical experience in a mentored environment. **Methods:** One student was selected to begin the fellowship in Fall 2019. Knowledge of MS was determined by questionnaires, and the student had strong knowledge of nursing but little of MS nursing. The student was then provided with a plan for training and a 6-month clinical experience in a mentored environment. **Results:** For the first 3 months, the student was precepted by a certified MS nurse practitioner and MS neurologists. A mid-fellowship student evaluation was developed which showed progress in both knowledge and skills in MS. During the second 3 months, the student will work with neuro-radiology, neuro-ophthalmology, neuro-urology, neuro-psychology and the outpatient department social worker. **Conclusions:** At the completion of the program, both the student and preceptors will complete evaluations documenting outcomes of this unique pilot project. It is anticipated that this program will generate similar training programs nationally and internationally.

Supported by: *This pilot project was funded by the International Organization of MS Nurses (IOMSN) supported by an educational grant from EMD Serono.*

Disclosure: *Patricia Melville: EMD Serono (speakers' bureau).* *Marijean Buhse, June Halper: Nothing to disclose.*

Keywords: *Nursing fellowship, Nursing management in MS*

(PGM11) Collaborative Working Between Multiple Sclerosis (MS) Nurses and a Pharmaceutical Company: An Educational Project from the Consortium of Multiple Sclerosis Centers (CMSC) Conference, Seattle, 2019

Mavis G. Ayer,¹ Karen Vernon,² Carmel Wilkinson,³ Lynda Kearney,⁴ Brenda Hamill⁵

¹Multiple Sclerosis, University Hospital Southampton Trust, Southampton, United Kingdom;

²Neurosciences, Salford Royal NHS Trust, Salford, United Kingdom; ³Multiple Sclerosis,

South Tyneside and Sunderland NHS Foundation Trust, Sunderland, United Kingdom;

⁴Neurology, NHS Fife, Fife, United Kingdom; ⁵Multiple Sclerosis, Northern Health and Social Care Trust, Belfast, United Kingdom

Background: Work collaboratively with key multiple sclerosis (MS) nurses from the United Kingdom and a pharmaceutical company in an educational project that has been shared with the wider UK MS community. **Objectives:** To share recent experience and reflections at the Consortium of Multiple Sclerosis Centers (CMSC) conference as part of a small focus group of UK MS nurses. **Methods:** Five nurses were supported by Roche to participate as a focus group of delegation to CMSC. The group met prior to the conference with the Roche team and a medical writer. The agenda was reviewed, and relevant topics/sessions were chosen and divided between the delegates, dependent on skill set and areas of interest and experience. The sessions chosen were areas of interest not only from the delegate's point of view, but also what would benefit the wider MS community in the United Kingdom. **Results:** The delegates attended the sessions (including the poster session) and fed back key learning messages attained to the medical writer using an agreed designed template. The delegates also had the opportunity to share with each other current practices and challenges and share experiences from areas of practice. This provided not only clinical supervision, but also reflection of own practice. From the feedback of the delegates, the medical writer produced a slide deck. The slide deck was given to the delegates to keep and to present to local members of their team and to the wider region. In addition, this slide deck is the intellectual property of the 5 delegates and will also be available via the United Kingdom Multiple Sclerosis Specialist Nurse's Association (UKMSSNA) slide deck library. **Conclusions:** This innovative project not only benefits the delegates, but also disseminates all learning and knowledge acquired to the wider MS communities. This way of working also provides greater transparency between the relationship of the sponsoring pharmaceutical companies and the delegates. The

delegates unanimously recommend this strategy of working during conferences and perhaps should be adopted by the rest of the pharmaceutical industry and health care professionals.

Supported by: None

Disclosure: *Mavis G. Ayer:* Biogen, Roche (sponsored delegate); Celgene, Merck, Novartis, Sanofi (consulting fee); Teva (sponsored education). *Karen Vernon, Carmel Wilkinson:* Biogen, Merck, Novartis, Roche, Sanofi, Teva (consulting fee). *Lynda Kearney:* Biogen, Novartis, Roche, Sanofi (consulting fee). *Brenda Hamill:* Biogen, Novartis (sponsored delegate); Roche (consulting fee).

Keywords: *Sharing best practice*

PSYCHOSOCIAL FACTORS

(PSF01) Differences in Depressive Symptomology Between Females and Males with Relapsing-Remitting Multiple Sclerosis

Chantel D. Mayo,^{1,2} Colleen Lacey,¹ Jodie R. Gawryluk^{1,2,3,4}

¹Psychology, University of Victoria, Victoria, BC, Canada; ²Institute on Aging and Lifelong Health, Victoria, BC, Canada; ³Division of Medical Sciences, University of Victoria, Victoria, BC, Canada; ⁴Island Health, Victoria, BC, Canada

Background: Up to 50% of individuals with multiple sclerosis (MS) experience depression, which greatly affects quality of life (Feinstein et al, 2014). Previous studies on depression in the general population have found that prevalence and reported symptoms differ as a function of gender, with higher rates in women (Salk et al, 2017). However, there are few studies on gender differences in symptoms of depression for people with MS, and the limited findings to date have been mixed (Patton et al, 2003; Théaudin et al, 2016). **Objectives:** The current study aimed to investigate whether there are differences between females and males with relapsing-remitting MS (RRMS) in overall depression scores as well as the types of depressive symptoms reported (somatic or cognitive). **Methods:** Demographic and Beck Depression Inventory, 2nd edition (BDI-2) raw scores for females and males with RRMS were downloaded with permission from the Multiple Sclerosis Outcome Assessments Consortium database (LaRocca et al, 2018). In addition to BDI-2 Total Scores, BDI-2 Somatic and Cognitive scores were also calculated for each participant (Beck et al, 1996; Vanheule et al, 2008). All statistical analyses were performed using RStudio. Data were first visually inspected using QQ-plots, followed by the Shapiro-Wilk Test of Normality, which indicated that the data deviated significantly from a normal distribution ($P < .001$). Thus, nonparametric Wilcoxon rank sum tests were used to compare BDI-2 Total Scores, BDI-2 Somatic Scores, and BDI-2 Cognitive Scores between females and males with RRMS. **Results:** Responses from 354 females and 140 males with RRMS were included in the analysis (mean age: females, 35.8 ± 9.5 years; males, 37.4 ± 10.2 years). Females reported significantly higher levels of overall depression (median = 9) compared to males (median = 7), $P = .032$. Furthermore, females endorsed significantly greater somatic symptoms (median = 7) than males (median = 5), $P = .026$. There were no significant differences in females' reports of cognitive symptoms (median = 2) compared to males' (median = 1), $P = .12$. **Conclusions:** Females with RRMS report higher levels of overall depression and somatic depressive symptoms compared to males with RRMS. Future research should focus on individuals with primary and secondary progressive MS to evaluate whether patterns of depressive symptomatology differ between females and males with progressive forms of MS.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Depression, Psychological issues and MS*

(PSF03) Me Too MS: Physical, Sexual, and Other Forms of Violence Experience in Women with Multiple Sclerosis

Jeta Pol-Patil,¹ Maria Claudia Manieri,² Tatenda Mahlanza,¹ Bonnie Ilene Glanz,¹ Elizabeth Misasi,¹ Laura T. Safar,³ Maria K. Houtchens¹

¹Partners Multiple Sclerosis Center, Brigham and Women's Hospital, Harvard Medical School, Boston, MA; ²Georgia State University, Department of Biology, Atlanta, GA; ³Lahey Health, Department of Psychiatry, Burlington, MA

Background: Sexual and physical violence against individuals with disabilities is widespread and linked to negative public health and social outcomes. In the past decade, emotional, physical, and sexual abuse of women with chronic illnesses and physical or cognitive disabilities has become increasingly recognized by public and health care providers. Women with disabilities have a higher likelihood of abuse, due to dependency on others, and stigma that surrounds their illness. **Objectives:** To explore the real-world prevalence of sexual, physical, and other forms

of abuse and their correlation with neurologic disability in women with multiple sclerosis (MS). **Methods:** Women with MS who saw a neurologist at the Partners MS Center between July 31, 2019, and October 21, 2019, were invited to participate in the study. They received a standardized anonymous questionnaire querying their previous sexual, physical, verbal, and emotional abuse experiences. Demographic information was collected on age, education, employment, and marital status. **Results:** 200/830 women (24%) completed the questionnaire. Mean (SD) age was 49 (11.39) years, and mean (SD) disease duration was 13 (8.67) years. 129 women (64.5%) were married or with a long-term partner. 134 patients (67%) were employed full or part-time. 186 women self-identified as heterosexual (93%), 8 (4%) were bisexual, and 5 (2.5%) were homosexual. 76/200 respondents (38%) reported some form of abuse at any point in time, 11/76 (14.4%) reported abuse over the preceding 12 months; 15 (20%) patients reported physical and/or sexual abuse, 19 (25%) reported verbal abuse, and 42 (55%) reported both physical and verbal abuse. 36 (47%) patients received professional help in dealing with the aftermath of the experience. 2/76 (3%) patients identified themselves and asked to be contacted by the health care team. Married women or those in domestic partnership were more likely to report any form of abuse than single patients ($P < .0207$). Any level of current neurologic disability was associated with a higher likelihood of ever experiencing verbal abuse ($P = .021$). There was no association between physical ($P = .0734$) or verbal abuse ($P = .363$) and neurologic disability using the χ^2 test. There was no association between these variables when adjusting for age and education based on a logistic regression model ($P = .860$). **Conclusions:** 38% of respondents acknowledged some form of abuse in their past. This proportion is higher than that reported in the general female population (29%). Less than 50% of affected women ever sought professional help in dealing with the trauma. Intimate partner violence in married or partnered women with MS may be more prevalent than previously recognized. Lower than expected anonymous questionnaire response rates suggest persistent patient discomfort in addressing this difficult matter. This study will continue recruiting subjects to increase the power of our observations.

Supported by: None

Disclosure: *Jeta Pol-Patil, Maria Claudia Manieri, Tatenda Mahlanza, Elizabeth Misasi, Laura T. Safar:* Nothing to disclose. *Bonnie Ilene Glanz:* Merck Serono, Verily Life Sciences (grant support). *Maria K. Houtchens:* Biogen, Celgene, Genzyme Sanofi, Mallinckrodt, Serono (consulting fee).

Keywords: *Physical and sexual violence, Psychological issues and MS*

(PSF04) Predictors for Self-Efficacy for People Living with Multiple Sclerosis

Joan Augustyn, Carly Robertson, Melissa Naylor, Alicia Pille

Occupational Therapy Education and MS Achievement Center, University of Kansas Medical Center, Kansas City, KS

Background: The Multiple Sclerosis Achievement Center (MSAC) is a community wellness rehabilitation program for individuals with progressed multiple sclerosis (MS). This site provides services to enhance quality of life and provide a sense of community for members. Services provided at this site include occupational therapy, physical therapy, social activities, and mental health groups. There is little evidence that focuses on predictors for self-efficacy in individuals with progressed MS in the literature. This research studied the effect of participation at MSAC in relation to feelings of self-efficacy from the perspective of individuals with MS by reviewing and analyzing data that are regularly collected on MSAC participants. **Objectives:** The objectives of this study were to analyze data on pain, fatigue, outside activities, and participation in program activities for members of the MSAC and analyze data on self-efficacy scale for participants of MSAC. **Methods:** This study looked at data collected as part of MSAC. These data included information from forms about levels of fatigue, pain level, medical concerns, outside activities, and social isolation that is collected each week from members. Scores from the Multiple Sclerosis Self-Efficacy Scale (MSSES), which consists of person-rated perception of ability to overcome challenges one is faced with, were also collected and analyzed. Sample size included 50 individuals (36 women, 14 men) who are members of the MSAC. This was an observational, cross-sectional design, and correlational analysis investigated the relationships between pain, fatigue, activity outside of the program, and participation in program activities and feelings of increased self-efficacy. **Results:** The results indicate no significant correlations between the total MSSE score and individual participant factors. However, results suggest that doing-based questions have a stronger relationship with self-efficacy ($\beta = .505$; $P < .001$) than feeling-based questions ($\beta = .411$; $P < .001$). When completing a correlational analysis of the total self-efficacy scores, doing-based questions also have a higher

correlation ($\beta = .640$) than feeling-based questions ($\beta = .577$) in comparison to total MSSE scores. **Conclusions:** The results indicate that both the ability to perform activities of daily living and feelings related to MS have an impact on levels of self-efficacy in this population. However, it suggests that the ability to perform tasks and activities of daily living had the most significant impact on participants' self-efficacy scores.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Psychological issues and MS, Wellness

(PSF05) The Association Between Health Literacy, Health Outcomes, and Medication Adherence in Patients with Multiple Sclerosis

Justin D. Williams,¹ Andrew Goodman,² Jessica F. Robb³

¹Neurology, University of Rochester School of Medicine & Dentistry, Rochester, NY;

²Neurology, University of Rochester, Rochester, NY; ³University of Rochester Medical Center, Rochester, NY

Background: Multiple sclerosis (MS) is an immune-mediated disorder characterized by demyelination of nerve cells within the brain and spinal cord over a period of time. Historically, MS was considered an untreatable disease; however, there are currently over a dozen disease-modifying therapies approved by the US Food and Drug Administration. A potential barrier to receiving a diagnosis and treatment of MS is health literacy, which is described as the degree to which individuals have the capacity to obtain, access, and understand basic health information and services needed to make appropriate decisions. **Objectives:** The purpose of this study is to investigate the relationships between health literacy, health outcomes, and medication adherence in patients with MS. **Methods:** This study was a single-site, prospective study done at the MS center at the University of Rochester Medical Center. Health literacy was measured using the Short Test of Functional Health Literacy in Adults. **Results:** Of the 179 subjects included in the analysis, 178 had adequate health literacy, 1 had marginal health literacy, and 0 had inadequate health literacy. **Conclusions:** The relationship between health literacy, health outcomes, and medication adherence cannot be determined in this sample given the lack of variability in health literacy.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Health literacy

(PSF06) It Takes a Village: The Veterans Health Administration (VHA) MS Centers of Excellence and National Multiple Sclerosis Society Partnership for Facilitating Communication, Collaboration, and Coordination of Services for Veterans with Multiple Sclerosis

Margaret Kazmierski,¹ Alicia Sloan,² Allison Krehbiel,³ Lynette Coleman,⁴ Doris Freire-Lill⁵

¹VA MS Centers of Excellence—East, Department of Veterans Affairs, VHA, Baltimore, MD;

²VA MS Centers of Excellence—West, Department of Veterans Affairs, VHA, Seattle, WA;

³MS Navigator Services, National Multiple Sclerosis Society, New York, NY; ⁴Advocacy, Services and Research, National Multiple Sclerosis Society, New York, NY

Background: Currently, there are 22 million veterans. Only 8 million of these veterans are enrolled with the Veterans Health Administration (VHA). 70% of these veterans receive additional health, support services, and care coordination from private sector providers outside of the VHA. The VA MS Centers of Excellence (MSCOE) supports comprehensive specialty care teams across the VHA who evaluate, treat, and provide ongoing care management to over 24,000 veterans with MS in the United States. MS Clinics across the VHA system provide comprehensive MS care through MS specialists including rehabilitation services, neurology, nursing, social work, and neuropsychology. In addition, VHA system provides home health aide services and caregiver support to assist veterans with MS remain independent in their community. **Objectives:** Through a collaborative partnership, VHA MSCOE and The National MS Society (NMSS) developed a formal process for mutual communication and coordination of resources for veterans with MS: 1) VA MSCOE social workers provide VHA 101 educational webinars to MS navigators who address unique needs of veterans with MS and their families, providers, and care partners. 2) To establish a process of case consultations that involve veterans who contact the MS navigators with complex resource, support, or benefit needs. 3) Identify veterans who could benefit from MS Navigator Program and send referrals from VHA. **Methods:** 1) The MSCOE Social Work Staff developed and provided training presentations (VHA 101) to educate MS navigators about veteran culture, VHA eligibility, enrollment, programs, and care navigation. 2) Point of contacts were established

for both the MSCOE and NMSS for discussing complex veteran cases. 3) Case consultations between MS navigators and MSCOE social work staff proceeded routinely. **Results:** Over 80 MS navigators participated in VHA 101 webinars provided by VHA MSCOE social work staff. Case consultations between MS navigators and VHA MSCOE social work staff were successfully resolved. Types of referrals between VHA and the MS Navigator Program were identified and increased including veterans' benefits, VA MS specialty care services, VA and NMSS funding. For example, VA Puget Sound referred 22 veterans to the MS Navigator Program for MS educational material, support groups, and financial assistance for veterans for bills, gym membership, driver's license, adaptive driving equipment not covered by VA, scooter lift installation, and bed bug eradication. **Conclusions:** Preliminary outcomes from educational trainings, individual case consultations, and the referral process have been effective between VHA/MSCOE and NMSS's MS Navigator Program. Education on how each organization operates and provides care and services for veterans has enhanced the level of information sharing and referrals, thus improving care. Additional presentations and trainings are being planned for both organizations.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, MS and the caregiver/family, Psychological issues and MS

(PSF07) Discussing Multiple Sclerosis (MS) Progression with Patients: Experiences of UK Health Care Professionals from the Spectrum Project

Martin Duddy,¹ Carmel Wilkinson,² Katherine Rhys³

¹The Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle, United Kingdom;

²Multiple Sclerosis, South Tyneside and Sunderland NHS Foundation Trust, Sunderland, United Kingdom; ³Novartis Pharmaceuticals UK Ltd, London, United Kingdom

Background: Receiving a diagnosis of secondary progressive multiple sclerosis (SPMS) can have a significant psychological impact on patients. Identifying how and when to initiate discussions about multiple sclerosis (MS) progression may be challenging for health care professionals (HCPs). **Objectives:** To understand how HCPs in the United Kingdom discuss the progression from relapsing-remitting MS (RRMS) to SPMS with patients. **Methods:** Interviews were conducted in 2019 with 59 HCPs from geographically dispersed UK MS centers (MS neurologist, $n = 41$; MS specialist nurse, $n = 15$; other HCP, $n = 3$) using a structured questionnaire. Topics covered included current practices for defining, diagnosing, and managing SPMS, and discussing SPMS with patients. This analysis focuses on discussing SPMS with patients. $n < 59$ indicates missing responses. **Results:** Progression from RRMS to SPMS is most commonly discussed with patients at the following time points (not mutually exclusive): when the SPMS diagnosis is confirmed ($n = 56/58$ [97%]), when a patient asks about SPMS after researching their condition ($n = 56/58$ [97%]), or when SPMS is first suspected ($n = 45/58$ [78%]). Only 20/58 HCPs (34%) discuss SPMS at initial RRMS diagnosis and 28/58 (48%) during the RRMS disease course. Most HCPs ($n = 43/57$ [75%]) reported that a neurologist is usually the first person to discuss progression with the patient. The most common terms used by HCPs when discussing SPMS with patients were "progression or progressive" ($n = 45/59$ [76%]), followed by "transition" ($n = 19/59$ [32%]), "worsening" ($n = 16/59$ [27%]), and "disability" ($n = 9/59$ [15%]). However, a number of HCPs reported that they would specifically avoid using the same terms [disability ($n = 13/59$ [22%]), progression or progressive ($n = 10/59$ [17%]), worsening ($n = 8/59$ [14%]), transition ($n = 4/59$ [7%])]. The median estimated time between first suspecting and diagnosing SPMS was 12.0 (IQR 12.0-24.0) months ($n = 45$). The most common explanations for reluctance to diagnose SPMS were concerns over withdrawing treatment ($n = 49/59$ [83%]) and psychological impact on patients ($n = 39/59$ [66%]). **Conclusions:** There is substantial variation in the United Kingdom in both how and when HCPs discuss the transition from RRMS to SPMS with patients. Discussions may be delayed until SPMS is suspected or even confirmed, which can take a year or more. Further training and support for HCPs may be needed to facilitate discussions with patients about MS progression and provide them with appropriate support during the transition phase.

Supported by: None

Disclosure: *Martin Duddy:* Novartis, MedDay (contracted research). *Carmel Wilkinson:* Novartis (consulting fee). *Katherine Rhys:* Novartis (salary).

Keywords: Progressive MS, Psychological issues and MS

(PSF08) Development and Implementation of a Patient Education and Cognitive Wellness Program for Veterans with Multiple Sclerosis

Terry Lee-Wilks,¹ Jessica Dalrymple,² Alicia Sharma,³ Jessica Stillman,² Amy Olzmann,⁴ Moira Dux⁴

¹MS Center of Excellence-East, VA Maryland Health Care System, Baltimore, MD; ²Neuropsychology, VA Maryland Health Care System, Baltimore, MD; ³Adult and Pediatric Medical Psychology, Life Bridge Health/Sinai Hospital, Baltimore, MD; ⁴VA Maryland Health Care System, Baltimore, MD

Background: Multiple sclerosis (MS) is a complex chronic disease that affects neurologic, psychiatric, and cognitive functions. Symptoms in these domains can adversely affect functioning and quality of life. Recent epidemiologic studies document higher MS incidence rates among military personnel compared to the general population. To date, no comprehensive group psychotherapeutic wellness intervention addressing the various factors affecting veterans with MS has been disseminated to our knowledge. **Objectives:** 1) To describe the development of a cognitive rehabilitation and psychotherapeutic wellness group intervention for veterans with MS; 2) To improve cognitive functioning in daily life and facilitate implementation of strategies for coping with cognitive, emotional, physical, and social challenges posed by MS. **Methods:** Content for an introductory 7-week group entitled MS Intervention and Development of Skills (MINDS) and an advanced, part 2, 7-week group entitled Master MINDS was adapted from various existing cognitive rehabilitation programs and tailored to veterans with MS. Sessions included psychoeducation regarding MS symptoms; compensatory strategies to address the cognitive domains often affected by MS (attention, memory, processing speed, executive functioning); methods for enhancing positive health behaviors, including strategies to cope with fatigue, depression, psychosocial stress, and social role changes; and discussion of relevant VA and community resources. Guest speakers from other disciplines in MS care were invited to facilitate 2 didactic and question-and-answer sessions. Importantly, veteran feedback was sought throughout each session and used to develop 2 new modules related to parenting with MS (Caring MINDS) and specific issues related to men with MS (Mr. MINDS). **Results:** Six Veterans participated in the groups and completed self-report questionnaires assessing mood (PHQ-9) and subjective cognitive impairment (MSNQ) pre- and postintervention. Examination of scores revealed that most reported stable or improved mood while half reported stable or lower risk of depression and/or cognitive impairment post-MINDS. Post-Master MINDS, most reported stable or improved mood and stable or lower risk of depression and/or cognitive impairment (1 participant missing). **Conclusions:** Group intervention for veterans with MS is a viable treatment modality, and content tailored to this population is subjectively useful in improving mood and increasing awareness of the cognitive, emotional, physical, and social challenges associated with MS. Therapeutic factors such as the instillation of hope, social support, validation of concerns, and interpersonal learning were identified by group members as key elements of the intervention. Finally, patient-centered feedback was critical in developing tailored treatment plans and additional modules.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Cognitive rehabilitation, Psychological issues and MS

(PSF09) The Effects of Customized Psychoeducation-Based Neurocounseling Interventions on the Coping Flexibility of African American Women with Multiple Sclerosis

Whitney McLaughlin

North Carolina State University, Raleigh, NC

Background: The importance of adaptive (ie, effective) coping strategies among people with multiple sclerosis (PwMS) has been well documented in the literature. However, a gap in the body of knowledge related to African American women living with this chronic disease still exists. Historically, their coping behaviors, emotional support needs, and mental health have not been a focus in the multiple sclerosis (MS) literature. As a result, the challenges, needs, and perspectives of black women with MS are limited and deserve more scholarly attention, and a great need exists to help this underserved and under-researched population with their coping efforts. This study examined the effectiveness of a Brain-Based Education and Wellness (BE WELL) intervention on the coping flexibility of African American women with MS. **Objectives:** 1) To learn about the conceptual framework for the BE WELL program. 2) To examine the effects of customized psychoeducation-based neurocounseling interventions on the coping flexibility of African American women with MS. 3) To assess participants' social validity ratings of the BE WELL intervention pro-

gram. **Methods:** An N-of-1/A-B-A single-case research design (SCRD) was used for this 12-week intervention study. Data were examined using both visual and statistical analysis. This involved using descriptive statistics including measures of central tendency and variability, autocorrelations, and regression analysis to look for trends. The G index was used to calculate effect sizes, and the nonparametric test Conservative Dual-Criteria (CDC) was used as a robust statistical analysis tool to compare the phases of each coping measure. **Results:** The participants were 4 African American women with MS ages 34 to 60. For 3 participants, "A," "C," and "T," there were large-to-medium effect sizes ($ES = 0.50$ to $ES = 1$) for 1 or more coping measures in the baseline to withdrawal phase. For participant 2, "Y," there was only a medium effect size for evaluation coping ($ES = 0.50$) from treatment to withdrawal phase. Participants' social validity ratings from the ATT ranged from 87-98 indicating that each participant found the intervention to be valuable. **Conclusions:** The customized BE WELL intervention seemed to have positive effects for each of the participants' coping flexibility. Participant 2 was the exception as only slight effects were observed. For 3 participants, the most profound effects occurred in evaluation coping. Results from the ATT validated the findings from the visual and quantitative analysis as all participants' ratings revealed that each participant experienced positive treatment effects from the BE WELL program.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Management of activities of daily living in MS, Psychological issues and MS

(PSF10) The Conformity of Masculine Norms and the Effects on Coping, Health Behaviors, and Quality of Life in Men with Multiple Sclerosis

Bryan Davis

Mellen Center for MS/Neurological Institute, Cleveland Clinic, Cleveland, OH

Background: The aim of the project is to explore how the conformity of masculine norms affects coping, health behaviors, and quality of life (QOL) in men with multiple sclerosis (MS). Despite some documented concerns of men with MS (eg, prognosis, QOL, targeted interventions) (Upton and Taylor, 2015), the specific conformity of socialized masculinity and gender norms affecting men with MS has not received attention in the academic literature. According to Levant and Wimer (2014), masculinity conformity is complex in nature due to finding some gender performance to be protective and others to be detrimental when it comes to men's health. It is unknown how masculinity conformity affects men with MS, specifically in terms of its association with coping, health behaviors, and QOL. This study aims to explore these factors to evaluate the experiences of men with MS. Such information will fill a void in the literature and inform future interventions to better serve men with MS. **Objectives:** Examine and describe the association between masculinity conformity, coping, health behaviors, and QOL in a representative sample of men with MS. 1. **Hypothesis:** Higher scores on the masculinity inventory will produce lower/worse scores on health-related QOL. Subaim: Investigate the differential effect of coping and masculinity conformity on QOL. 2. **Hypothesis:** Men with higher scores on the masculinity inventory will have less/worse coping, which will negatively impact QOL more than for men with lower scores on the masculinity inventory (independent of coping ability). 3. **Hypothesis:** Higher scores on the masculinity inventory will produce lower/worse scores on health-related QOL. Subaim: Investigate the differential effect of coping and masculinity conformity on health behaviors. **Methods:** Participants in this study will include adult (18+ years) men with MS who are established patients at the Mellen Center (Cleveland Clinic). Demographic information (age, race, marital status, household income, education status) and clinical characteristics (years since MS diagnosis, Patient-Determined Disease Steps), and 5 patient-reported outcomes will be collected: The Conformity of Masculinity Inventory-46 which is a short version of the CMNI (Mahalik et al, 2003) will be used to assess the conformity to 9 masculine norms. Health-related QOL will be assessed with the PROMIS Global Health (Hays 2009). 1) The Ways of Coping Questionnaire (Lazarus and Folkman, 1985) will be used to measure coping. 2) Health behaviors will be measured using the Health Behaviors Inventory-20. 3) Disease impact will be evaluated using MS Performance Scales. **Results:** Research initiated January 2020. Results expected in spring of 2020. **Conclusions:** Research initiated January 2020. Results expected in spring of 2020.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Gender Issues, MS and the caregiver/family, Psychological issues and MS

(PSF11) Understanding the Lived Experience of Health Through the Exploration of Well-Being of Women with Multiple Sclerosis: Preliminary Findings

Jennifer L. Howard,¹ Yolanda Babenko-Mould,¹ Kimberley Jackson,¹ Tracy Smith-Carrier²

¹Health Science–Nursing, Western University, London, ON, Canada; ²School of Social Work, King's University College at Western University, London, ON, Canada

Background: There are approximately 100,000 Canadians with multiple sclerosis (MS), a condition that is 3 times more likely to affect women compared to men. Existing evidence proposes that women may experience disability, social determinants of health, and their overall health differently than men. Therefore, current evidence may benefit from researchers exploring gender as a category for investigation of phenomena related to health. The World Health Organization defines health as physical, mental, and social well-being and is underpinning the exploration of health in women with MS for this study. Limited evidence is presently available as to how women in Canada with MS experience their health through physical, social, and mental well-being. **Objectives:** The purpose of this hermeneutic phenomenological study is to understand the essence of the lived experiences of women affected by MS living in Southwestern Ontario, Canada, for health in the context of physical, social, and mental well-being. **Methods:** van Manen's hermeneutic phenomenological approach was used to explore the experience of health through the context of physical, social, and mental well-being for participants. Participants included 20 women affected by MS living in Southwestern Ontario, Canada, who were interviewed using semistructured interview guides that were audio-recorded, transcribed verbatim, and analyzed using van Manen's interpretive phenomenological approach, which consisted of extracting significant statements and themes for the investigated phenomena. **Results:** Preliminary results from this study will be presented at this conference as to how women experience health through their well-being and will be discussed in an oral presentation. **Conclusions:** A preliminary review of the findings of health through the context of social well-being, such as barriers or factors that promote the overall health and well-being for women affected by MS, will be presented. Further exploration as to how social well-being and various social determinants of health affect health through the context of physical, social, and mental well-being is needed to further extrapolate these preliminary findings of the study.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Nursing management in MS, Self care and MS

QUALITY OF LIFE AND OUTCOMES

(QOL01) Alemtuzumab Effects on Urogenital Function: Results Pooled From the CARE-MS 9-Year Functional Assessment of Multiple Sclerosis Quality-of-Life Survey

Aaron Boster,¹ Rafael Arroyo,² Antonio Bertolotto,³ Samuel F. Hunter,⁴ Carolina Ionete,⁵ Bart Van Wijmeersch,⁶ Ericka M. Bueno,⁷ Nadia Daizadeh,⁷ Elizabeth M. Poole,⁷ Bhupendra O. Khatri⁸

¹Boster MS Center, Columbus, OH; ²Hospital Universitario Quironsalud, Madrid, Spain; ³SCDO Neurologia & CRESM, AOU San Luigi, Orbassano, Turin, Italy; ⁴Advanced Neurosciences Institute, Franklin, TN; ⁵University of Massachusetts Memorial Medical Center, Worcester, MA; ⁶Faculty of Medicine and Life Sciences, Hasselt University, Hasselt, Belgium; ⁷Sanofi, Cambridge, MA; ⁸Center for Neurological Disorders at Wheaton Franciscan Healthcare, Milwaukee, WI

Background: Nearly all patients with multiple sclerosis (MS) experience urogenital dysfunction within 10 years, resulting in significant impacts on quality of life (QoL). Few studies have measured the effects of disease-modifying therapies (DMTs) on this aspect of disability. In the CARE-MS trials (trial registration: NCT00530348, NCT00548405), alemtuzumab showed greater improvements in clinical, magnetic resonance imaging, and QoL outcomes vs subcutaneous interferon beta-1a (SC IFNβ-1a) over 2 years in patients with relapsing-remitting MS, and efficacy was maintained in 2 consecutive extension studies (NCT00930553, NCT02255656). **Objectives:** Determine the effect of alemtuzumab treatment on urogenital function over 9 years in pooled CARE-MS patients. **Methods:** Patients received either SC IFNβ-1a 44 µg 3x/week or 2 alemtuzumab courses (12 mg/day; baseline: 5 consecutive days; 12 months later: 3 consecutive days) in the core studies, with additional alemtuzumab as needed in the extensions (3 consecutive days, ≥12 months after the most recent course) or other DMTs. Items on the Functional Assessment of MS (FAMS) questionnaire pertaining to urination frequency, bladder control, and sexual function (for patients who had been sexually

active within 1 year) were assessed. Patients rated their perceptions on a 0- to 4-point scale and considered the previous 7 days in their responses. **Results:** In the alemtuzumab group (n = 798), mean urination frequency rating was 1.04 at baseline, improved to 0.93 at year 2 (P = .002 vs baseline), and was 0.96 at year 9 (P = .94); 73% of responses at year 9 rated at 0 or 1 (indicating normal frequency). Mean urination frequency rating in SC IFNβ-1a patients (n = 386) was 1.11 at baseline and 1.04 at year 2 (P = .01 vs alemtuzumab at year 2). Mean urine control rating in alemtuzumab patients was 0.91 at baseline, 0.86 at year 2 (P = .28 vs baseline), and 0.93 at year 9 (P = .035); 76% of responses at year 9 rated at 0 or 1 (indicating good control). SC IFNβ-1a patients rated urine control as 0.88 at baseline and 0.84 at year 2. FAMS urine control ratings and Expanded Disability Status Scale bowel/bladder functional system scores in the alemtuzumab group were highly correlated (Spearman coefficient at year 2, 0.5695; P < .0001). Mean sexual satisfaction rating for alemtuzumab patients was 2.50 at baseline, 2.57 at year 2 (P = .20 vs baseline), and 2.62 at year 9 (P = .56); >60% of responses at year 9 rated at 3 or 4 (indicating high satisfaction). Mean sexual satisfaction rating in SC IFNβ-1a patients was 2.44 at baseline and 2.45 at year 2 (P = .047 vs alemtuzumab at year 2). **Conclusions:** Alemtuzumab-treated patients' perception of urine frequency and bladder control and satisfaction with sexual function were stable over 9 years after initiating alemtuzumab.

Supported by: Sanofi, Bayer HealthCare Pharmaceuticals

Disclosure: Aaron Boster: Biogen, Mallinckrodt, Medtronic, Novartis, Sanofi Genzyme, Teva (consulting fees and/or fees for non-CME services); Sanofi (advisor for participating on the managed care review panel). Rafael Arroyo: Almirall, Bayer, Biogen, Merck, Novartis, Roche, Sanofi, Teva (speaking fees, advisory board). Antonio Bertolotto: Biogen (consulting fee, lecture fees); Genzyme (consulting fee); Merck, Novartis, Roche, Sanofi, Teva (lecture fees). Samuel F. Hunter: AbbVie, Acorda, Actelion, Adamas, Alkermes, Avanir, Bayer HealthCare, Biogen, Genzyme, Novartis, Osmotica, Questcor, Roche, Sanofi, Synthron, Teva (consulting agreements, speaker honoraria, grant/research support). Carolina Ionete: Biogen, Roche (contracted research); Sanofi (compensation for advisory board participation, contracted research). Bart Van Wijmeersch: Actelion, Bayer Schering, Biogen, Merck Serono, Novartis, Roche, Sanofi, Teva (research and travel grants, honoraria for MS-expert advice, speaking fees). Ericka M. Bueno, Nadia Daizadeh, Elizabeth M. Poole: Sanofi (salary). Bhupendra O. Khatri: Acorda, Celgene, Serono, Teva (consulting/honorarium [consulting work, and speaker programs]); Alexion, Biogen, Genentech, Novartis, Sanofi (consulting/honorarium [consulting work, speaker programs], contracted research); Ra Pharmaceuticals (contracted research).

Keywords: Disease-modifying treatments in MS, Management of activities of daily living in MS

(QOL02) The Impact of Relapses on Quality of Life in Patients with Neuromyelitis Optica Spectrum Disorder: Data from the Phase 3 PREVENT Study

Achim Berthele,¹ Michael Levy,^{2,3} Karissa Johnston,⁴ Meagan Harwood,⁴ Adrian Kiehlhorn,⁵ Mynying Royston,⁵ Guido Sabatella,⁵ Jacqueline Palace⁶

¹Technical University of Munich, Munich, Germany; ²Johns Hopkins University, Baltimore, MD; ³Massachusetts General Hospital and Harvard Medical School, Boston, MA; ⁴Broadstreet HEOR, Vancouver, BC, Canada; ⁵Alexion Pharmaceuticals, Boston, MA; ⁶John Radcliffe Hospital, Oxford, United Kingdom

Background: Data are lacking on the impact of relapses on patients with neuromyelitis optica spectrum disorder (NMOSD). PREVENT (trial registration: NCT01892345) was a randomized, double-blind study of eculizumab in patients with aquaporin-4 immunoglobulin-G-positive NMOSD, which showed a 94% reduction in the risk of adjudicated relapse vs placebo; it also included survey assessments of health-related quality of life (QOL). **Objectives:** To evaluate the effect of relapse on QOL in patients with NMOSD, using data from the phase 3 PREVENT study. **Methods:** Patients' health-related QOL was assessed using the EuroQol 5-Dimensions questionnaire (EQ-5D) and the Medical Outcomes Study Short-Form (36-item) Health Survey (SF-36), with higher values in both indicating better health-related QOL. In the current post hoc analysis, data from the eculizumab and placebo groups were pooled and the last recorded EQ-5D and SF-36 scores before an adjudicated relapse were compared (at relapse level) with postrelapse scores (recorded ≥30 days after relapse) using a paired t test. **Results:** In the absence of relapse, EQ-5D and SF-36 scores were stable over time, as expected. Mean scores before (n = 24) and after (n = 22) relapse were EQ-5D index (possible range 0-1): 0.656 and 0.595, respectively (pre-post difference, -0.067; P = .012); EQ-5D visual analog scale score (possible range 0-100): 60.458 and 56.500, respectively (pre-post difference, -4.227;

$P = .226$); SF-36 physical component summary score (possible range 0-100): 38.487 and 36.431, respectively (pre-post difference, -2.602 ; $P = .049$); and SF-36 mental component summary score (possible range 0-100): 45.300 and 41.774, respectively (pre-post difference, -3.164 ; $P = .041$). There were significant differences between pre- and postrelapse scores in the SF-36 domains of bodily pain ($P = .027$), physical functioning ($P = .007$), role-emotional ($P = .021$), and vitality ($P = .025$). **Conclusions:** This analysis suggests that relapses in patients with NMOSD are associated with a significant reduction in certain aspects of QOL beyond the immediate relapse period.

Supported by: None

Disclosure: *Achim Berthele:* Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent, speakers' bureau). *Michael Levy:* Alexion Pharmaceuticals (consulting fee, contracted research, fees for non-CME/CE services received directly from commercial interest or its agent); Genentech, Quest Diagnostics, Viela Bio (consulting fee, fees for non-CME/CE services received directly from commercial interest or its agent). *Karissa Johnston, Meagan Harwood:* Nothing to disclose. *Adrian Ktelborn, Minying Royston:* Alexion Pharmaceuticals (salary). *Guido Sabatella:* Alexion Pharmaceuticals (ownership interest, salary). *Jacqueline Palace:* Biogen, Chugai (contracted research); LEK, Viela Bio, Guidepoint (consulting fee); Merck Serono (meeting/lecture/workshop participation); Novartis, Roche, Argenx (speakers' bureau); UCB, Viela Bio, Roche (conference/lecture participation).

Keywords: Quality of life in NMOSD

(QOL04) Understand Common Multiple Sclerosis Symptoms Experienced Among MS Patients Participating in an Online MS Community

Beth Schneider

MyHealthTeams, San Francisco, CA

Background: People with multiple sclerosis (MS) often seek perspectives and support from each other by connecting on patient social networks. These organic interactions provide unfiltered, rich insight into the day-to-day challenges and needs of patients, including both the emotional and physical issues they face. Directly understanding the holistic impact of MS on patients is crucial to treating patients, including improving doctor-patient interactions and enabling patients to better manage their MS. **Objectives:** Leverage the largest MS patient social network in the world to understand the most prevalent symptoms of patients with MS as they reach out to one another for information and support. **Methods:** Research conducted on de-identified organic discussions within MyM-Team.com, a social network of >127,000 people with MS in the United States. Using natural language processing, 178,884 verbatim discussions from April-September 2019 were analyzed. Key themes were identified and used to determine common symptoms and sentiment. **Results:** 40% of the discussions were about symptoms, which were highly negative (60%). Most prevalent discussion was pain (35% of symptom discussions), especially leg pain. While members turned to medicines such as gabapentin, they also sought out less traditional approaches like CBD in search of pain relief. Other common symptoms discussed included mobility issues and fatigue, but also the emotional impact of MS (depression and anxiety). While slowing MS progression is being addressed by their health care provider, MS symptoms were often not being treated, especially depression and anxiety. **Conclusions:** Understanding the physical and emotional symptoms that accompany MS but that are not always shared between patients and their doctors can help health care providers provide a more holistic approach to treating patients with MS. This includes helping patients with MS understand what symptoms they are likely to experience and how they can mitigate them.

Supported by: None

Disclosure: *Beth Schneider:* MyHealthTeams (contracted research).

Keywords: Comprehensive care and MS, Management of activities of daily living in MS, Psychological issues and MS

(QOL05) Uncovering the Needs and Gaps in Care Among Multiple Sclerosis Patients Participating in an Online MS Community

Beth Schneider

MyHealthTeams, San Francisco, CA

Background: People with multiple sclerosis (PwMS) often seek perspectives and support from each other by connecting on patient social networks. These organic interactions provide unfiltered, rich insight into the day-to-day challenges and needs of patients that are not always raised with health care providers (HCPs). They also uncover attitudes towards

and perceptions of disease-modifying therapies that sometimes are a surprise to HCPs. Directly understanding the key challenges, concerns, and habits of PwMS can elevate doctor-patient interactions and enable patients to better manage their multiple sclerosis (MS). **Objectives:** Leverage MS social network to obtain key insights, and understand key issues and challenges of patients with MS that could be helpful to HCPs treating people with MS. **Methods:** Research conducted with de-identified organic verbatims within MyMTeam.com, a social network of >127,000 PwMS in the United States. Using natural language processing, 178,884 verbatim discussions from April-September 2019 were analyzed. Key themes were identified and used to determine positive or negative sentiments. **Results:** 40% of the discussions were about symptoms with pain being the most discussed, 15% about the doctor-patient relationship, 10% disease-modifying treatments, and 6% on symptomatic medications. Discussions revealed that depression and anxiety that accompany MS are often undertreated and that pain can be relentless, especially leg pain. Approximately 18% of the conversations focused on supporting one another. Social conversations were extremely positive and well received ("I feel less isolated since joining this team."), while conversations focused on living with MS were generally negative especially the profound impact of MS on quality of life. **Conclusions:** The organic interactions among patients with MS provided deeper understanding of symptoms, challenges, attitudes to treatment, and steps taken to mitigate MS in ways that are not often shared between patients and doctors. Understanding the needs of these patients provides significant opportunities for HCPs to better support and educate their patients. This includes setting the right expectations; addressing the impact of MS holistically, including pain, depression and other symptoms; and providing tools and education materials to help stay the course. The research also suggests that encouraging patients with MS to connect with one another can help alleviate some of the isolation and depression.

Supported by: None

Disclosure: *MyHealthTeams* (contracted research).

Keywords: Comprehensive care and MS, Management of activities of daily living in MS, Psychological issues and MS

(QOL06) System-Level Variation in All-Cause Hospitalizations in Multiple Sclerosis: Year-1 Results of the Multiple Sclerosis Continuous Quality Improvement (MS-CQI) Research Collaborative

Brant J. Oliver,^{1,2,3,4} Fal S. Mehta⁵

¹Geisel School of Medicine at Dartmouth, Lebanon, NH; ²Dartmouth Hitchcock Medical Center, Lebanon, NH; ³The Dartmouth Institute, Hanover, NH; ⁴Geisel School of Medicine, Hanover, NH; ⁵Community & Family Medicine, Dartmouth, Lebanon, NH

Background: Multiple Sclerosis Continuous Quality Improvement (MS-CQI) is the first randomized, multicenter, prospective, longitudinal, systems-level, improvement science research study for multiple sclerosis (MS). MS-CQI is a 3-year study of system-level variation in performance outcomes, and leverages benchmarking to inform improvement using an informatics-enabled learning health system approach. MS-CQI collects 11 clinical electronic health record (EHR) outcome measures longitudinally, including MS treatment, all-cause emergency department utilization, and all-cause hospitalizations. We also collect demographic information and comorbidities. **Objectives:** To describe year-1 (baseline/preintervention) findings on system-level variation in selected clinical outcomes for individual sites, between sites, and for MS-CQI collectively. **Methods:** Four US MS centers are participating in MS-CQI: an urban academic center, a rural academic center, a rural community hospital, and a large urban private practice (total N = 5000 persons with MS). We collected approximately 7500 clinical measures abstracted from EHR data in year 1 from nearly 3000 clinical encounters. Demographic characteristics and longitudinal variation in measures did not vary significantly between sites. Encounter volume between centers was similar. We used analysis of variance, multiple regression, and maximum likelihood estimation methods to conduct inferential analyses. **Results:** Univariate analyses found significant differences ($P < .05$) between sites for multiple clinical outcomes including exacerbations, disease-modifying treatment, magnetic resonance imaging utilization, emergency department utilization, and hospitalizations. Controlling for individual level factors, including comorbidities, significant site (system)-level effects (with high-performing center specified as the referent group) were found for all-cause hospitalizations, with comparator sites demonstrating odds ratios ranging as high as 2.4 (95% CI: 1.34, 4.4). **Conclusions:** We found that significant geographic system-level variation in MS outcomes exists for all-cause hospitalizations for people with MS followed by participating MS-CQI centers. Findings suggest that a focus on system-level variation and improvement may be needed to reduce all-cause hospitalizations for people with MS.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Epidemiology of MS, System-level variation

(QOL07) Relapse Rate Is Influenced by System-Level Variation: Year-2 Results of the Multiple Sclerosis Continuous Quality Improvement (MS-CQI) Research Collaborative

Brant J. Oliver,¹ Karen Walsh,² for the MS-CQI Investigators

¹Geisel School of Medicine, Hanover, NH; ²College of Population Health, Thomas Jefferson University, Philadelphia, PA

Background: Multiple Sclerosis Continuous Quality Improvement (MS-CQI) is the first multicenter improvement science research collaborative for multiple sclerosis (MS), and includes a systems-level study of variation in MS outcomes. MS-CQI is a 3-year study that leverages benchmarking results to inform system-level improvement efforts targeting clinical outcomes using an informatics-enabled learning health system approach. **Objectives:** Here we present relapse rate results for year 1 (baseline/preintervention) compared to year 2 (first year of intervention). We also describe system-level variation in relapse rate for individual sites, between sites, and for MS-CQI collectively. **Methods:** We collect administrative data and 11 clinical electronic health record (EHR) outcome measures longitudinally across 4 clinical MS care centers in the United States. We conduct statistical process control analyses for benchmarking. Logistic regression and maximum likelihood estimation methods are used for inferential analyses. **Results:** Four US MS centers are participating: an urban academic center (n = 1000), a rural academic center (n = 1000), a rural community hospital (n = 1500), and an urban private practice (1500), following a total N = 5000 persons with MS. We have collected approximately 7200 clinical encounter measures from EHR data in year 1 and 10,000 in year 2. Demographic characteristics and longitudinal variation in measures did not vary significantly between sites. For year 1, center-specific proportions of persons with MS with at least 1 relapse ranged from 5%-16.9%. Mean relapse rate varied significantly ($P < .01$) across all centers. SPC analyses demonstrate a MS-CQI reduction of relapse rate from 11.5% (year 1) to 4.3% (year 2). Two sites were below the MS-CQI average of 7% (3.3%, 6.3%), and 2 were above the average (8.5%, 10.3%). Controlling for individual factors and covariates, logistic regression analyses identified significant center-level effects on relapse rate in year 1, with comparator sites demonstrating odds ratios as high as 2.61 (95% CI: 1.8, 3.8). **Conclusions:** MS-CQI has observed a significant reduction in population-level relapse rate by 7.2% during the first year of quality improvement intervention. We also found significant geographic system-level variation in MS relapse, suggesting that a focus on system-level variation and improvement may be needed to optimize outcomes.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Improvement science in MS

(QOL09) Living with Secondary Progressive Multiple Sclerosis: Results from an MS Coalition Survey

Hollie Schmidt,¹ Bari Talente,² Amanda Montague,³ Emily Willingham⁴

¹Accelerated Cure Project for MS, Waltham, MA; ²National MS Society, Washington, DC; ³Multiple Sclerosis Association of America, Cherry Hill, NJ; ⁴San Francisco, CA

Background: Many people diagnosed with relapsing multiple sclerosis (MS) will eventually transition to secondary progressive MS (SPMS). Until recently, only mitoxantrone had been approved by the US Food and Drug Administration as a disease-modifying therapy (DMT) for this MS subtype. In 2018, however, the agency accepted an application for siponimod for treatment of SPMS. In anticipation of this drug's approval, the MS Coalition, a network of 9 independent MS organizations, surveyed people with SPMS to understand how the disease affects their lives and their experiences with and attitudes about DMTs. The survey was developed with the Institute for Clinical and Economic Review (ICER), which at the time was conducting a review of siponimod for use in SPMS. **Objectives:** 1) To understand the challenges that people with SPMS face, including impacts on quality of life. 2) To learn about current use of DMTs among people with SPMS and their perspectives on future DMTs that could be developed for SPMS. **Methods:** Representatives of MS Coalition member organizations and ICER developed this web-based survey, which was then disseminated by MS Coalition members whose constituents include people with MS. **Results:** A total of 2263 respondents completed the full survey, and 2966 answered at least 1 question. Of those who participated in the survey, 51% reported being unable to work because of disability, 69% needed help with activities of daily living, and 86% used a mobility aid.

About three-fourths chose "fewer available treatment options" (78%) and "decline in quality of life" (72%) as impacts associated with their form of MS. Regarding use of DMTs, 37% reported no current use, and 22% indicated that they were taking Ocrevus. The other DMTs included in the survey were each used by fewer than 10% of respondents. When asked to consider a hypothetical new drug for SPMS, respondents expressed broad interest in possible benefits the drug could provide, including prevention of brain atrophy and improvement in SPMS symptoms. Side effects and long-term risks topped the list of reasons respondents gave for why they might not stay on a new drug. In responses to open-ended questions about impacts on daily life and family, common themes related to isolation, burden on family members, stress, and loss of mobility. **Conclusions:** Our respondents with SPMS described experiencing profound challenges, and their information provides insights into target areas of unmet need warranting continued therapeutic development.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Disease-modifying treatments in MS, Management of activities of daily living in MS, MS and the caregiver/family

(QOL10) Corticosteroid Pulse Therapy-Hyperglycemia: Protocol for Capillary Blood Glucose Control

Ivone R. Fernandes, Ana C. Sena, Maria A. Paschoal, Adakson M. Franco

Santa Casa of Misericórdia, São Paulo, Brazil

Background: Pulse therapy is the preferred therapy for treatment of multiple sclerosis (MS) outbreaks, which is a neurologic, demyelinating, and inflammatory condition, with intermittent periods of outbreak-remission. The recommendation is corticosteroid doses (500 mg to 1 g), every 3-5 days, administered at the hospital, as an inpatient, or as outpatient at an infusion clinic. The treatment reduces the inflammation during the outbreak phase of the MS and seeks to stabilize the crisis. Despite their adverse effects, the glucocorticoids are potent anti-inflammatories in the treatment of autoimmune pathologies. The glucocorticoids block the entrance of glucose in the tissues and increase the proteolysis, decreasing their synthesis in muscles, skin, bones, connective tissue, fat cells, and lymphoid tissue. Before the pulse therapy, it is important to eliminate the possibility of an active infection and to always administer the antiparasitic to control possible infestations. Checking the blood pressure, body weight and capillary glucose are very important during the infusion. Daily checks pre- and postinfusion are required as hyperglycemia may occur as an adverse effect of the therapy. The capillary blood glucose check is a blood test that gives immediate results about glucose concentration in the capillaries and the digital pulp. **Objectives:** To describe a protocol for puncture site rotation for the capillary blood glucose test performed during pulse therapy. **Methods:** A rotation pattern was established for the digital pulp punctures. The patient is directed to properly wash the hands and dry well. The nurse professional punctures the selected site on the right or left side of the distal phalanx of the finger chosen for the test, by alternating the puncture sites, and records the date, time, location and blood glucose value. **Results:** Not applicable. **Conclusions:** The introduction of a protocol for puncture site rotation, based on a simple code, assists in the communication between nursing professionals and promotes patient safety. The protocol allowed the participation and cooperation of the client thus establishing self-care.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Hyperglycemia, Nursing management in MS

(QOL11) Natalizumab Is Associated with Improvement in Cognitive Processing Speed and Health-Related Quality of Life: STRIVE 4-Year Results

Jai Perumal,¹ Roumen Balabanov,² Laura Balcer,³ Steven Galetta,³ Paula Read,⁴ Kuangnan Xiong,⁵ Denise Campagnolo,⁵ Christophe Hotermans,⁵ Lily Lee,⁵ Danette Rutledge,⁵ Robert J. Fox⁷

¹Weill Cornell Medical College, Cornell University, New York, NY; ²Northwestern University, Chicago, IL; ³New York University School of Medicine, New York, NY; ⁴Biogen, Maidenhead, MA, United Kingdom; ⁵Biogen, Cambridge, MA; ⁶Biogen (at time of this analysis), Cambridge, MA; ⁷Mellen Center for Multiple Sclerosis, Cleveland Clinic, Cleveland, OH

Background: Multiple sclerosis (MS) negatively affects cognitive function and quality of life (QoL), interfering with a patient's ability to work, pursue leisure activities, and perform activities of daily living. Natalizumab is a highly effective treatment for patients with relapsing-remitting MS (RRMS) and has been associated with improved cognitive function

and QoL. **Objectives:** To examine 4-year, end-of-study cognitive processing speed and patient-reported QoL outcomes for natalizumab-treated patients with early RRMS. **Methods:** STRIVE is a completed, 4-year, multicenter, observational, open-label, single-arm study of anti-JC virus antibody-negative patients starting natalizumab <3 years after RRMS diagnosis. Cognitive processing speed was assessed by the Symbol Digit Modalities Test (SDMT). QoL was assessed via the patient-reported Multiple Sclerosis Impact Scale (MSIS-29). Outcomes were assessed annually. Changes from baseline were analyzed via a Wilcoxon signed rank test. **Results:** At baseline, patients in the intention-to-treat (ITT) population (N = 222) had active disease with a mean (SD) of 1.4 (1.2) relapses in the prior year. Baseline mean (SD) SDMT score was 52.1 (14.0). ITT patients showed improvements in SDMT score from baseline to year 1 (n = 191; mean change from baseline [CFB]: 2.29 [95% CI 0.84, 3.73]; P = .0006) with clinically significant improvements (ie, an increase ≥ 4 points) in year 2 (n = 158; mean CFB: 4.3 [95% CI 2.4, 6.2]; P < .0001) and year 4 (n = 174; mean CFB: 4.6 [95% CI 2.9, 6.2]; P < .0001). Patients with SDMT data at all time points (N = 145) had clinically significant SDMT improvements in years 2-4 (mean CFB [95% CI]: year 2, 4.4 [2.5, 6.2]; year 3, 4.3 [2.9, 5.8]; year 4, 5.1 [3.4, 6.7]). ITT patients also showed significant improvements in MSIS-29 score starting in year 1 (mean CFB [95% CI]: physical, n = 186; -3.9 [-5.8, -2.1]; P < .0001; psychological, n = 186; -1.7 [-2.8, -0.6]; P = .0012) that were sustained through year 4 (mean CFB [95% CI]: physical, n = 174; -4.7 [-6.9, -2.4]; P < .0001; psychological, n = 172; -2.6 [-3.9, -1.4]; P < .0001). **Conclusions:** Natalizumab treatment over 4 years was associated with a clinically significant improvement in cognitive processing speed as measured by the SDMT as well as improvement in patient-reported physical and psychological health. The potential for long-lasting (up to 4 years) improvement supports natalizumab's effectiveness in patients with early RRMS.

Supported by: Biogen

Disclosure: *Jai Perumal:* Acorda, Biogen, Genzyme, Teva (consulting fee). *Roumen Balabanov:* Biogen (consulting fees; grant/research support); Sanofi, Teva (consulting fee). *Laura Balcer:* Biogen, Genzyme (consulting fee). *Steven Galetta:* Biogen (consulting fee). *Paula Read, Kuangnan Xiong, Denise Campagnolo, Christophe Hotermans, Danette Rutledge:* Biogen (employee and may hold stock and/or stock options). *Lily Lee:* Biogen (employee at time of this analysis, may hold stock). *Robert J. Fox:* Actelion (consulting fees; advisory board participant); Biogen, Novartis (consulting fees; advisory board participant; clinical trial contract, research grant funding); EMD Serono, Genentech, Teva (consulting fee).

Keywords: Disease-modifying treatments in MS, Psychological issues and MS

(QOL12) Multiple Sclerosis Patient Perspectives: Disease Education and Communication Needs

Jessica O'Neill,¹ Mavis G. Ayer,² Samantha R. Colhoun,³ Nicola Daykin,⁴ Brenda Hamill,⁵ Simone Gabriele,⁶ Jordanne Florio,¹ Serena Pulcini,¹ Alison Thomson⁷

¹Roche Products Ltd, Welwyn Garden City, United Kingdom; ²Multiple Sclerosis, University Hospital Southampton Trust, Southampton, United Kingdom; ³Neuro, Queen Elizabeth Hospital Birmingham, Birmingham, United Kingdom; ⁴Nottingham City Care, Nottingham, United Kingdom; ⁵Multiple Sclerosis, Northern Health and Social Care Trust, Belfast, United Kingdom; ⁶Ipsos MORI, London, United Kingdom; ⁷Blizard Institute, Queen Mary University, London, United Kingdom

Background: Multiple sclerosis (MS) is a progressive neurologic disease that can significantly affect quality of life (QoL). In recent years there has been a shift towards increased patient engagement to inform person-centered care. It is therefore important that people with MS (PwMS) are equipped with the knowledge needed for informed, shared decision making and maintaining independence outside of a formal care setting. **Objectives:** This study aimed to identify unmet disease education and communication needs in PwMS to empower informed decisions, enable self-management, and to maintain independence for as long as possible. **Methods:** In October 2018 a roundtable of patient representatives, PwMS, carers, and MS nurses agreed on key themes associated with maintaining independence. In 2019, an official project steering group formed, and 2 studies for PwMS were co-developed: a qualitative Online Patient Community activity and a quantitative online survey. The qualitative activity used Ipsos' Syndicated MS Online Patient Community (a consistent panel of PwMS), and the quantitative survey was recruited through the MS Trust monthly newsletter and Facebook group. Results were discussed and prioritized by the steering group. **Results:** Data were analyzed from 28 and 117 respondents with relapsing-remitting MS from the Ipsos' Syndicated MS Online Patient Community and quantitative survey, respectively. Maintaining independence was a key goal in the qualitative findings, however only 33% reported they felt that they could "plan for the long term," and 21% said they set short-term/long-term goals for managing their MS. Overall, 66% "strongly/somewhat" agreed MS had

prevented them from reaching their full potential and had a "very/slightly negative" impact on QoL in terms of work (73%), social life (69%), and caring for family (57%). Data from this study will explore the relationship between perceived MS knowledge, planning for the future, and impact on QoL. **Conclusions:** There is evidence of a negative impact of MS on QoL among participants of our study. This highlights the importance of understanding unmet needs in education, linking disease progression and impact on future QoL in the wider MS population.

Acknowledgments: This project was sponsored by Roche Products Ltd. Ipsos MORI UK Ltd provided support in developing and conducting the surveys and performing data analysis, which was funded by Roche Products Ltd.

Disclosure: *Jessica O'Neill:* Roche Products Ltd (works for). *Mavis G. Ayer:* Biogen, Roche (sponsored delegate); Celgene, Merck, Novartis, Sanofi (consulting fee); Teva (sponsored education). *Samantha R. Colhoun, Alison Thomson:* Biogen, Novartis, Roche Products Ltd (consulting fee). *Nicola Daykin:* Roche Products Ltd (consulting fee). *Brenda Hamill:* Biogen, Novartis (sponsored delegate); Roche (consulting fee). *Simone Gabriele:* Roche Products Ltd (works for Ipsos MORI, which was funded by Roche to undertake this research). *Jordanne Florio:* Roche Products Ltd (works for). *Serena Pulcini:* Roche Products Ltd (works for).

Keywords: Comprehensive care and MS, Management of activities of daily living in MS, Patient education

(QOL13) Multiple Sclerosis Patient Perspectives: Impact on Employment

Jessica O'Neill,¹ Mavis G. Ayer,² Samantha R. Colhoun,³ Nicola Daykin,⁴ Brenda Hamill,⁵ Maria Fei,⁶ Jordanne Florio,¹ Serena Pulcini,¹ Alison Thomson⁷

¹Roche Products Ltd, Welwyn Garden City, United Kingdom; ²Multiple Sclerosis, University Hospital Southampton Trust, Southampton, United Kingdom; ³Neuro, Queen Elizabeth Hospital Birmingham, Birmingham, United Kingdom; ⁴Nottingham City Care, Nottingham, United Kingdom; ⁵Multiple Sclerosis, Northern Health and Social Care Trust, Belfast, United Kingdom; ⁶Ipsos MORI, London, United Kingdom; ⁷Blizard Institute, Queen Mary University, London, United Kingdom

Background: Multiple sclerosis (MS) is a progressive neurologic disease that can significantly impact quality of life (QoL). MS is typically diagnosed between the ages of 20 and 40, affecting individuals during critical employment years. Continued employment in people with MS (PwMS) has been shown to be associated with higher QoL and better disease management. **Objectives:** This study aimed to identify unmet communication and disease education needs in PwMS to empower informed decisions, enable self-management, and to maintain independence for as long as possible. In particular, we sought to gain an understanding of the impact of MS on employment and highlight the main challenges for people to remain in work. **Methods:** In October 2018 a roundtable meeting of patient representatives, PwMS, carers, and MS nurses agreed on key themes associated with maintaining independence. In 2019, an official project steering group formed, and 2 studies for PwMS were codeveloped: a qualitative online patient community activity and a quantitative online survey. The qualitative activity used Ipsos' Syndicated MS Online Patient Community (a consistent panel of PwMS), and the quantitative survey was recruited through the MS Trust monthly newsletter and Facebook group. Results were discussed and prioritized by the steering group. **Results:** Data were analyzed from 28 and 117 respondents with relapsing-remitting MS, from the Ipsos Syndicated MS Online Patient Community and quantitative survey, respectively. Data from the quantitative survey revealed the following: 66% of respondents "strongly/somewhat" agreed their MS had prevented them from reaching their full potential, and had an impact on QoL, with 73% of participants reporting that their MS had a "very/slightly negative impact" on work life. Overall, 69% were in full- or part-time employment, with only 26% remaining in the same job, with the same hours, since their diagnosis. Data from this study will further explore the impact on work for PwMS. **Conclusions:** Findings from our study highlight the impact of MS on working life among participants. It is important to understand these issues in the wider MS population to prevent problems at work and future loss of work.

Supported by: None

Disclosure: *Jessica O'Neill:* Roche Products Ltd (works for). *Mavis G. Ayer:* Biogen, Roche (sponsored delegate); Celgene, Merck, Novartis, Sanofi (consulting fee); Teva (sponsored education). *Samantha R. Colhoun, Alison Thomson:* Biogen, Novartis, Roche Products Ltd (consulting fee). *Nicola Daykin:* Roche Products Ltd (consulting fee). *Brenda Hamill:* Biogen, Novartis (sponsored delegate); Roche (consulting fee). *Maria Fei:* Roche Products Ltd (works for Ipsos MORI, which was funded by Roche to undertake this research). *Jordanne Florio:* Roche Products Ltd (works for). *Serena Pulcini:* Roche Products Ltd (works for).

Keywords: Disease education, Employment in MS, Management of activities of daily living in MS

(QOL14) Determining the Relationship of Demographic and Clinical Variables with Fatigue in Multiple Sclerosis, Using the 5-Item Modified Fatigue Impact Scale (MFIS-5)

Natasha Hameed,¹ Jennifer A. Ruiz,^{2,3} Heather M. DelMastro,^{2,4} Mary Bailey^{1,5}

¹Mandell Center for Multiple Sclerosis, Mount Sinai Rehabilitation Hospital, Trinity Health Of New England, Hartford, CT; ²Mandell Center for Multiple Sclerosis, Mount Sinai Rehabilitation Hospital, Trinity Health Of New England, Hartford, CT; ³Departments of Rehabilitative Medicine and Medical Sciences, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT; ⁴Department of Rehabilitative Medicine, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT; ⁵Assistant Clinical Professor Neurology, Yale School of Medicine, New Haven, CT

Background: Fatigue is reported as one of the most prevalent and disabling symptoms in patients with multiple sclerosis (MS). The Modified Fatigue Impact Scale (MFIS) is a self-reported tool that captures the degree of fatigue in relation to its impact on physical, cognitive, and psychosocial functioning. A recent article by Rooney et al has provided an improved understanding of the relationship of a variety of both clinical and demographic variables with the impact of fatigue on MS using the MFIS. To our knowledge, such a robust comparison has not been reported using the shortened version, the MFIS-5. **Objectives:** To determine the association between the impact of fatigue and demographics and clinical characteristics among persons with MS (PwMS). **Methods:** This study was a secondary analysis of a cross-sectional study of 253 PwMS. Demographic variables included age, gender, race (white/nonwhite), smoking status (smoker/nonsmoker), and employment status (employed/unemployed). Clinical characteristics included disease duration (DD), body mass index (BMI), level of disability (Patient-Determined Disease Steps [PDDS]), depression (Center for Epidemiologic Studies-Depression Scale [CES-D]), cognitive processing speed (Symbol Digit Modality Test [SDMT]), and use of disease-modifying therapy (DMT [on a DMT/not on DMT]). All measures, including fatigue impact (MFIS-5), were collected at a single visit. Spearman correlation coefficient was used to determine the strength of the associations and Mann-Whitney U for comparison. **Results:** The sample had a mean age of 48.6 ± 11.6 (range: 20-73) years, DD of 12.3 ± 8.7 (1-47) years, PDDS score of 2.7 ± 2.1 (range: 0-7), BMI of 28.9 ± 7.2 (range: 17.5-59.8), and MFIS-5 score of 9.6 ± 5.3 (range: 0-20). This sample was mostly female (75.9%), with relapsing MS (94.5%), unemployed (51.1%), white (84.5%), nonsmokers (84.9%), and on a DMT (83.8%). Fatigue impact was moderately correlated with disability ($r = 0.571$, $P = .000$) and depression ($r = .552$, $P = .000$) and weakly associated to age ($r = 0.173$, $P = .006$) and SDMT ($r = -0.297$, $P = .000$). No relationship was observed with DD or BMI. No difference was observed among dichotomized variables of gender, race, smoking status, or use of a DMT. However, the impact of fatigue was found to be different ($P = .000$) between those employed (median MFIS-5 score: 8) and unemployed (median MFIS-5 score: 12). **Conclusions:** Our study demonstrates the impact of fatigue, as measured by the MFIS-5, is associated with higher disability, depression, age, and decreased cognition and is greater among unemployed PwMS. Given its ease of administration, the MFIS-5 serves as a practical measure for assessing the impact of MS fatigue in clinical settings. The current study provides increased understanding of its relationship with various clinical and demographic variables.

Supported by: None

Disclosure: *Natasha Hameed, Jennifer A. Ruiz, Heather M. DelMastro: Nothing to disclose. Mary Bailey: Alexion, Biogen, Genentech, Genzyme, Novartis (speakers' bureau); Celgene (consulting fee).*

Keywords: *Comprehensive care and MS, Fatigue in MS*

(QOL15) Multiple Sclerosis Management and Expanded Disability Status Scale: A Great Start, but a Reason for Change Was Never So Apparent and Needed

Mark Gudesblatt,¹ Jared Srinivasan,¹ Olivia Kaczmarek,¹ Taylor Drost,¹ Barbara Bumstead,¹ Lori Fafard,¹ Kaitlyn Jaenicke,¹ Marijean Buhse,¹ Myassar Zarif,¹ Daniel Golan,² Jeffrey Wilken,³ Cynthia Sullivan,⁴ Timothy Fratto,⁵ Gavin Giovannoni⁵

¹South Shore Neurologic Associates, Patchogue, NY; ²Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel; ³Washington Neuropsychology Research Group, Washington, DC; ⁴Nursing, State University of New York at Stony Brook, Stony Brook, NY; ⁵Queen Mary University of London, Blizard Institute, Barts and the London School of Medicine and Dentistry, London, UK, London, United Kingdom

Background: Since the Expanded Disability Status Scale (EDSS) was pioneered by Dr John Kurtzke in 1967, it has been incorporated into clinical trial measurements in people with multiple sclerosis (PwMS). When combined with reported relapse rates and magnetic resonance imaging measurements of disease activity, EDSS has been the basis for approval of >15 disease-modifying therapies (DMTs). No evidence of disease

activity (NEDA) has been proposed as the goal for optimizing DMT. EDSS remains critical for both NEDA and treatment. Use of a nonlinear scale to measure disability can be problematic if there is great variability of PwMS within homologous EDSS-defined disability levels. Functional ability reflects the combined impact of cognitive function, manual dexterity, ambulation, and other factors. If the degree of variability of these "abilities" exceeds 20%, this scale would no longer be valid. **Objectives:** To explore the combined variability of functional performance with groups of similar disability across important aspects of ability. **Methods:** Retrospective review of prospective registry of PwMS who were evaluated by multidimensional computerized cognitive testing, digital gait analysis, and patient-reported outcomes (PROs) for hand function that had simultaneous measurements of PDDS or EDSS. **Results:** 258 PwMS, 73% female, age 46 ± 10 , multidomain computerized cognitive testing global summary score of 7 domains had adjacent EDSS score overlap (0-2.5, 3-4.5, 5-6.5, and >7) of 65% and extreme EDSS group overlap was 42%. Accumulative cognitive impairment (# cognitive domains impaired >1 SD) was 72% across adjacent EDSS groups, and extreme EDSS group overlap was 38%. 254 PwMS, 72% female, age 46 ± 10 , mean normalized velocity for preferred walking speed varied >20% within EDSS groups (A: 0-2.5 [24%], B: 3-4.5 [34%], C: 5-6.5 [53%]) and overlapped >20% (AB: 29%, BC: 25%). 783 PwMS, 74% female, age 49 ± 11 , completed PDDS and NARCOMS PRO for both hand function and tremor demonstrated variability >50% across all PDDS (0-1, 2-4, >4) and overlap >50% of adjacent PDDS groups and >32% across extreme PDDS groups. **Conclusions:** While the EDSS greatly advanced the treatment of MS, the degree of variability of disease impact within and across disability groups warrants immediate abandonment of this measure of care. This should be replaced by clinical trials with objective patient-centric multidimensional measures of disease impact to improve treatment selection and monitoring for progression.

Supported by: None

Disclosure: *Mark Gudesblatt: Acorda, Amgen, Medtronic, Saol Therapeutics (speakers' bureau); Biogen, EMD Serono, Novartis, Sanofi, Teva (contracted research). Jared Srinivasan, Olivia Kaczmarek, Taylor Drost, Lori Fafard, Kaitlyn Jaenicke, Daniel Golan, Timothy Fratto: Nothing to disclose. Barbara Bumstead, Marijean Buhse: Biogen, Genzyme (speakers' bureau). Myassar Zarif: Acorda, Biogen, Genzyme, Teva (speakers' bureau). Jeffrey Wilken: Biogen (contracted research); EMD Serono (speakers' bureau); Genzyme (contracted research, speakers' bureau). Cynthia Sullivan: Roche (contracted research). Gavin Giovannoni: AbbVie, Bayer Schering, Biogen, Canbec, Eisai, Elan, Five Prime, Genentech, Sanofi Genzyme, GlaxoSmithKline, GW Pharmaceuticals, Ironwood, Merck Serono, Novartis, Pfizer, Roche, Synthon BV, Teva (consulting fee); Multiple Sclerosis and Related Disorders (Elsevier) (co-chief editor); UCB Pharma (grants).*

Keywords: *Comprehensive care and MS, Equipment in MS, Natural history of MS*

(QOL16) A Coach-Supported Standardized Approach for Quality Improvement (QI) in the Multiple Sclerosis Continuous Quality Improvement (MS-CQI) Research Collaborative

Randall S. Messier

Randy Messier LLC, Fairfield, VT

Background: Multiple Sclerosis Continuous Quality Improvement (MS-CQI) is the first randomized, multicenter, prospective, longitudinal, randomized, systems-level, improvement science research collaborative for multiple sclerosis (MS). MS-CQI is a 3-year study of system-level variation in performance outcomes, and leverages performance benchmarking to inform improvement using an informatics-enabled learning health system approach. MS-CQI has 3 components: 1) collection and reporting of benchmarking data, 11 clinical outcome and 21 patient-reported outcome measures; 2) a quality improvement (QI) coach-supported QI intervention utilizing a standardized Tool Kit; and 3) a 3-year, step-wedge randomized study to test the effect of QI vs control on MS health outcomes. Four MS centers are participating: an urban academic center, a rural academic center, a rural community hospital, and a large urban private practice (total N = 5000). **Objectives:** Describe system readiness to engage in QI and characteristics of MS centers randomized to coach-supported QI intervention. **Methods:** MS centers randomized to QI participate in site visits and regular meetings with the QI coach. Teams use MS-CQI benchmarking data, local level data, and a standardized QI Tool Kit to develop their improvement theme and improvements. The QI coach conducts regular assessments of team QI knowledge, skills, attitudes, and readiness to engage in QI work at monthly intervals, including QI assessment scales (IHI Improvement Progress and QI Knowledge Application and Skills). **Results:** MS-CQI is in year 3. One center was randomized

to QI in year 2 (private for profit) of the study, and 2 centers were randomized (1 rural, 1 urban academic) in year 3. Site teams have identified QI foci including patient access, patient orientation, previsit planning, social work and behavioral health, and emergency room utilization. The teams are showing progress in their QI culture and development; IHI Improvement Progress Scale scores have changed from initial score of 1.5 (planning for the project has begun) to 3.5 (some improvements in measurements and outcomes and continuing to improve). **Conclusions:** MS center teams randomized to intervention in MS-CQI have successfully engaged in the coach-supported QI intervention. Site readiness and capability assessments reveals a greater understanding of their QI needs and a standard and consistent approach to improvement suggesting an initial positive response to the intervention.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Nursing management in MS, Quality improvement

(QOL18) Does Participation in a 6-Week Mindfulness Course Improve Mood and Overall Emotional Wellness for People Living with Multiple Sclerosis?

Tiffany Malone,¹ Lacey Sayre,¹ Brian Hutchinson²

¹Multiple Sclerosis Achievement Center, Dignity Health, Citrus Heights, CA; ²Multiple Sclerosis Achievement Center, Dignity Health, Sacramento, CA

Background: People with multiple sclerosis (MS) often experience emotional distress about the life they had prior to the diagnosis of MS and what that diagnosis means for their future. The Multiple Sclerosis Achievement Center (MSAC) conducts day wellness programs to address physical, cognitive, and social well-being. Program activities include exercise, brain training, education, socialization, and community outings. As part of the MSAC program, members have the opportunity to participate in a 6-week mindfulness course to address emotional wellness. **Objectives:** To determine, through the use of patient-reported outcomes (PROs), if members participating in the mindfulness course demonstrate improvements in mood and overall emotional wellness. **Methods:** Members of the MSAC will be asked to complete 4 paper/pencil outcome measures in January 2020 as part of their annual participation in the MSAC's programs, including the Multiple Sclerosis Impact Scale-29 (MSIS-29), Multiple Sclerosis Self-Efficacy Scale-10 item (MSSE), Godin Leisure-Time Exercise Questionnaire (GTLEQ), and Neuro-QoL (Anxiety, Depression, Emotion & Behavior, Positive Affect, Cognition, Ability to Participate, and Social Roles sections are used). Members will be offered the opportunity to participate in a professionally facilitated 6-week mindfulness course, starting in January 2020, to provide education, strategies, resources, and emotional support to achieve what they need and/or want for the present moment. Participants of the mindfulness course will be asked to complete PRO measures upon completion of the course. **Results:** One-year and 2-year comparison data of PROs, from MSAC members, have indicated correlations between Self-Efficacy, Anxiety, Ability to Participate, and Positive Affect (per MSSE & Neuro-QoL). Additionally, increased MSIS-29 scores directly correlate with anxiety while inversely correlating with self-efficacy and GTLEQ. Mean scores of depression, reported from the Neuro-QoL, remained the same ($x = 48$) over a 2-year reporting period. These comparisons did not specifically measure a member's participation in the mindfulness course. **Conclusions:** Data collection and analysis will be completed for those members participating in the 6-week mindfulness course. Analysis will include pre and post measures, as well as comparison with those not participating in the mindfulness course.

Supported by: None

Disclosure: Tiffany Malone, Lacey Sayre: Nothing to disclose. Brian Hutchinson: Biogen (speakers' bureau).

Keywords: Mindfulness

(QOL19) The Mediterranean Diet and Fatigue, Depression, and Emotional Well-being in Multiple Sclerosis: A Study in Patient-Reported Outcomes

Jamie Bolling,¹ Carley T. Rusch,² Alison Kraus,³ Nicole Tester,² Nicole Herndon,³ Aaron Carlson,¹ Carlos Vervloet Sollero,¹ Tirisham V. Gyang¹

¹Neurology, University of Florida, Gainesville, FL; ²Food Science and Human Nutrition, University of Florida, Gainesville, FL; ³Rehabilitation, UF Health, Gainesville, FL

Background: Multiple sclerosis (MS) is a chronic demyelinating inflammatory disease of the central nervous system and a leading cause of disability among young adults. It presents with a variety of neurologic symptoms, which can significantly affect the quality of life of affected individuals. Although no specific diet exists for MS, dietary factors show

potential for beneficial effects on inflammation, neuroprotection, and repair. Increased dietary quality has been associated with lower disability and symptom burden in MS. Various diets, including ketogenic, fasting, Mediterranean, and plant-based diets produced mixed results in studies. The Mediterranean diet (MD) is high in antioxidants, fiber, and mono- and polyunsaturated fatty acids as it emphasizes higher intake of fish, olive oil, fruits, and vegetables. Previous studies have shown the MD may be beneficial in risk reduction for cardiovascular diseases, type 2 diabetes, obesity, cognitive impairment, and brain atrophy. Limited studies have researched the relationship between MS outcomes and MD adherence. Preliminary evidence suggests that a MD may be beneficial in the MS population as it may be associated with reduced fatigue, impact of MS symptoms, and disability. Early findings warrant further investigation on the impact of an MD on MS outcomes to provide more evidence on the relationship of this diet on fatigue, cognitive function, and emotional well-being in MS. **Objectives:** To examine the relationship between adherence to the MD and fatigue, cognitive function, and emotional well-being, as captured by validated patient-reported outcomes in MS. **Methods:** Subjects will be recruited, consented, and enrolled from the University of Florida MS clinic. Adherence to the MD will be determined by using the validated 14-Item Mediterranean Diet Assessment Tool. Neuro-QOL questionnaires will score cognitive function and emotional well-being, while the validated Modified Fatigue Impact Scale will score fatigue. Data analysis will determine the relationship between adherence to the MD and fatigue, cognitive function, and emotional well-being. **Results:** Data collection is ongoing. **Conclusions:** Diet is an important factor to consider in comprehensive MS care and may have significant impact on quality of life. We hypothesize that MD will be associated with improved fatigue, cognitive function, and emotional well-being scores, although further research will better characterize its impact.

Supported by: None

Disclosure: Jamie Bolling, Carley T. Rusch, Alison Kraus, Nicole Tester, Nicole Herndon, Carlos Vervloet Sollero, Tirisham V. Gyang: Nothing to disclose. Aaron Carlson: Novartis Pharmaceutical (contracted research); Sanofi Genzyme (consulting fee).

Keywords: Complementary/alternative therapies in MS, Comprehensive care and MS, Nutrition and MS

REHABILITATION

(REH01) Assistive Technology for Progressive Deficits in Communication and Access in People with Advanced Multiple Sclerosis: Case Studies in Iterative Design

Alexander Burnham

Rehabilitation Services, The Boston Home, Boston, MA

Background: People with advanced multiple sclerosis (PwAMS; Expanded Disability Status Scale score > 6.5) can experience dynamic and progressive changes in motor speech intelligibility, vocal projection, and upper extremity control and coordination, which can interfere with access to typical expressive communication modalities (eg, telephone and computer access). In most cases, assistive technology (AT) can be useful in supplementing or replacing these functions to promote effective communication with partners and caregivers to maximize functional communication independence with minimal need for caregiver assistance. **Objectives:** Participants will have the opportunity to understand how PwAMS can maximize functional independence with appropriate and supported access to AT. Participants will recognize how increasingly mainstream and ubiquitous technology can be modified and interfaced with more customized AT to meet unique access challenges in the MS population. Participants will have a better understanding of how nursing and allied health/rehabilitation disciplines can collaboratively support PwAMS to complete electronic activities of daily living even in the presence of severe motor disability. **Methods:** This case series will describe the experiences of 3 PwAMS at the Boston Home, a specialized residence for individuals with advanced neurodegenerative disorders, who have benefited from using AT with ongoing customization and intervention to adapt the technology as appropriate to accommodate for multiple sclerosis-associated progressive motor, sensory, and cognitive deficits using currently available AT within the past 5 years. One individual will demonstrate concepts of iterative design as motor access to augmentative communication devices deteriorated; another will illustrate accommodations to speech-recognition technology despite progressive dysarthria; the third highlights cumulative adaptations to nurse call and speakerphone access options to compensate for worsening hypophonia and quadriplegia in order to maintain contact with remote family and caregivers. Photos of the applied AT will

illustrate how postural limitations and desire for wheeled mobility guided the decision-making process for selecting and modifying systems to meet the needs of the users. The roles of rehabilitation, nursing, and adaptive technology specialists in developing, modifying, and implementing appropriate iterations of communication AT in this particular care setting will be described as part of the interdisciplinary team caring for this cohort of PwAMS. **Results:** Recommendations for providing caregiver education and eliciting feedback and considerations for designing, implementing, and adapting communication AT for PwAMS in other residential environments will be discussed. Outcomes in terms of satisfaction with AT access, services, and devices from PwAMS and caregivers will also be described using formal and informal assessment options. **Conclusions:** NA.

Supported by: None

Disclosure: *Alexander Burnham: The Boston Home (salary).*

Keywords: *Comprehensive care and MS, Equipment in MS, Management of activities of daily living in MS*

(REH02) Predictors of Improvement in Respiratory Function Following Resistive Inspiratory Muscle Training in Advanced Multiple Sclerosis

Alexander Burnham,¹ Lisa Doyle,² Min H. Huang,² Donna Fry³

¹Rehabilitation Services, The Boston Home, Boston, MA; ²Physical Therapy, Franklin Pierce University, Manchester, NH; ³University of Michigan-Flint, Flint, MI

Background: Respiratory compromise in people with advanced multiple sclerosis (PwAMS = Expanded Disability Status Scale [EDSS] score ≥ 6.5) worsens as the disease progresses and is a major cause of morbidity and mortality. There is evidence that exercise can improve respiratory muscle function in PwAMS even in later stages of the disease. **Objectives:** It is not known if certain characteristics of PwAMS contribute to their ability to benefit from a respiratory exercise program. This study identified some possible predictive factors. **Methods:** Thirty-eight subjects were recruited at a SNF specializing in care for PwAMS. Inclusion criteria were age > 18 , multiple sclerosis (MS) diagnosis, and EDSS score ≥ 6.5 . The current study used a repeated measures within-subject design in which participants performed 3 sets of 15 repetitions of resistive inspiratory muscle exercises daily for 10 weeks using the Threshold Inspiratory Muscle Trainer (IMT). Demographics, number of comorbidities, body mass index (BMI), EDSS score, and years post-MS diagnosis were obtained at time of enrollment. Maximum inspiratory pressure (MIP) and maximum expiratory pressure were obtained as measurements of respiratory muscle strength at several intervals over the 28-week duration of the study, including 10-week baseline phase, 10-week intervention phase, and 8-week retention period postintervention. Progression of IMT resistance was adjusted weekly by the research team based on symptoms, rate of perceived exertion, and baseline MIP. Secondary outcomes assessed included fatigue (evaluated with MFIS-5) and cognitive processing speed (measured with oral version of Symbol Digit Modalities Test [SDMT]). **Results:** Correlation analysis of baseline characteristics with MIP change scores (MIP-CS) was performed to identify potential predictors of improvements in MIP for the regression analysis. Two separate linear regression models with MIP-CS and predicted values (MIP%-CS) as dependent variables were constructed. The regression model with BMI, fatigue (MFIS-5), and cognition (SDMT) as independent variables and MIP-CS as the dependent variable was significant ($F_{3,25} = 3.19, P = .041, R^2 = 0.53$). SDMT was a significant predictor in the model ($P = .035$) (higher SDMT scores significantly associated with better outcomes of IMT training); BMI as predictor approached significance ($P = .056$). The regression model with BMI, MFIS-5, and SDMT as independent variables and MIP%-CS as the dependent variable was significant ($F_{3,25} = 3.19, P = .027, R^2 = 0.55$). BMI was a significant independent predictor in the model ($P = .029$), with higher BMI associated with worse outcomes with IMT training; SDMT as predictor approached significance ($P = .053$). **Conclusions:** Participant scores in BMI, MFIS-5, SDMT, and age/gender-adjusted MIP at baseline significantly predicted MIP-CS ($F_{4,33} = 6.34, P = .001, R^2 = 0.44, R^2$ adjusted = 0.37). Factors such as age, gender, duration of disease, EDSS score, and number of comorbidities were not significant predictors of MIP-CS.

Supported by: None

Disclosure: *Alexander Burnham: The Boston Home (salary). Lisa Doyle, Min H. Huang, Donna Fry: Nothing to disclose.*

Keywords: *Comprehensive care and MS, Equipment in MS, Respiratory intervention in MS*

(REH03) Rehabilitation of Dysarthria in Multiple Sclerosis

Alice Estevo Dias

Scientific Research, Brazilian Association of Multiple Sclerosis, Sao Paulo, Brazil

Background: Multiple sclerosis (MS) can affect the speech motor system and result in dysarthria. The Lee Silverman Voice Treatment (LSVT-LOUD) is an intensive behavioral treatment program to improve overall voice and speech functions in individuals with neurologic diseases. **Objectives:** The main goal of this study is to inspect the efficacy of LSVT-X treatment in patients with MS. **Methods:** Thirty-four participants (relapsing-remitting MS = 16, secondary progressive MS = 14, primary progressive MS = 4), 27 female and 7 male, with age ranging from 31 to 81 years, received the LSVT-X, administered twice a week in 1-hour sessions over 8 weeks. The evaluation was performed by 3 expert speech therapists. The auditory-perceptive analysis of their voices was carried out, based on the GRBASI scale. The acoustic analysis was also conducted by Praat software (version 6.0.50), considering the following measures: fundamental frequency and voice intensity. Data were statistically analyzed to compare pre- and postrehabilitation. **Results:** Several signs and symptoms related to voice and speech were verified in different proportions and impacts on the intelligibility of verbal communication. Four altered domains have been identified: articulation (slow articulation, imprecise consonants), voice (pitch and intensity instability, impairment in vocal quality), respiration (decreased phonatory time, short cycles), and prosody (longer and frequent pauses, deficient loudness control). After treatment, there was improvement in parameters analyzed, dysarthria was attenuated, and positive results were achieved. **Conclusions:** In addition to conventional pharmacologic therapy, dysarthria treatment should be emphasized as part of a management plan focused on overall health and well-being, regardless of the type of MS, course of disease, and manifestation of speech and voice symptom.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *SLP intervention, Dysarthria*

(REH04) Daily Occupational Performance in Multiple Sclerosis

Alice Estevo Dias,¹ Gina Corsi²

¹Scientific Research and ²Rehabilitation, Brazilian Association of Multiple Sclerosis, Sao Paulo, Brazil

Background: Multiple sclerosis (MS) is a chronic, inflammatory, and degenerative disease that affects the central nervous system. Occupational performance is often compromised and negatively impacts daily activities and activities. **Objectives:** To understand the perception of people affected by MS on occupational performance and identify the main difficulties in routine activities. **Methods:** 55 people with MS participated, being 40 (73%) women and 15 (27%) men, aged between 27 and 60 years. The 5 major impairments in occupational performance were observed, according to the degree of importance, according to the Canadian Occupational Therapy Model (COTM), then the participants self-assessed their performances and satisfactions by means of a scale of 1 to 10 points. **Results:** The analyses revealed that participants considered their ability to perform routines and perform roles and tasks related to moderate to poor personal care, leisure, and productivity. **Conclusions:** Signs and symptoms of muscle weakness, fatigue, cognitive and visual changes, and sensitivity were determinant to impair occupational performance appropriate to the needs and interests of the participants. The evaluation of occupational therapy and the rehabilitation of disabilities organized and facilitated the daily lives of people with MS.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Occupational therapy*

(REH05) Assessing the Benefits of Using Telehealth in Conjunction with a Fitbit to Improve Walking in Veterans with Multiple Sclerosis

Kelley Volpenhein,¹ Amy Banks,¹ Lindsay J. Riegler,² Sarah Kiefer Luhring¹

¹Cincinnati VA Medical Center, Cincinnati, OH; ²Rehab Care Line, Cincinnati VA Medical Center, Cincinnati, OH

Background: Patients with multiple sclerosis (MS) often have limitations with mobility due to fatigue, weakness, and impairments with balance and coordination. Many of these patients decrease their activity and walking due to these limitations, which can lead to further deconditioning. **Objectives:** The objective of this study was to examine improvements in walking distance and perceived impact of walking ability in patients with MS using a Fitbit in conjunction with telerehabilitation. **Methods:** Currently 2 patients are enrolled in pilot with ongoing recruitment. Patients are asked for a self-perception of daily step count at evaluation. The 2-minute walk test and 10-meter walk test are administered at initial evalu-

ation for baseline measures. Patients are given a Fitbit and asked to do their normal activity until the first telehealth session. At the first telehealth session, the therapist views the patients walking log and establishes a weekly goal. Weekly sessions to monitor and progress home walking program occur over telehealth. After 12 sessions of telehealth, the patient attends a clinic post program evaluation to obtain outcome measures. The patient is given the Fitbit to continue with program independently. **Results:** To be determined. **Conclusions:** To be determined.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Equipment in MS, Management of activities of daily living in MS, Telehealth

(REH06) Feasibility of Telehealth Rehabilitation for Veterans with Progressive Neuromuscular Disease

Benjamin Dons

Physical Medicine and Rehabilitation, Veterans Health Administration, Saint Louis, MO

Background: Individuals who live in rural communities have difficulty accessing specialized medical services such as physical therapy. Individuals with progressive neurologic diseases who do have access to physical therapy are limited to what is available in their community. Exercise is feasible and can improve fitness and improve quality of life for individuals with progressive neurologic disease. Clinical video telehealth (CVT) provides these veterans with access to specialists for their condition and significantly reduces the energy and financial costs of traveling to specific appointments. Barriers exist whether an individual lives in a rural community or within a few miles of a health center. Utilizing CVT can eliminate these barriers and greatly improve adherence to a physical therapy rehabilitation program. Finally, the ability to view an individual in their home environment gives providers the ability to problem solve physical challenges and safety issues that may be present in a person's home. **Objectives:** 1) Extend specialty care from Neurology into Physical Medicine and Rehabilitation for Veterans with MS. 2) Determine the feasibility of the CVT devices in a variety of settings. 3) Decrease travel hours and costs. 4) Minimize caregiver time, burden, and other costs. **Methods:** There were 19 veterans evaluated during the study period, 18 of whom were diagnosed with multiple sclerosis (MS) and 1 with amyotrophic lateral sclerosis (ALS). Veterans were referred to PM&R physical therapy and evaluated by neurologic clinical specialist. They were evaluated before and after the intervention period using standardized functional outcome measures. Follow-up visits were scheduled at least once every week for 30-minute sessions and were re-evaluated every 30 days for up to 90 days. Veterans had the opportunity to extend their rehabilitation session for another 90 days if they were making improvements. Sessions took place with the clinical specialist in a private office with necessary rehabilitation equipment available for demonstration of exercises. **Results:** N = 18, 18 had preintervention assessments, and 10 patients agreed to participate. Four patients had a discharge visit where postintervention measures were collected. Total visits: 63. Travel miles saved: 6770 miles. Travel dollars saved: \$3724. **Conclusions:** Telehealth visits for patients with progressive neuromuscular disease such as MS and ALS can be effective and feasible in an outpatient setting. There were no adverse effects, and this program resulted in a significant reduction in miles traveled and cost savings for veterans and the Veteran Health Administration. Barriers to adopting new technology were an issue for some veterans, and greater improvements were seen with those who incorporated new technology easily.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS, MS and the caregiver/family

(REH07) Systematic Review on Exercise Training as a Neuroplasticity-Inducing Behavior in Multiple Sclerosis

Catherine D. Jones, Jessica F. Baird, Robert W. Motl, Brian M. Sandroff

Physical Therapy, University of Alabama at Birmingham, Birmingham, AL

Background: Exercise training is associated with improvements in physical fitness, walking mobility, balance, and possibly cognition in persons with multiple sclerosis (MS), perhaps based on neuroplasticity. However, it is difficult to characterize exercise training as a neuroplasticity-inducing behavior among persons with MS based on changes in functional outcomes alone, as neuroplasticity reflects true brain-behavior relationships. **Objectives:** The current systematic review provided a critical evaluation of exercise training as a neuroplasticity-inducing behavior in persons with MS based on a well-established conceptual model. This involved prioritizing exercise training studies in persons with MS that included both

functional and neuroimaging outcomes and further examined associations among these outcomes. **Methods:** We performed an open-dated search of online scholarly databases in July 2019 using a targeted and comprehensive search strategy. To be eligible for full-review, papers had to be published in English and include the following components: a) exercise training, b) neuroimaging outcomes, and c) functional outcomes (ie, measures of physical fitness, walking mobility, balance, and/or cognition) in persons with MS. Acceptable study designs included randomized controlled trials, single-group pre/post designs, and quasi-experimental designs. Four independent reviewers extracted relevant data from each eligible paper, including information on participant characteristics, exercise intervention characteristics, neuroimaging outcome characteristics, functional outcome characteristics, and pattern of study results. **Results:** The literature search returned only 9 papers (involving 7 original interventions) that met eligibility criteria wherein inferences regarding neuroplasticity could be drawn, based on the inclusion of both neuroimaging and functional end points. Within those 9 papers, there is mixed evidence for exercise training as a neuroplasticity-inducing behavior in persons with MS. **Conclusions:** There is insufficient data necessary to draw definitive conclusions on exercise as a neuroplasticity-inducing behavior in MS. Future research efforts might consider examining specific neural changes that would be expected to result from exercise prescriptions that are specifically designed to induce certain functional changes among persons with MS.

Supported by: None

Disclosure: Nothing to disclose

Keywords: CNS repair, Exercise, Imaging and MS

(REH08) Aerobic Reserve in People with Multiple Sclerosis: Measurement and Correlates

Corey D. Feasel, Brian M. Sandroff, Robert W. Motl

Physical Therapy, University of Alabama at Birmingham, Birmingham, AL

Background: The concept of aerobic reserve reflects the available potential energy to perform essential tasks to maintain independent living and is calculated by subtracting the energetic demand for activities of daily living from peak aerobic power. To date, the concept of aerobic reserve has not been applied in multiple sclerosis (MS), and there are limited data on its measurement and correlates in this population. **Objectives:** This study described the measurement and correlates of aerobic reserve in MS. **Methods:** The sample included 23 people with MS who were fully ambulatory (median [range] Expanded Disability Status Scale [EDSS] score = 3.5 [2.0]). Participants completed a single session that included obtaining informed consent, EDSS examination, demographic questionnaire, as well as administration of the Symbol Digit Modalities Test (SDMT), Timed 25-Foot Walk Test (T25FW), 6-minute walk distance (6MWD), and a cardiopulmonary exercise test (CPET) performed on a treadmill (modified Balke protocol). Aerobic reserve was calculated by subtracting the person's steady-state VO_2 extracted during the first stage of the CPET from peak VO_2 obtained from the CPET. **Results:** Twenty-one of 23 participants met criteria for providing a maximal effort during the CPET. The mean (SD) aerobic reserve was 9.5 (\pm 3.7) mL/kg/min. Aerobic reserve strongly correlated with peak VO_2 (mean [SD], 22.4 [\pm 5.4] mL/kg/min), $r = 0.77$, $P < .01$. Aerobic reserve was positively correlated with SDMT raw score (mean [SD], 49.3 [\pm 7.2]), $r = 0.45$, $P = .03$ and time to exhaustion in seconds on the CPET (mean [SD], 592.8 [\pm 205.5]), $r = 0.63$, $P < .01$. Aerobic reserve was negatively correlated with resting heart rate (mean [SD], 79.4 [\pm 11.5] bpm), $r = -0.50$, $P = .02$ and BMI (mean [SD], 29.6 [\pm 5.8] kg/m²), $r = -0.56$, $P < .01$. Aerobic reserve did not correlate with age, sex, EDSS score, or T25FW or 6MWD results. **Conclusions:** Aerobic reserve can be measured during CPET in people with MS, and might have implications for understanding symptomatic and functional outcomes in people with MS.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Aerobic exercise training, Complementary/alternative therapies in MS, Management of activities of daily living in MS

(REH09) The Impact of Vascular Comorbidities on Perceived Functional Impact in Persons with Multiple Sclerosis

Elizabeth S. Gromisch,^{1,2,3} Heather M. DelMastro,^{1,4} Lindsay O. Neto,^{1,4} Gabriele C. DeLuca,^{5,6,7} Jennifer A. Ruiz,^{1,3} Aaron P. Turner,^{8,9,10} Thomas P. Agresta,^{11,12,13} Albert C. Lo¹

¹Mandell Center for Multiple Sclerosis, Mount Sinai Rehabilitation Hospital, Trinity Health Of New England, Hartford, CT; ²Department of Neurology, University of Connecticut School of Medicine, Farmington, CT; ³Departments of Rehabilitative Medicine and

Medical Sciences, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT; ⁴Department of Rehabilitative Medicine, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT; ⁵Nuffield Department of Clinical Neurology, Oxford University, Oxford, United Kingdom; ⁶Department of Clinical Neurosciences, John Radcliffe Hospital, Oxford, United Kingdom; ⁷Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, United Kingdom; ⁸Department of Rehabilitation Medicine, University of Washington, Seattle, WA; ⁹Rehabilitation Care Service, VA Puget Sound Health Care System, Seattle, WA; ¹⁰Multiple Sclerosis Center of Excellence West, Veterans Affairs, Seattle, WA; ¹¹Department of Family Medicine, St. Francis Hospital and Medical Center, Trinity Health Of New England, Hartford, CT; ¹²Connecticut Institute for Primary Care Innovation, Hartford, CT; ¹³Center for Quantitative Medicine, University of Connecticut School of Medicine, Farmington, CT

Background: Persons with multiple sclerosis (PwMS) can have a number of comorbidities and secondary conditions, which can complicate care and negatively affect health-related quality of life. In particular, chronic vascular conditions, such as diabetes, hypertension, hyperlipidemia, and heart disease, have been associated with more rapid accumulation of irreversible disability in PwMS. However, little is known about the impact of co-occurring chronic vascular conditions on perceived functional impact in PwMS. **Objectives:** To examine differences between PwMS with and without vascular comorbidity with regards to their self-reported functional impact. **Methods:** Participants (n = 257) were randomly selected PwMS who participated in the MS Characterization of Upper Extremity Functioning (MS-CUE) study. The MS Impact Scale (MSIS-29) was used to measure perceived physical and psychological impact on daily life, while the Functional Status Index (FSI) was used to assess functional performance in 5 domains: gross mobility, hand activities, personal care, home chores, and social/role activities. Due to non-normally distributed data, Mann-Whitney U analyses were conducted to examine differences between PwMS with and without a vascular comorbidity, with effect size reported as r. **Results:** On average, PwMS were 48.72 ± 11.56 (0-73) years old and had MS for 12.40 ± 9.78 (1-47) years, with a median Patient-Determined Disease Steps (PDDS) score of 3 (0-7). A total of 112 of 257 (43%) PwMS had at least 1 co-occurring vascular condition, with hyperlipidemia (n = 72, 64.3%) and hypertension (n = 66, 58.9%) being the most common. PwMS with at least 1 vascular comorbidity reported higher levels of physical (r = -0.29, P < .001) and psychological impact (r = -0.16, P = .009), as well as more issues with gross mobility (r = -0.28, P < .001), hand activities (r = -0.17, P = .007), personal care (r = -0.19, P = .002), home chores (r = -0.24, P < .001), and social/role activities (r = -0.24, P < .001). **Conclusions:** PwMS with vascular comorbidity have worse perceived functional performance and physical and psychological well-being compared to PwMS without vascular comorbidity. These findings suggest that the presence of chronic vascular conditions in PwMS negatively affects perceived functioning, which has important implications for provision of care and quality of life for PwMS. Future work unraveling the mechanism by which vascular comorbidity influences perceived functional outcomes warrants further evaluation.

Supported by: None

Disclosure: Elizabeth S. Gromisch, Heather M. DelMastro, Lindsay O. Neto, Jennifer A. Ruiz, Aaron P. Turner, Thomas P. Agresta, Albert C. Lo: Nothing to disclose. Gabriele C. DeLuca: Merck-Serono (contracted research); Novartis (speakers' bureau); Sanofi Genzyme (travel).

Keywords: Management of activities of daily living in MS, Psychological issues and MS, Vascular comorbidity

(REH10) Is Treadmill Walking Analogous to Overground Walking in Persons with Multiple Sclerosis?

Evan T. Cohen,¹ Herbert Karpatkin,² Lourdes Rodriguez,² Rosangelis Rodriguez,² Rachel Rhodes,² Anna Rubeo²

¹Rehabilitation and Movement Sciences, Rutgers, The State University of New Jersey, Blackwood, NJ; ²Physical Therapy, Hunter College, New York, NY

Background: Gait and walking impairment is a common finding in persons with multiple sclerosis (PwMS) and is often studied using a treadmill. However, studies in other neurologic conditions have questioned whether treadmill walking accurately represents overground walking. If it does not, it may be questionable whether gait research performed on a treadmill can be generalized to overground walking. Similarly, it may suggest that the use of a treadmill for clinical gait evaluation and intervention for PwMS may not be supported. **Objectives:** The purpose of this study is to examine whether treadmill walking speed is similar to overground walking speed in PwMS. We hypothesized that treadmill walking speed would not be significantly different from overground walking speed in PwMS. If our hypothesis is correct, it will suggest that clinicians can be confident that use of a treadmill for the examination and treatment of gait and walking dysfunction is representative of overground walking. If

our hypothesis is not supported, however, it would suggest that clinicians who treat PwMS for gait and walking impairment should reconsider the use of a treadmill as a tool for evaluation and treatment. **Methods:** 19 people with MS (11 women and 8 men; median [IQR; range] Expanded Disability Status Scale score 4.5 [2.5; 2.0-6.5]) performed an overground 2-Minute Walk Test (2MWT) to determine their average walking speed, then were asked to walk at the same speed on a treadmill. Participants were given up to 10 minutes to familiarize themselves with the treadmill before trying to achieve their overground walking speed. **Results:** Pearson correlations were used to examine the relationship between predicted walking speed (PWS) (based on the mean walking speed during the 2MWT) and the actual walking speed (AWS) attained while on the treadmill. There was a positive correlation between AWS and PWS (r = 0.841 [r² = 0.707], n = 19, P = .000). 70.7% of the variance in PWS could be predicted by the AWS. This, however, leaves 29.3% of the variance unexplained. **Conclusions:** Although overground walking and treadmill walking may seem very similar, there are contextual differences in these activities that may limit generalizability between the two. Researchers should be cautious when generalizing outcomes in physical performance in PwMS measured during treadmill walking to expected performance in overground walking. Clinicians should also be cautious in expecting generalized training effects of treadmill walking on overground walking in PwMS. The walking techniques used on a treadmill may not be the same ones used for overground walking, and therefore treadmill walking may not generalize to overground walking.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS, Gait, Management of activities of daily living in MS

(REH11) Telerehabilitation Compared to Outpatient Rehabilitation for Patients with Multiple Sclerosis and Mobility Disorders

Heather Barksdale,¹ Brittany McHugh,¹ Wayne Hodges,² Jessica Peters,² Paul M. Hoffman³

¹Rehabilitation, UF Health Jacksonville, Jacksonville, FL; ²UF Health Jacksonville, Jacksonville, FL; ³Neuroscience Institute, UF Health Jacksonville, Jacksonville, FL

Background: Multiple sclerosis (MS) is a multifocal disease of the central nervous system, often producing variable and longstanding symptoms that may lead to loss of function and disability. Access to specialized rehabilitation specialists may be limited by loss of mobility and distance from rural areas to outpatient rehabilitation centers. This pilot study was undertaken to determine the feasibility of conducting a physical therapy-guided telerehabilitation (TR) program for individuals with mobility deficits resulting from MS. Data on mobility, quality of life (QOL), fatigue, and travel cost were examined. The TR group was then compared retrospectively to an outpatient-based therapy (OP) group to review effect on mobility scores. **Objectives:** 1) Determine if a TR program prescribed and monitored by an MS-certified physical therapist (PT), delivered on a home, web-based platform is a feasible delivery technique for individuals with MS. 2) Identify if a TR program can improve access to a component of comprehensive MS care while reducing travel cost. 3) Examine changes to mobility scores for a TR program compared with an OP program to compare the effects of the 2 delivery options for individuals with MS. **Methods:** Subjects with confirmed MS and mobility deficits were recruited from the MS Center of Excellence at UF Health Jacksonville for the TR group. Initial and final face-to-face examinations were performed by a board-certified neurologist and an MS specialist PT. Subjects underwent 8 weeks of physical therapy-guided TR using the Jintronix software platform and a kinetic tracking system. Subjects seen in OP by the same PTs performing the TR were selected by a chart review process from January 2018 through September 2019 with a diagnosis of MS (identified by a search of the practice's electronic database of the ICD 10 code G35). Subjects were selected for comparison based on duration of treatment and matching outcome measures completed. Data were then reviewed for effect on mobility and travel between the 2 groups. **Results:** Eight participants completed the TR program. All TR subjects demonstrated improvement in either fatigue, QOL, or mobility measures. No adverse events were noted during or following completion of the program. The 8 subjects saved a combined \$8487.23 in projected travel costs. Results of measures between groups showed equivalence in terms of meeting minimal detectable change for the outcome measures examined. **Conclusions:** The eight-week TR program was feasible and safe for subjects with MS and mobility impairments. Compared to the OP group, the TR group demonstrated the same number of subjects meeting the minimal detectable change for the functional measures. Savings were superior in the TR group for travel cost and time. Further studies are needed to guide the design and establish the efficacy of TR compared with OP rehabilitation programs.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Tele-rehabilitation

(REH12) Management of Low Back Pain for Individuals with Multiple Sclerosis: A Case Series

Heather Barksdale, Joseph Sugden

Rehabilitation, UF Health Jacksonville, Jacksonville, FL

Background: Individuals with multiple sclerosis (MS) are often referred to a physical therapist (PT) for evaluation of immobility, pain, and functional impairment. These individuals may be referred during the diagnosis process, a relapse, or while clinically stable. Self-reported pain symptoms for individuals with MS can also be multifactorial and originate from several areas including symptoms of central, peripheral, orthopedic, or a combination of these origins. Therefore, the underlying diagnosis of MS may complicate the evaluation, treatment plan, and progression for individuals with MS presenting with orthopedic complaints, including low back pain (LBP). This retrospective case series presents 4 cases of individuals with MS who were referred to physical therapy who also reported symptoms of LBP. **Objectives:** 1) Identify special considerations for evaluation techniques, treatment modifications, and progression modifications specific to individuals with MS and LBP. 2) Initiate guidelines that should be considered when establishing a plan of care for similar individuals with MS to establish the most effective approach for functional improvement or stability. **Methods:** All charts reviewed included subjects who attended the UF Health Jacksonville outpatient rehabilitation downtown location and were seen by the participating PTs from January 2018 through August 2019. Of the charts reviewed, 4 satisfied inclusion-exclusion criteria and are reviewed here. Charts were reviewed for PT plan of care, pain reports, and functional measures. **Results:** In each case, subjects reported improvement or resolution in pain measures and also demonstrated improvement in functional measures examined. Specific interventions and functional measures were tailored for each patient and were found to vary due to individual differences in clinical presentation and differences in response to a given intervention. **Conclusions:** The variability between subject presentation and complexity for individuals with LBP and underlying MS diagnosis were found to result in noted variance in treatment duration and approach between subjects. This points to the importance of thorough initial evaluation to include both neurologic and orthopedic standard of care. The evaluation ensures an appropriate plan of care for individualized treatment as well as identification of potential barriers to progression of treatment. As MS is a progressive condition, it is important to educate and train patients in ways to self-manage their musculoskeletal pain and functional deficits once an appropriate treatment plan has been established. Further research is needed to establish specific outcome measures and screening tools to identify individuals who will best benefit from outpatient physical therapy directed toward the impairment of LBP. The types of assessments and treatments reviewed in these cases may facilitate improved identification and standardization for these individuals.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Physical therapy

(REH13) Predicting Fall Risk in Persons with Multiple Sclerosis Utilizing the 12-Item Multiple Sclerosis Walking Scale

Heather M. DelMastro,^{1,2} Elizabeth S. Gromisch,^{1,3,4} Helen Dawes,^{5,6} Gabriele C. DeLuca,^{7,8,9} Robert Krug,^{1,4} Anton Pick,¹⁰ Jennifer A. Ruiz^{1,4}

¹Mandell Center for Multiple Sclerosis, Mount Sinai Rehabilitation Hospital, Trinity Health Of New England, Hartford, CT; ²Department of Rehabilitative Medicine, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT; ³Department of Neurology, University of Connecticut School of Medicine, Farmington, CT; ⁴Departments of Rehabilitative Medicine and Medical Sciences, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT; ⁵Department of Clinical Neurology, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, United Kingdom; ⁶Centre for Movement, Occupation and Rehabilitation Sciences, Faculty of Health and Life Sciences, Oxford Brookes University, Oxford, United Kingdom; ⁷Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, United Kingdom; ⁸Nuffield Department of Clinical Neurology, Oxford University, Oxford, United Kingdom; ⁹Department of Clinical Neurosciences, John Radcliffe Hospital, Oxford, United Kingdom; ¹⁰Oxford Centre for Enablement, Oxford University Hospitals NHS Foundation Trust, Oxford, United Kingdom

Background: Evidence suggests that 50%-80% of persons with multiple sclerosis (PwMS) have difficulty walking and impaired balance, with half of them falling at least once a year. Falls can lead to increased risk of

injury and fear of falling, which may further impair a person's function. Studies have previously shown that patient-reported outcomes predict PwMS's risk of falling, but small sample sizes and variable cutoff scores have limited generalization of the findings. **Objectives:** To determine the predictive value of a cutoff score for the 12-Item Multiple Sclerosis Walking Scale (MSWS-12) to identify PwMS with greater fall risk. **Methods:** A total of 135 PwMS were included as part of a preliminary analysis of an ongoing, larger cross-sectional study in which the MSWS-12 and frequency of falls (self-reported over past 6 months) were collected. PwMS were designated as "faller" if they had >1 fall in the past 6 months. Descriptive statistics were used to describe the clinical characteristics of the fallers (n = 82) and nonfallers (n = 53) (age, gender, disease duration, use of assistance, and Patient-Determined Disease Steps [PDDS] score). Clinical characteristics and MSWS-12 scores of the faller and nonfaller groups were compared. A receiver operating characteristic curve was used to estimate the classification accuracy of the MSWS-12. Optimal cutoff scores were calculated using the Youden index, and sensitivity and specificity were calculated. **Results:** There were no differences in age, gender, or disease duration between fallers and nonfallers. Fallers had higher median PDDS scores [3; 0-6 vs 1, 0-6; [P < .01]] and higher median MSWS-12 scores [67.5 vs 38.3; P < .001] than nonfallers. Fallers were more dependent on assistive devices compared to nonfallers (P < .01). The MSWS-12 cutoff score for fallers was ≥ 45.83 (Youden index: 0.46), with a sensitivity of 78.1%, specificity of 67.9%, and a classification accuracy of 76.7% to detect fallers. **Conclusions:** MSWS-12 was found to be predictive of fall risk in PwMS with a cutoff score much lower than previously reported. These findings indicate a lower threshold of the MSWS-12 score may help clinicians identify PwMS at greatest fall risk so that appropriate fall risk prevention interventions may be implemented.

Supported by: None

Disclosure: Heather M. DelMastro, Elizabeth S. Gromisch, Helen Dawes, Anton Pick, Jennifer A. Ruiz: Nothing to disclose. Gabriele C. DeLuca: Merck-Serono (contracted research); Novartis (speakers' bureau); Sanofi Genzyme (travel). Robert Krug: Novartis (contracted research).

Keywords: Comprehensive care and MS, Falls

(REH14) The Impact of Lower Limb Strength on Walking in Persons with Multiple Sclerosis: A Preliminary Analysis

Heather M. DelMastro,^{1,2} Helen Dawes,^{3,4} Gabriele C. DeLuca,^{5,6,7} Robert Krug,^{1,8} Anton Pick,⁹ Jennifer A. Ruiz^{1,8}

¹Mandell Center for Multiple Sclerosis, Mount Sinai Rehabilitation Hospital, Trinity Health Of New England, Hartford, CT; ²Department of Rehabilitative Medicine, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT; ³Centre for Movement, Occupation and Rehabilitation Sciences, Faculty of Health and Life Sciences, Oxford Brookes University, Oxford, United Kingdom; ⁴Department of Clinical Neurology, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, United Kingdom; ⁵Nuffield Department of Clinical Neurology, Oxford University, Oxford, United Kingdom; ⁶Department of Clinical Neurosciences, John Radcliffe Hospital, Oxford, United Kingdom; ⁷Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, United Kingdom; ⁸Departments of Rehabilitative Medicine and Medical Sciences, Frank H. Netter MD School of Medicine at Quinnipiac University, North Haven, CT; ⁹Oxford Centre for Enablement, Oxford University Hospitals NHS Foundation Trust, Oxford, United Kingdom

Background: Persons with multiple sclerosis (PwMS) report weakness and walking difficulty as some of their most disabling symptoms. Lower limb (LL) weakness is prevalent in PwMS and is associated with more significant disability, impaired balance, and increased difficulty walking. However, limited research exists describing the relationship between strength of specific LL muscle groups and walking in the same cohort. **Objectives:** To determine the impact of dominant (D) and nondominant (ND) LL strength on patient-reported outcomes (PROs) and objective walking outcome measures in PwMS. **Methods:** A cross-sectional sample of PwMS (n = 137) derived from a larger, ongoing study was used. The following walking measures were collected at a single visit: 12-item MS Walking Scale (MSWS-12), Timed 25-Foot Walk (T25FW), and D and ND stride length (StrL), step length (SL), and double support time (DStime). Isometric peak torque of hip extension and flexion (HExt; Flex), knee extension and flexion (KExt; Flex), ankle plantar and dorsiflexion (APF; DF), and hip abduction (HAbd) were also collected. Descriptive statistics were performed (age, gender, disease duration, and disability level: Patient-Determined Disease Steps [PDDS] score) and a correlational analysis was used to determine the strength of the association of walking to strength in muscle groups. **Results:** The MS cohort had a mean age of 51.4 (range: 21-75) years, disease duration of 14.5 (range: 0.3-40.0) years, and median PDDS score of 2.5 (range: 0-7), with 74.1% being female. All muscle groups were correlated with SL and StrL, and inversely correlated with T25FW, MSWS-12, and DStime. Strong associations were observed between D HFlex and StrL (D: r = 0.621, P < .001; and ND: r = 0.636, P < .001), D HFlex and ND SL (r = 0.608, P < .001),

ND KFlex and StrL (D: $r = 0.610$, $P < .001$; and ND: $r = 0.622$, $P < .001$), ND HAbd and ND SL ($r = 0.640$, $P < .001$), and ND HAbd and StrL ($r = 0.605$, $P < .001$). Weak to moderate correlations ($r = \pm 0.190$ to 0.599 , $P < .05$) were found for all remaining strength and walking measures assessed. **Conclusions:** All LL muscle groups (HExt, HFlex, KExt, KFlex, APF, ADF, and HAbd) were associated with the PRO (MSWS-12) and objective walking variables (T25FW, gait parameters: StrL, SL, and DStime) collected. These findings suggest that strength training interventions of these muscles may improve walking in PwMS. Importantly, this study improves understanding of the relationship between different major LL muscle groups with both walking performance and perceived difficulty walking in PwMS.

Supported by: None

Disclosure: *Heather M. DelMastro, Helen Dawes, Anton Pick, Jennifer A. Ruiz: Nothing to disclose. Gabriele C. DeLuca: Merck-Serono (contracted research); Novartis (speakers' bureau); Novafi Genzyme (travel). Robert Krug: Novartis (contracted research).*

Keywords: *Comprehensive care and MS, Walking*

(REH15) The Effects of Intermittent Versus Continuous Walking on Distance to Fatigue in Persons with Multiple Sclerosis

Herb Karpatkin, Rachel Rhodes, Lourdes Rodriguez, Rosangelis Rodriguez, Anna Rubeo
Physical Therapy, Hunter College, New York, NY

Background: Diminished walking endurance is common in persons with multiple sclerosis (PwMS). Previous studies have shown that PwMS walk farther in 6 minutes when using intermittent walking (IW) (ie, with interspersed rest breaks) than with continuous walking (CW), but it is unknown whether PwMS can walk greater distances or longer duration when using IW for periods of longer than 6 minutes. **Objectives:** The purpose of this study is to compare distance and time walked on a treadmill at a fixed velocity utilizing IW or CW. We hypothesized that PwMS would be able to walk greater distances and for longer duration when walking intermittently than when walking continuously. If our hypothesis was correct, it would indicate that PwMS can increase the total distance that they walk before being limited by fatigue using intermittent as opposed to continuous walking training. **Methods:** A randomized crossover design was used. Participants were randomized into 2 order groups: IW then CW, or CW then IW. The IW condition included alternating 30 seconds of walking and 30 seconds of seated resting. The CW condition consisted of continuous walking. Participants wore an overhead harness for safety. Baseline walking speed was determined with a 2-minute walk test (2MWT). Participants walked at the fastest pace up to the 2MWT speed until they either lost their balance or asked to stop. Walking time and walking distance were recorded. After 1 week, participants returned and performed the crossover condition. **Results:** 19 subjects (Expanded Disability Status Scale score 4.7 ± 1.4 , 10 female) completed the study. Participants had significantly longer walking distance in the intermittent condition than in the continuous condition (1575.4 ft SD ± 498.4 vs 1035.9 ft SD ± 356.2 , $P = .028$). IW enabled participants to walk at best-pace for greater distances than CW. **Conclusions:** These findings further support the use of IW training to improve walking endurance in PwMS. Adding rest breaks during endurance training enabled participants in this study to walk farther and longer, increasing the "dosage" of the walking activity. Comparative effectiveness studies should be conducted to determine whether IW training is superior to the traditional model of CW training to improve walking endurance. In PwMS, greater walking distance can be achieved with IW than with CW, suggesting that greater walking endurance gains can be made in these patients using this approach.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Complementary/alternative therapies in MS, Gait, Management of activities of daily living in MS*

(REH16) The Effects of Cooling Vests on Gait Fatigability in Persons with Multiple Sclerosis

Herbert Karpatkin, Sean Riva, Kaitlin Russo, Sandon Stevens, Winnie Yu
Physical Therapy, Hunter College, New York, NY

Background: Gait dysfunction is a ubiquitous and multifactorial finding in persons with multiple sclerosis (PwMS). A major reason for gait dysfunction in PwMS is objective fatigability (OF), characterized by progressive worsening of gait parameters over the course of a walk. Although OF is also multifactorial, it is presumed to be due in large part to multiple sclerosis (MS) thermosensitivity, where increased heat leads to

decreased conduction through demyelinated nerves. Prolonged exercise such as long walks can lead to increased core temperature in PwMS, and therefore lead to progressive worsening of gait over the course of the walk. An intervention to prevent core temperature rise could limit OF during gait in PwMS. **Objectives:** The purpose of this study was to investigate whether the use of a commercially available cooling vest would result in decreased OF of gait in PwMS. We hypothesized that wearing the vest for 30 minutes prior to a 6-minute walk (6MW) would result in less evidence of gait fatigability in PwMS when compared to performing the 6MW without prior cooling. If our hypothesis is correct, it will suggest that PwMS who experience gait fatigability can mitigate this by the use of cooling garments. **Methods:** A randomized crossover design is being used. Ambulatory patients with a diagnosis of MS are randomized into a cooled and uncooled condition. Cooling is accomplished by the wearing of a commercially available cooling vest for 30 minutes while seated. The uncooled condition is sitting for 30 minutes without wearing the vest. Immediately after the 30 minutes, subjects perform a 6MW test. Objective fatigability is measured by comparing the speed of the walk in the first minute to the speed of the walk in the sixth minute. Subjective fatigue is measured using the Visual Analog Scale of Fatigue (VASF). Data collection began fall of 2019 and will conclude winter of 2020. **Results:** To date, 5 subjects (Expanded Disability Status Scale score 4.4) have completed the study. Due to the small sample size, only descriptive statistics are reported. Mean 6MW test distance was higher in the cooled condition ($1137.3'$) than in the uncooled condition ($1087.9'$). Mean differences between the distance walked in first minute and sixth minute was less in the cooled condition ($-1.6'$) than in the uncooled ($-12.4'$). Subjects experienced less subjective fatigue as measured by the VASF in the cooled condition (7.4 mm) than in the uncooled (13.8 mm). **Conclusions:** These findings, although preliminary, support our hypothesis that cooling may diminish OF of gait in PwMS and thereby improve gait endurance. Once we have achieved an adequate sample size, a more in-depth analysis will be performed. If our hypothesis is then reaffirmed, it will suggest that the use of a commercially available cooling vest may decrease the impact of fatigue on gait in PwMS.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Equipment in MS, Gait fatigue, Management of activities of daily living in MS*

(REH17) A Combination of Core Exercise and Balance-Based Torso Weighting for Women with Multiple Sclerosis

Jeannie B. Stephenson,¹ Douglas E. Haladay,¹ Stephanie L. Carey,² Jason W. Beckstead,³ Derrick Robertson⁴

¹School of Physical Therapy & Rehabilitation Sciences, University of South Florida, Tampa, FL;

²Department of Mechanical Engineering, University of South Florida, Tampa, FL;

³College of Public Health, University of South Florida, Tampa, FL; ⁴College of Medicine, Neurology, University of South Florida, Tampa, FL

Background: Multiple sclerosis (MS) is a neurodegenerative disease that often results in fatigue and balance and walking impairment. Core exercise has been shown to reduce fatigue and improve balance and walking in people with MS. However, no studies have investigated the effects of a combination of core exercise and balance-based torso weighting (BBTW). **Objectives:** The purpose of this study was to investigate whether the combination of BBTW plus core exercise leads to greater improvement in self-reports of fatigue, balance confidence, and walking ability compared to core exercise alone in women with MS. **Methods:** Eighteen women with MS (Expanded Disability Status Scale score 3.0-5.0) were randomly assigned to 1 of 2 groups: core exercise (CE) or BBTW plus core exercise (BBTW + CE). Subjects completed 3 questionnaires at baseline and after a 6-week intervention period: Modified Fatigue Impact Scale (MFIS), Activities-Specific Balance Confidence Scale (ABC Scale), and the MS Walking Scale (MSWS-12). All subjects participated in a Pilates-based CE program once a week with a physical therapist along with a daily home exercise program. In addition to the CE, 1 group also participated in the BBTW protocol. This involved fitting subjects with a vest worn on the torso and application of small weights to the vest at baseline followed by biweekly sessions to adjust weights and gradually increase wearing time up to 6 hours daily. **Results:** Following the 6-week intervention period, both groups demonstrated positive change indicating improvements in self-reported fatigue, balance confidence, and walking ability. The percent change for each measure was as follows: MFIS: CE group = 14.1% decrease, BBTW + CE group = 19.9% decrease; ABC Scale: CE group = 9.7% increase, BBTW + CE group 15.6% increase; and MSWS-12: CE group: 11.9% decrease, BBTW + CE group = 19.3% decrease. Despite these improvements, none of the

change in scores exceeded the MDC_{95} estimates for each measure (MFIS = 49%, ABC = 20%, MSWS-12 = 53%). **Conclusions:** Core exercise with or without BBTW led to decreased self-perceived fatigue and improved balance confidence and walking ability, however, the percent change for both groups did not exceed MDC_{95} estimates. The percent change in perceived fatigue, balance, and walking was greater in the BBTW + CE group. The balance wear vest may provide individuals with added truncal proprioceptive input and recruitment of core stabilizers, however, the mechanism of improvement needs to be further investigated.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS, Core exercise

(REH21) Impact of Restless Legs Syndrome Severity on Cognitive Function in Adults with Multiple Sclerosis and Restless Legs Syndrome

Katie L.J. Cederberg,¹ Morgan L. Schuetz,¹ Brianna Mathison,¹ Tiffany J. Braley,² Arthur S. Walters,³ Robert W. Motl¹

¹Physical Therapy, University of Alabama at Birmingham, Birmingham, AL; ²Neurology, University of Michigan, Ann Arbor, MI; ³Neurology, Vanderbilt University, Nashville, TN

Background: Restless legs syndrome (RLS) is a sleep disorder present in as many as 26% of persons with multiple sclerosis (PwMS) and may exacerbate many of the symptoms and consequences of multiple sclerosis (MS), including cognitive function. Additionally, RLS symptoms often impair sleep quality, which could further exacerbate the neuropsychological symptoms associated with MS. **Objectives:** The present study examined the relationship between RLS severity and cognitive impairment in adults with MS and RLS. **Methods:** Participants with MS were screened for RLS using the Cambridge-Hopkins Restless Leg Syndrome Short Form Diagnostic Questionnaire. Participants attended 1 session wherein a rater performed an examination for scoring the Expanded Disability Status Scale (EDSS) and participants completed the International Restless Legs Syndrome Study Group Scale (IRLS), the Pittsburgh Sleep Quality Index (PSQI), and the Epworth Sleepiness Scale (ESS) followed by the Brief International Cognitive Assessment for Multiple Sclerosis battery consisting of the Symbol Digit Modalities Test (SDMT), California Verbal Learning Test-II (CVLT-II), and Brief Visuospatial Memory Test-Revised (BVMTR). **Results:** All participants (N = 22) had RLS (mean IRLS = 20.4; SD = 6.1). Non-parametric bivariate correlations indicated strong, negative associations between scores from the IRLS and CVLT-II ($p = -0.627$; $P < .01$), BVMTR ($p = -0.608$; $P < .01$), and years of education ($p = -0.632$; $P < .01$). There were no significant associations among scores from the IRLS and the SDMT, PSQI, or ESS. We performed 2 multivariate linear regressions with forward stepwise selection wherein we regressed scores from 1) CVLT-II on IRLS scores and 2) BVMTR on IRLS scores in step 1, and included variables that were significantly correlated with cognitive scores and IRLS scores in bivariate correlation analyses in step 2 (ie, years of education). IRLS scores significantly predicted CVLT-II ($R^2 = 0.398$) and BVMTR performance ($R^2 = 0.371$); however, the relationship with BVMTR performance was attenuated by including years of education ($\Delta R^2 = 0.144$). **Conclusions:** Our findings suggest that worse RLS severity could contribute to worse immediate verbal recall and memory and worse immediate visual recall and visuospatial memory in PwMS. Additional research is necessary to explore mechanisms that may underlie this association. If a causal pathway exists, diagnosis and treatment of RLS symptoms may offer new opportunities to reduce cognitive impairment in adults with MS.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Psychological issues and MS, Sleep and MS

(REH22) Utilization of Therapy Services Among Patients with Multiple Sclerosis

Laura K. Miller

Exercise Science, Bellarmine University, Louisville, KY

Background: Multiple sclerosis (MS) is a chronic, progressive neurologic disease estimated to be affecting nearly 1 million individuals in the United States. Common symptoms of MS include pain, impaired mobility or activities of daily living, and fatigue, among many others. The use of rehabilitation in all settings has been shown to improve symptoms and function while helping to minimize disability. **Objectives:** The purpose of this study was to examine the current usage of various rehabilitation services in the Kentucky area and understand barriers to rehabilitation among patients with MS. A secondary objective was to identify areas in which patients require additional education related to managing their disease. **Methods:** Eighty-nine participants completed a voluntary online survey that included 19 questions about their disease and experiences with being referred to various types of rehabilitation. Additional questions

were asked about their knowledge and beliefs and any barriers related to rehabilitation. Surveys were completed in Survey Monkey and results analyzed. Although the sample size was small, important information about therapy usage was gathered during this survey. **Results:** Over half of participants are not participating in regular exercise, despite current literature that shows the benefits of exercise. Approximately one third of individuals surveyed are unaware of the benefits of various types of rehabilitation. Half of the participants have plans to make changes to either their diet or exercise programs over the next year. Around 50% of participants expressed interest in learning more about nutrition, supplements, exercise, stress management, and stretching. These results highlight the need for continued patient education regarding management of MS symptoms. **Conclusions:** Despite significant research that indicates the benefits of therapy for patients with MS, there continue to be many patients who have not been referred to skilled therapy services. Furthermore, most patients are interested in obtaining additional information related to managing their disease, which could be addressed by a comprehensive rehabilitation team.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS, Comprehensive care and MS, Management of activities of daily living in MS

(REH23) Proximal Movement Compensations Are Related to Muscle Function and Walking Capacity in People with Multiple Sclerosis

Mark M. Manago,^{1,2} Enrique Alvarez,² Paul Kline,^{1,3} Cory Christiansen^{1,3}

¹Department of Physical Medicine and Rehabilitation, University of Colorado School of Medicine, Aurora, CO; ²Department of Neurology, University of Colorado School of Medicine, Aurora, CO; ³Geriatric, Research, Education, and Clinical Center, VA Eastern Colorado Healthcare System, Aurora, CO

Background: Distal lower extremity movement compensations are associated with muscle weakness and mobility limitations in people with multiple sclerosis (MS), however less is known about pelvis and trunk compensations during walking. **Objectives:** To 1) compare differences in pelvis and trunk kinematics during walking between participants with MS and a control group, and 2) determine associations of trunk and pelvis kinematics with muscle function, spatiotemporal parameters, and walking capacity in the participants with MS. **Methods:** In this cross-sectional study, 20 people with MS (Expanded Disability Status Scale 1.5-5.5) and 10 people with comparable age and sex (CTL) underwent 3-dimensional gait analysis. The primary kinematic variables of interest were frontal and sagittal plane pelvis and trunk angular displacement during the stance period of walking. All participants also underwent muscle function assessments (hip and trunk strength and endurance), and walking capacity measures (Timed 25-Foot Walk [T25FW], 2-Minute Walk Test [2MWT]). **Results:** Compared to the CTL group, the MS group had significantly greater sagittal plane trunk and pelvis angular displacement for both the stronger ($P = .031$) and weaker ($P = .042$) sides, less frontal plane trunk and pelvis angular displacement for both the stronger ($P = .008$) and weaker ($P = .024$) sides, and more sagittal plane trunk angular displacement for the stronger side ($P = .047$) during stance phase. There were low-to-moderate correlations in the MS group for sagittal plane pelvis angular displacement with trunk flexion endurance ($r = -0.369$, $P = .019$), and frontal plane pelvis angular displacement with lateral trunk flexion strength ($r = 0.353$, $P = .030$), step length ($r = 0.529$, $P < .001$), stance time ($r = -0.433$, $P = .005$), T25FW ($r = 0.496$, $P = .001$), and 2MWT ($r = 0.582$, $P < .001$). **Conclusions:** In people with MS, movement compensation at the pelvis during walking, particularly decreased frontal plane motion, was associated with worse walking capacity, muscle function, and spatiotemporal parameters. Future studies may consider targeting proximal muscle function to improve walking outcomes in people with MS. Rehabilitation clinicians may consider evaluation of proximal muscle function and gait compensations when planning rehabilitation interventions to improve walking capacity in people with MS.

Supported by: None

Disclosure: Mark M. Manago, Paul Kline, Cory Christiansen: Nothing to disclose. Enrique Alvarez: Actelion, Biogen, Celgene, EMD Serono, Genentech, Genzyme, Novartis, Teva, TG Therapeutics (consulting fee); Biogen, Genentech, Novartis, Rocky Mountain MS Center (contracted research).

Keywords: Rehabilitation in MS

(REH24) Effects of a Weight-Based Training Program on Bone Density, Cognition, and Quality of Life of Multiple Sclerosis Patients

Mary Ann Picone

Holy Name Medical Center, Teaneck, NJ

Background: Multiple sclerosis (MS) is a chronic neurodegenerative disease caused by the lesion forming demyelination of the central nervous system. Some of the issues associated with MS include cognitive impairment, increased mental and physical fatigue, and decreased bone mineral density. Using different variations of exercise is a common practice in the care for patients diagnosed with MS. It has been supported that there is a positive impact associated with exercise and the long-term progression of functional limitations and quality of life measures in patients with MS. **Objectives:** Patients with MS have lower bone mineral density and a higher prevalence of osteoporosis. Physical activity has had a positive effect in bone health of patients with MS. The primary purpose of this study is to investigate if there is a correlation between weekly body weight exercise and bone density in patients with MS. Cognitive functioning and psychological well-being have also been shown to improve through the intervention of regular exercise. Additionally, cognitive and quality-of-life measures will also be investigated as part of this study. **Methods:** This study will enroll a total of 25 patients. Patients eligible for the study are between the ages of 40-55, diagnosed with MS, and having an Expanded Disability Status Scale score below 5.5. Each patient will receive a baseline dual energy X-ray absorptiometry (DEXA) scan, a verbal Symbol Digit Modalities Test (SDMT), and a Multiple Sclerosis Impact Scale (MSIS-29). Following the baseline visit, patients begin a 6-week body weight exercise program. The program consists of one 30-minute group session, under trained physical therapist supervision, and 1 video-guided at-home session for a total of 2 sessions per week. Upon completion of the training program, each patient completes a midpoint SDMT and MSIS-29. An end-point DEXA scan, SDMT, and MSIS-29 are conducted 8 weeks after the completion of the training program. Statistical analysis will be conducted for potential changes in bone density values, SDMT scores, and MSIS-29 physical and psychological scores. **Results:** To date, 4 patients completed, 5 patients dropped, and 4 patient enrolled/awaiting enrollment. Data are being collected and further analysis is required. **Conclusions:** Thus far patients have reported positive experiences, but until analysis no conclusions can be made or supported with regard to bone density and cognitive effects.

Supported by: None

Disclosure: *Mary Ann Picone: Biogen Inc (speakers' bureau).*

Keywords: *Body weight exercise for MS, Complementary/alternative therapies in MS, Management of activities of daily living in MS*

(REH25) Orchestrating a New Path for Multiple Sclerosis Rehabilitation: Empowering Patients Through Both Physical and Music Therapies

Megan Weigel,¹ Renee Fleming,² Brian Hutchinson,³ Wendy L. Magee,⁴ Wendy Su⁵

¹First Coast Integrative Medicine, Jacksonville Beach, FL; ²John F. Kennedy Center for the Performing Arts, Washington, DC; ³Multiple Sclerosis Achievement Center, Dignity Health, Sacramento, CA; ⁴Temple University, Philadelphia, PA; ⁵Novartis Pharmaceuticals Corporation, East Hanover, NJ

Background: Although disease-modifying therapies (DMTs) are available for multiple sclerosis (MS) to delay disability progression and reduce relapses, as MS progresses additional support and management of symptoms become increasingly important. **Objectives:** To assess the role of nonpharmacologic therapies, focusing on physiotherapy (PT) and music therapy (MT) that can lead to improvements in most of the physical and psychosocial domains that are negatively affected in patients with MS. **Methods:** MEDLINE was searched without date restriction to identify studies on the efficacy of PT and MT in MS. A panel of PT and MT experts was convened to identify important themes and research studies. **Results:** PT can lead to improvements in mobility and balance. A review of 16 randomized controlled trials showed that treadmill training 3 times per week for 8 weeks improved walking endurance by 26.5 m from baseline in patients with MS. PT can also provide clinically meaningful improvements in fatigue, health-related quality of life (HRQoL), mood, and cognition. In a group of 20 patients with MS who performed high-intensity resistance training twice a week for 12 weeks, patients achieved statistically significant reductions in anxiety ($P = .002$), depression ($P = .019$), and fatigue ($P = .001$). Likewise, MT can improve physical symptoms and HRQoL in MS. In a trial comparing rhythmic-cued motor imagery, metronome-cued motor imagery, and no intervention, patients in the 2 intervention groups could, respectively, walk a mean of 62.1 m and 60.9 m further after 4 weeks vs baseline; the mean change in the no-intervention group was -17.1 m. Significant improvements in HRQoL measures were also seen in both intervention groups vs the no-intervention groups for physical function, general health perception, vitality, social function, and mental health ($P < .05$). **Conclusions:** While DMTs aim to reduce disability progression and inflammatory activities in MS, additional nonpharmacologic therapies are an important adjunct for managing daily life with MS, particularly in improving or maintaining mobility,

cognition, and other functional systems. Current studies regarding the use of PT and MT in MS indicate these are highly beneficial tools in delaying the loss of both fine and gross motor skills, improving overall well-being and psychosocial health factors, and, ultimately, preserving HRQoL. Further research on combined PT and MT interventions may further improve outcomes in MS.

Supported by: None

Disclosure: *Megan Weigel: Acorda, EMD Serono, Mallinckrodt Pharmaceuticals (speakers' bureau); Biogen Inc, Celgene Corporation, Inc (consulting fee); Novartis Pharmaceuticals Corporation, Sanofi Genzyme (consulting fee, speakers' bureau). Renee Fleming, Brian Hutchinson, Wendy L. Magee: Novartis Pharmaceuticals Corporation (consulting fee). Wendy Su: Novartis Pharmaceuticals Corporation (salary).*

Keywords: *Complementary/alternative therapies in MS, Comprehensive care and MS, Management of activities of daily living in MS*

(REH26) Improving Quality of Life Using an End-Effector Robotic Rehabilitation Approach in Progressive Multiple Sclerosis

Savanur Deepak Rajendra

Neurorehabilitation and Robotics, IU Health, Indianapolis, IN

Background: Progressive multiple sclerosis (MS) is characterized clinically by gradual disease progression and accumulation of neurologic disability, independent of relapses. Rehabilitation has been recommended as a means to reduce disability and restore function. High-quality evidence supporting progressive MS rehabilitation is limited. An end-effector robot-assisted gait trainer (RAGT) addresses many of the limitations of therapist-assisted gait training while providing an environment for regaining mobility and independence. **Objectives:** The objective of this study was to establish the safety and feasibility of RAGT and determine its impact on movement capacity, fatigue, and quality of life in patients with progressive MS. **Methods:** Single-blinded, randomized clinical trial using RAGT. Subjects trained 2 times per week for 10 weeks for a total of 20 training sessions. Five subjects with progressive MS have completed the RAGT protocol: 4 women and 1 man ranging in age from 33 to 63. The group has a range of Expanded Disability Status Scale scores from 4.5 to 6.5. Physical therapists individualized training intensity and RAGT characteristics to maximize benefits for each subject. Motor capacity outcomes (walking speed and endurance [2MWT]) and quality of life measures (Modified Fatigue Impact Scale [MFIS] and the Multiple Sclerosis Impact Scale 29 [MSIS-29]) were assessed at baseline and after the final training session (20th session). Subjects were monitored at each visit for adverse events. **Results:** There was no reported adverse event for any subject. Three of the 5 subjects had a 10% or greater increase in walking speed with an average improvement of 0.062 m/s. The group averaged 13% improvement in fast walking speed. Subjects had an average improvement of 10% on the MFIS and 15% on the MSIS-29. MFIS subscales revealed the greatest amount of improvement in the physical domain (44%). The MSIS-29 subscales indicated that individuals had a significant decrease in physical disability (18%). **Conclusions:** These 5 subjects with progressive MS tolerated the treatment dosage of 2 times per week for 20 weeks and did not experience any adverse event throughout the robotic training. Focused gait training using RAGT resulted in improvement in walking speed. Subjects reported that training reduced their disability and fatigue, enhancing their overall quality of life.

Supported by: None

Disclosure: *Nothing to disclose*

Keywords: *Complementary/alternative therapies in MS, Equipment in MS*

(REH27) Strategies to Foster Buy-in for Physical and Occupational Therapists: Engagement Across Multiple Sites and States for a Study on Tele-Exercise and Multiple Sclerosis (TEAMS)

Tracy Flemming Tracy,¹ Hui-Ju Young,² Tapan Mehta,² Mohanraj Thirumalai,³ Dorothy Pekmezci,⁴ Emily Riser,⁵ James Rimmer²

¹Rehabilitation, Tanner Center for MS, Birmingham, AL; ²School of Health Professions, University of Alabama at Birmingham (UAB)/Lakeshore Foundation Collaborative, Birmingham, AL; ³School of Health Professions, UAB, Birmingham, AL; ⁴School of Public Health, UAB, Birmingham, AL; ⁵Tanner Center and Foundation for Multiple Sclerosis, Birmingham, AL

Background: Tele-Exercise and Multiple Sclerosis (TEAMS) is a research project funded by the Patient-Centered Outcomes Research Institute (PCORI) that aims to deliver a 12-week exercise-based rehabilitation intervention to 820 people with multiple sclerosis (MS) who live in Alabama, Mississippi, and Tennessee. Participants are randomized into 1 of 2 study arms: TeleCAM and DirectCAM. The TeleCAM consists of 4

testing visits with the intervention delivered via videos accessible through a tablet application. The DirectCAM arm involves 4 testing visits and 20 clinic visits where the intervention is delivered. Physical and occupational therapists at 43 outpatient clinics across the 3 states are trained to deliver the intervention to enrolled participants. This researcher-provider model is an integral part of the study design and is critical to the sustainability and success of the project. **Objectives:** The purpose of this presentation is to describe a multitiered approach used to facilitate therapist engagement and form partnerships between the therapists, researchers, and study participants. **Methods:** A Therapist Manual of Operating Procedures (T-MOP) was developed for each clinic to ensure consistency of intervention delivery. The Clinical Research Coordinator used the T-MOP along with a rehabilitation guideline established by the Consortium of Multiple Sclerosis Centers (CMSC) to train clinicians at each clinic. Continuing Education Units (CEUs) were approved allowing each clinician to claim 4-6 hours for on-site training. Videos were created to provide an instructional guide on how to administer each outcome measure. GoToMeeting was used as a platform to deliver ongoing study updates. Information pertaining to participants is communicated through a HIPPA-compliant portal (Box). **Results:** 43 clinics across the 3 states and 86 therapists were trained for the study. Each clinic has a copy of the comprehensive (81 pp) T-MOP available in print form with access to a digital copy on Box. 18 GoToMeetings have been held and recorded for training and study updates, and each of the 86 trained therapists has used the outcome-measure videos. 43 therapists have access to Box and have uploaded approximately 1900 documents within 26 months. There are 823 participants enrolled in the study, and 731 have been baseline tested. **Conclusions:** Our implementation science researcher-provider model has established partnerships with therapists, research staff, and participants and has resulted in seamless communications in intervention delivery and data management across the participating clinics in 3 states. Investments in training and engagement of therapists should be considered critical to recruitment, enrollment, retention, and dissemination in research. The model can be adapted for other similar projects that require strong emphasis on researchers engaging clinicians in implementation science.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Engagement, MS and the caregiver/family

(REH28) Correlates of Change in First Trial Exposures Across 2 Days of Protective Steps Among Those with Multiple Sclerosis

Charles Van Liew,¹ Leland E. Dibble,² Bo Foreman,² Daniel S. Peterson^{1,3}

¹College of Health Solutions, Arizona State University, Phoenix, AZ; ²Physical Therapy and Athletic Training, University of Utah, Salt Lake City, UT; ³Research, Phoenix Veterans Affairs Medical Center, Phoenix, AZ

Background: Multiple sclerosis (MS) is a common, debilitating, neurodegenerative disorder that causes myriad symptoms. Gait and balance dysfunctions are common and manifest early in the disease, increasing fall risk. In particular, the ability to quickly and effectively react to a loss of balance is worse in people with MS. Therefore, improving reactive balance among those with MS is desirable. However, for maximum ecological validity, improvements in reactive balance through training would be demonstrable upon first loss-of-balance exposure. **Objectives:** The aim of this study is to evaluate first trial changes in people with MS before and after 1 day of protective stepping practice. The study also seeks to identify clinical correlates of first trial changes to begin evaluating for whom such training may provide benefit. **Methods:** Fourteen people with MS underwent 2 consecutive days of support-surface perturbations using an instrumented treadmill. Protective stepping outcomes were step length, step latency, and margin of stability. The backward step performance on the first trials on days 1 and 2 were compared, and difference scores were evaluated for relationships with correlates based on theoretical considerations. **Results:** There were no significant changes in first trial performance after training. However, some clinical and cognitive characteristics, such as mini-BESTest performance, improvement from day 1 to day 2 on the Symbol-Digits Modality test, type of MS diagnosis, and falls history were related with the amount of change individuals experienced. **Conclusions:** Although preliminary, these findings provide evidence that those with more favorable disease states may see more robust first-trial improvements after perturbation training. Greater doses, larger and more homogeneous samples, or longer delay between training and reassessment may be needed to understand the existence and relevance of first trial changes.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Rehabilitation

(REH29) Lifestyle Redesign for Multiple Sclerosis: A Case Series of Female Hispanic Patients

Vanessa Miller, Rebecca Cunningham

Mrs. T.H. Chan Division of Occupational Science and Occupational Therapy, University of Southern California, Los Angeles, CA

Background: Research indicates that the incidence rate and clinical presentation of multiple sclerosis (MS) varies between patients of different ethnic backgrounds. On average, the incidence rate of Hispanic patients with MS tends to be lower than that of non-Hispanic whites, while the average age of first reported MS symptom is earlier in the Hispanic population. Hispanic patients have higher incidence of cervical spinal lesions, mobility impairments, and optic neuritis at first MS-related event. Patients with MS of Hispanic descent may be at a higher risk of disability earlier in the disease process. Due to the identified clinical presentation and disability risks, it is critical to provide rehabilitation services that will support symptom and disease management. Evidence in the MS literature supports behavior and lifestyle interventions as critical components for symptom and disease management, as well as improved quality of life. Lifestyle Redesign is an occupational therapy (OT) approach that focuses on helping patients acquire health-promoting habits and routines to improve overall function, health, and quality of life, as well as to improve self-management of chronic conditions. This methodology involves education, occupational self-analysis, personal exploration, and goal setting interventions, to facilitate reflection and increase motivation for and the enactment of health-promoting behavior changes. **Objectives:** Describe the delivery of Lifestyle Redesign to address chronic disease and symptom management in patients with MS within an OT plan of care and provide a descriptive case series with clinical outcomes to demonstrate how this intervention can be applied clinically with Hispanic females with MS. **Methods:** The subjects included in this case series participated in an average of 11 OT sessions. All subjects are female, of Hispanic descent, and between the ages of 20 and 45. The Canadian Occupational Performance Measure (COPM), Multiple Sclerosis Quality of Life Inventory (MSQLI), and Health Related Quality of Life Short Form-36 (SF-36) were used at pre- and postintervention. **Results:** Clinically significant improvements occurred in the COPM overall performance and satisfaction scores, with patients demonstrating an average 5.3-point increase on performance and an average 7.2-point increase on satisfaction. On average, SF-36 scores improved in 7 subscales including emotional well-being, social functioning, and bodily pain, and MSQLI scores improved in 3 subscales including the MFIS. **Conclusions:** This case series supports the use of Lifestyle Redesign to address symptom and chronic disease management in Hispanic females with MS because of the demonstrated benefits in the areas of functional performance and symptom presentation. Additionally, this case series contributes to the broader evidence for the feasibility of Lifestyle Redesign services for neurologic populations.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Comprehensive care and MS, Lifestyle interventions, Management of activities of daily living in MS

(REH30) Cognitive Processing Speed as a Predictor of Motor Skill Learning in Healthy Adults and Persons with Multiple Sclerosis

Erin M. Edwards, Nora E. Fritz

Physical Therapy, Wayne State University, Detroit, MI

Background: Motor and cognitive deficits are frequently reported in individuals with multiple sclerosis (MS), resulting in a high incidence of rehabilitation enrollment. Presently, there is no way to predict whether a patient will benefit from a specific rehabilitation program and factors mediating exercise responsiveness in MS remain unknown. **Objectives:** This study aims to determine baseline cognitive and pathological predictors of an individual's ability to benefit from a balance-training program. We hypothesized that faster processing speed and increased myelin water fraction (MWF) in brain regions related to balance at baseline would result in greater automaticity at the trained task, as measured by the change in dual-task cost (DTC) following training. **Methods:** Four healthy participants and 1 participant with MS (1 male, 4 female; age 40 ± 14.3 year) underwent a magnetic resonance imaging examination and 4 consecutive days of balance training on the Neurocom Basic Balance Master. Each day involved a single session of twenty 2-minute blocks; participants performed weight shifts on a force platform in response to targets on a screen. Participants were evaluated pre- and post-training on their ability to perform a dual-task (Limit of Stability Test + N-back Test). **Results:** Following training, all participants demonstrated improvements in reaction time (14%), velocity (34%), directional control (5%), and target

accuracy (6%) on the challenging balance task. Reductions in DTC were seen across individuals, suggesting lower extremity motor skill training is feasible. Faster baseline processing speed on the Symbol Digit Modalities Test predicted reduced motor DTC in velocity ($r = 0.671$, 95% CI $-1.00, 0.00$) and directional control ($r = 0.783$, 95% CI $0.11, 1.00$) following training. Lastly, MWF values across brain regions related to balance were lower in the participant with MS compared to age-matched healthy controls. **Conclusions:** Data collection is ongoing; processing speed holds promise as a baseline indicator of the ability to benefit from a motor learning paradigm targeting postural control and balance. Given that demyelination is the pathologic hallmark of MS, and the MWF of the participant with MS is lower, our myelin water imaging data display feasibility to distinguish myelin-specific changes that may reflect exercise responsiveness. Identifying key variables associated with successful recovery of motor skills is a promising driving-force for improvements in the field of neurorehabilitation.

Supported by: None

Disclosure: Nothing to disclose

Keywords: *Imaging and MS, Rehabilitation methods*

RELAPSE THERAPY

(RTH01) Safety of Satralizumab Based on Pooled Data from Phase 3 Studies in Patients with Neuromyelitis Optica Spectrum Disorder (NMOSD)

Benjamin M. Greenberg,¹ Jerome de Seze,² Edward Fox,³ Albert Saiz,⁴ Takashi Yamamura,⁵ Carole Marcillat,⁶ Xiuqing Kou,⁷ Kristina Weber,⁸ Brian G. Weinschenker⁷

¹University of Texas Southwestern Medical Center, Dallas, TX; ²Hopital de Haute-pierre, Strasbourg, France; ³Central Texas Neurology Consultants, Round Rock, TX; ⁴Department of Neurology, Hospital Clinic and Institut d'Investigacions Biomèdiques August Pi i Sunyer, University of Barcelona, Barcelona, Spain; ⁵National Institute of Neuroscience, National Center of Neurology and Psychiatry, Tokyo, Japan; ⁶F. Hoffmann-La Roche Ltd, Basel, Switzerland; ⁷Mayo Clinic, Rochester, MN

Background: Interleukin-6 (IL-6) is implicated in the immunopathology of neuromyelitis optica spectrum disorder (NMOSD). Satralizumab, a humanized recycling monoclonal antibody that binds to the IL-6 receptor, demonstrated a reduction in NMOSD relapse risk in two phase 3 studies: SAKuraSky (satralizumab in combination with baseline immunosuppressants; trial registration: NCT02028884), and SAKuraStar (satralizumab monotherapy; NCT02073279). **Objectives:** To evaluate the safety of satralizumab vs placebo in a pooled population of patients with NMOSD from the SAKura studies, using the latest data from their open-label extension (OLE) periods. **Methods:** SAKuraStar and SAKuraSky are randomized studies comprising a double-blind (DB) period (satralizumab 120 mg every 4 weeks vs placebo) followed by an OLE period (satralizumab only). The combined DB and extension period was defined as the overall satralizumab treatment (OST) period (cutoff June 7, 2019). Safety was evaluated in the DB and OST periods and reported as adverse event (AE) rates per 100 patient-years (PYs). **Results:** The pooled DB population included 178 patients (satralizumab, $n = 104$; placebo, $n = 74$), and a total of 166 patients received satralizumab in the OLE. Mean and median satralizumab exposures in the OST period were 133.3 and 128.6 weeks. Rates of AEs and serious AEs were comparable between satralizumab and placebo groups in the DB period (AEs: 478.49 vs 506.51 events/100 PYs, respectively; serious AEs: 14.97 vs 17.98 events/100 PYs, respectively), and were consistent in the OST period. In the DB period, 4 patients (3.8%) in the satralizumab group and 6 (8.1%) in the placebo group withdrew from study due to an AE. Infection rates were lower with satralizumab vs placebo in the DB period (113.04 vs 154.85 events/100 PYs), with no increased risk of opportunistic infections. Infection rates with satralizumab were similar between the DB and OST periods. The injection-related reaction rate was higher with satralizumab vs placebo in the DB period (17.03 vs 8.99 events/100 PYs); injection-related reactions were mostly mild-to-moderate and did not lead to treatment discontinuation. No deaths or anaphylactic reactions were reported. **Conclusions:** In patients with NMOSD, satralizumab was well tolerated and showed a favorable safety profile. Results from the overall satralizumab treatment period, which expanded on the DB periods by adding data from the ongoing OLE periods, were consistent with the DB period results.

Supported by: None

Disclosure: Benjamin M. Greenberg: Abcam, Alexion, EMD Serono, Genentech, Novartis (consulting fee); Chugai, CELENE Nanomedicine, Guthy-Jackson Charitable Foundation for NMO, MedImmune, National MS Society, PCORI, Transverse Myelitis Association (contracted research). Jerome de Seze: Chugai, Roche (consulting fee). Edward Fox: AbbVie, Biogen, Celgene, EMD Serono, Sanofi Genzyme (personal fees and grants); Chugai (contracted research, personal fees and grants, personal fees and non-financial support – recruiting site for the

SAkuraStar study); Genentech, MedDay, Novartis, Roche, TG Therapeutics (consulting fee, contracted research, speakers' bureau). Albert Saiz: Bayer Schering, Biogen, Merck, Novartis, Roche, Sanofi, Teva (speakers' bureau). Takashi Yamamura: Alexion Pharma, Ono (consulting fee); Biogen, Novartis, Teijin Pharma (consulting fee, speakers' bureau); Chiome Bioscience, Miraca Holdings (contracted research); Chugai (consulting fee, contracted research, grant, speakers' bureau); CSL Behring, Mitsubishi Tanabe, Takeda, Teijin Home Healthcare (speakers' bureau). Carole Marcillat, Xiuqing Kou, Kristina Weber: F. Hoffmann-La Roche (salary). Brian G. Weinschenker: Alexion, Chugai, Mitsubishi Tanabe, Roche, Viela Bio (consulting fee); Hospices Civil de Lyon, MVZ Labor PD Dr Volkemann und Kollegen GbR, Oxford University, RSR Ltd (royalties).

Keywords: *Neuromyelitis optica spectrum disorder*

(RTH02) Adolescents with Neuromyelitis Optica Spectrum Disorder Achieved Similar Exposures and Favorable Safety Profile When Treated with the Adult Satralizumab Dosing Regimen

Cheryl Hemingway,¹ Hanna Silber Baumann,² Xiuqing Kou,³ Daniela Stokmaier,³ Patricia Sanwald Ducray,² Veronica G. Anania,⁴ Hajime Ito,⁵ H.-Christian von Buedingen,³ Sian Lennon-Chrimes⁶

¹Great Ormond Street Children's Hospital, London, United Kingdom; ²Roche Innovation Center, F. Hoffmann-La Roche Ltd, Basel, Switzerland; ³F. Hoffmann-La Roche Ltd, Basel, Switzerland; ⁴Genentech, San Francisco, CA; ⁵Chugai Pharmaceuticals Co Ltd, Tokyo, Japan; ⁶Roche Innovation Center, Welwyn Garden City, United Kingdom

Background: Interleukin-6 (IL-6) is implicated in the immunopathology of neuromyelitis optica spectrum disorder (NMOSD). Satralizumab, a humanized recycling monoclonal antibody that binds to the IL-6 receptor (IL-6R), demonstrated a reduction in NMOSD relapse risk in two phase 3 studies: SAKuraSky (satralizumab in combination with baseline immunosuppressants; trial registration: NCT02028884) and SAKuraStar (satralizumab monotherapy; NCT02073279). **Objectives:** To describe satralizumab exposure in adolescents with NMOSD to support dose selection. **Methods:** Patients in both studies ($N = 178$) received placebo or satralizumab 120 mg at weeks 0, 2, and 4, and every 4 weeks thereafter. Data on clinical and protocol-defined relapses (PDRs), aquaporin-4 autoantibody (AQP4-IgG) serostatus, safety end points, and pharmacokinetic (PK) and pharmacodynamic markers were evaluated in adolescent patients. A popPK model, developed using data from a phase 1 satralizumab trial (healthy volunteers) and both phase 3 studies, was used to analyze PK data. **Results:** Eight adolescent patients were enrolled in SAKuraSky (adolescents were not permitted in SAKuraStar). Seven were evaluated for efficacy (1 patient had PK data only). The mean age was 15.4 (range 13-17) years; mean weight (79.3 [range 47.5-140.4] kg) was similar to the adult population. Six patients were female; 3 patients were AQP4-IgG seropositive. The range of model-predicted exposures was similar to those in adults, correlating inversely with body weight, and not age. Treatment effects on soluble IL-6R levels, a marker of target engagement, were similar between adults and adolescents, with similar predicted median IL-6R occupancy (>95% maintained over the dose interval). One of 4 patients receiving satralizumab had a relapse (PDR, $n = 1$); all 3 patients receiving placebo relapsed (PDR, $n = 1$; clinical relapse, $n = 2$). The safety profile of satralizumab in adolescents was consistent with the adult patient population; no new safety signals were identified. **Conclusions:** These findings support the recommendation that adolescent patients with NMOSD receive the adult 120 mg loading and every 4 weeks maintenance regimen of satralizumab.

Supported by: None

Disclosure: Cheryl Hemingway: Biogen, Roche (consulting fee); Novartis (consulting fee, speakers' bureau). Hanna Silber Baumann: F. Hoffmann-La Roche, Innovation Center (salary). Xiuqing Kou, Patricia Sanwald Ducray, H.-Christian von Buedingen: F. Hoffmann-La Roche (salary). Daniela Stokmaier: Roche (salary). Veronica G. Anania: Genentech (salary). Hajime Ito: Chugai (salary). Sian Lennon-Chrimes: Roche Products Ltd (salary).

Keywords: *Neuromyelitis optica spectrum disorder*

(RTH03) Efficacy and Safety Outcomes from a Prospective Observational Registry of Repository Corticotropin Injection for Relapse of Multiple Sclerosis

Jeffrey Kaplan,¹ Tamara Miller,² Matthew Baker,³ Bryan Due,⁴ Enxu Zhao⁴

¹Kansas City Multiple Sclerosis Center, College Park Specialty Center, Overland Park, KS; ²Advanced Neurology of Colorado, Fort Collins, CO; ³Collier Neurologic Specialists, LLC, Naples, FL; ⁴Mallinckrodt Pharmaceuticals, Bedford, NJ

Background: Effective relapse treatment is critical for minimizing disability in patients with multiple sclerosis (MS). Repository corticotropin injection (RCI) is approved by the US Food and Drug Administration for the treatment of MS exacerbations. **Objectives:** This multicenter,

prospective, observational registry study aimed to characterize treatment response, recovery, and safety outcomes of RCI in the treatment of acute MS relapse. **Methods:** The following data were obtained upon initiation of RCI therapy (baseline) and again at various time points after. Clinical outcomes were assessed using the MS Impact Scale (MSIS-29v1), Expanded Disability Status Scale (EDSS), and Clinical Global Impression of Improvement (CGI-I) scale. Patient-reported outcomes were collected via the Work Productivity and Activity Impairment Questionnaire: MS (WPAI:MS) and Health Resource Utilization (HRU) questionnaire. Mean changes from baseline were evaluated at 2 and 6 months via 2-sided paired *t* tests. Serious and nonserious adverse events (SAEs/AEs) were reported throughout the study. **Results:** After treatment with RCI (N = 125), mean MSIS-29v1 physical subscale scores (primary end point) decreased from baseline (55.69) at 2 months (-7.99, *P* = .0002) and 6 months (-9.64, *P* < .0001). Post hoc analyses showed larger improvements in patients who received >5 doses of RCI (n = 23) vs ≤5 doses (n = 71) at 2 months (-10.74, *P* = .0180 vs -6.48, *P* = .0177) and 6 months (-14.62, *P* = .0415 vs -7.90, *P* = .0011). Mean EDSS scores decreased from baseline (3.92) at 2 months (-0.37, *P* < .0001) and 6 months (-0.45, *P* < .0001), with greater improvement in patients who received >5 doses vs ≤5 doses at 2 months (-0.50, *P* = .0068 vs -0.24, *P* = .0059) and 6 months (-0.64, *P* = .0430 vs -0.36, *P* = .0111). CGI-I scores improved in 63.38% of patients (*P* < .0001) at 2 months and 61.40% of patients (*P* < .0001) at 6 months postbaseline. Eighty-three AEs were reported by 35 patients (28%), and 16 SAEs were reported by 11 patients (8.8%). The most common AEs/SAEs were MS relapse (4% AE, 4% SAE) and urinary tract infection (3.2% AE, 1.6% SAE). WPAI:MS and HRU responses showed improvements from baseline for most end points at 2 and 6 months. **Conclusions:** Improvements in clinical MS scales and patient-reported measures of MS impact, along with the low incidence of AEs/SAEs, support the efficacy and safety of RCI as a treatment option for MS relapse. Treatment response showed greater improvements with >5 doses.

Supported by: None

Disclosure: *Jeffrey Kaplan:* Alexion, Allergan, Amgen, Biogen, EMD Serono, Eli Lilly, Teva (speakers' bureau). *Tamara Miller:* Acorda, Amgen, Mallinckrodt Pharmaceuticals, Reven (speaking and consulting fees); *Adamas, Elan, EMD Serono, Ipsen, Mallinckrodt Pharmaceuticals, Ono, Sun Pharma (contracted research); Allergan, Biogen, Genentech, Novartis, Sanofi Genzyme, Teva (contracted research, speaking and consulting fees). Matthew Baker:* Acorda, Avanir, Biogen, Celgene, Genentech, Mallinckrodt Pharmaceuticals, Sanofi Genzyme, Teva (consulting fee). *Bryan Due, Enxu Zhao:* Mallinckrodt Pharmaceuticals (employee).

Keywords: Comprehensive care and MS, Disease-modifying treatments in MS, Immunology and MS

(RTH04) A Prospective Clinical Trial Utilizing an iOS Mobile Application and Apple Watch to Manage Multiple Sclerosis and Predict Disease Relapses

Mary Ann Picone

Holy Name Medical Center, Teaneck, NJ

Background: Multiple sclerosis (MS) is a chronic neurodegenerative disease caused by the lesion forming demyelination of the central nervous system. Relapsing forms of MS are indicative of short periods of new or worsening symptoms, often associated with new lesions or increase demyelination. Early identification of disease relapses is crucial for facilitating earlier treatment and minimizing functional decline. The current standard of care relies on clinic visits at 3-, 6-, or 12-month intervals to assess a patient's disease status and facilitate necessary treatment. This format may lead to overlooked disease worsening. A new and expanding practice in the field of disease monitoring and treatment is the use of wearable data recording technology. Such technology can monitor physical, cognitive, and emotional status in patients with MS in real-time and may act as an additional resource in MS treatment. **Objectives:** The primary purpose of this study is to test the acceptability and feasibility of the comprehensive mobile application (app) "MS Health Connect" and its integrated wearable technology. Additionally, this study aims to investigate whether relapse indicators or objective evidence of disease progression can be identified and correlated to recorded data. The testing and use of this app aim to assess patient quality of life, MS severity, ambulatory changes, and fall detection. **Methods:** Study duration for each patient is 1 year. Patients will have access to a mobile and Apple watch-compatible version of the app. Patients are to wear the Apple watch daily and complete weekly or bimonthly mood, balance, and health surveys. Weekly 2-minute walk tests are also required. Walk tests are performed remotely using the inherent watch features. At baseline and every 3 months after onboarding, patients have a scheduled appointment in which patients will

complete cognitive, visual acuity, dexterity, and physical tests as well as complete quality-of-life and MS related surveys. Statistical analysis will be conducted for the validity of app components, the correlation of disease progression, and Apple watch-recorded data, and additional experimental aims related to the acceptability and feasibility of the app. **Results:** To date, 5 patients enrolled. Data are being collected. Further analysis is required. **Conclusions:** Positive feedback about the app and its components had been reported. Analysis of completed data is required for any conclusion to be supported.

Supported by: None

Disclosure: *Mary Ann Picone:* Biogen Inc (speakers' bureau).

Keywords: Complementary/alternative therapies in MS, Management of activities of daily living in MS, Wearable technology in MS

(RTH05) Characterization of the Pharmacokinetics and Pharmacodynamics of Satralizumab, a Recycling Antibody, to Support Once-Every-4-Weeks Dosing in Patients with Neuromyelitis Optica Spectrum Disorder

Sian Lennon-Chrimes,¹ Hanna Silber Baumann,² Gaelle Klingelschmitt,³ Xiujing Kou,³ Patricia Sanwald Ducray,² Veronica G. Anania,⁴ Hajime Ito,⁵ H.-Christian von Buedingen³

¹Roche Innovation Center, Welwyn Garden City, United Kingdom; ²Roche Innovation Center, F. Hoffmann-La Roche Ltd, Basel, Switzerland; ³F. Hoffmann-La Roche Ltd, Basel, Switzerland; ⁴Genentech, San Francisco, CA; ⁵Chugai Pharmaceuticals Co Ltd, Tokyo, Japan

Background: Interleukin-6 (IL-6) has been implicated in the immunopathology of neuromyelitis optica spectrum disorder (NMOSD). Satralizumab is a subcutaneously administered monoclonal antibody that binds to and blocks the IL-6 receptor (IL-6R). Satralizumab was engineered to be recycled back into circulation via the neonatal Fc receptor (FcRn), increasing its serum half-life and effecting prolonged inhibition of IL-6R signaling. **Objectives:** To define an effective, convenient, long-term dosing regimen for satralizumab in patients with NMOSD. **Methods:** The pharmacologic characteristics (pharmacokinetics [PK] and pharmacodynamics) of satralizumab were assessed in 72 Japanese healthy volunteers (HV); single dose, range 30-240 mg), 33 patients with rheumatoid arthritis (RA) (multiple doses, range 30-120 mg), and 104 patients with NMOSD from two phase 3 studies in NMOSD (SAkuraSky [trial registration: NCT02028884] and SAkuraStar [NCT02073279]; 120 mg loading, once every 4 weeks). A popPK model, based on HV and NMOSD data, was used to derive predictions for individual PK parameters. **Results:** Satralizumab provided significant inhibition of IL-6R signaling for 4 weeks; target engagement resulted in sustained increases in soluble IL-6R levels in HVs and patients with RA and NMOSD. In the NMOSD population, the PK of satralizumab was shown to be nonlinear, with an effective half-life of approximately 30 days at a dose of 120 mg; the median predicted IL-6R occupancy was maintained at >95% throughout the 4-week dose interval. Meaningful and comparable efficacy vs placebo was demonstrated in patients with NMOSD in both phase 3 studies: hazard ratio (95% CI) for reduction in protocol-defined relapse risk was 0.38 (0.16-0.88), *P* = .0184 in SAkuraSky; and 0.45 (0.23-0.89), *P* = .0184 in SAkuraStar. Satralizumab showed a favorable safety profile in patients with NMOSD when administered as monotherapy or in combination with baseline immunosuppressants. **Conclusions:** The recommended 120-mg loading and every-4-weeks maintenance regimen of satralizumab represents an effective, safe, and convenient treatment in NMOSD.

Supported by: None

Disclosure: *Sian Lennon-Chrimes:* Roche Products Ltd (salary). *Hanna Silber Baumann:* F. Hoffmann-La Roche, Innovation Center (salary). *Gaelle Klingelschmitt, Xiujing Kou, Patricia Sanwald Ducray, H.-Christian von Buedingen:* F. Hoffmann-La Roche (salary). *Veronica G. Anania:* Genentech (salary). *Hajime Ito:* Chugai (salary).

Keywords: Neuromyelitis optica spectrum disorder

SELF-CARE

(SEL01) Understanding Gaps in Knowledge Surrounding Flu Shots and Immunizations as They Relate to Multiple Sclerosis

Alexis Crispino

Multiple Sclerosis Association of America, Austin, TX

Background: Over the past 5 years the influx of US Food and Drug Administration-approved treatment therapies, and advances in symptom management for, multiple sclerosis (MS) has been astounding. Emphasis has been placed on a healthy lifestyle to maximize quality of life, and

many experts including the American Academy of Neurology and the Centers for Disease Control and Prevention agree that receiving vaccinations and flu shots are a part of staying healthy, including for most individuals with MS. However, this evolving landscape in care and treatment options has also heightened questions, misconceptions, and confusion surrounding the influence of flu shots and immunizations on MS. **Objectives:** Seeking to better understand gaps in knowledge and what patients with MS know/believe about flu shots and immunizations this assessment sought to analyze: 1) key areas of concern for receiving flu shots or immunizations; 2) how flu shots and immunizations are discussed with health care providers; and 3) understand current beliefs surrounding flu shots and immunizations. **Methods:** The Multiple Sclerosis Association of America (MSAA) developed and disseminated a 27-question survey on the topic of flu and immunizations as they relate to patients with MS that was emailed out to the MSAA client database. **Results:** 1926 patients with MS participated in the survey, with 32% of respondents reporting that they do not receive an annual flu shot and do not anticipate getting one this year. When asked why respondents do not receive flu shots, 36% opted concerns that flu shots are not good for people with MS, 32% are worried about side effects, and 28% do not trust or believe them to be safe. 68.26% reported an MS neurologist as their leading source of information, but still 37.74% said that they do not feel well informed about flu shots and 36.28% do not feel well informed enough about immunizations. 42% of respondents feel worried that if they receive an immunization or flu shot, they will have an adverse reaction and 38% believe that if they receive an immunization or flu shot it will interfere with their disease-modifying therapy or worsen their MS. Overall, 62.19% of individuals feel well informed about flu shots and immunizations while 37.81% feel that they need more information. **Conclusions:** These findings suggest that although experts agree that flu shots and immunizations are recommended for most individuals, there is still significant confusion among the MS patient community. "I believe that there are links between flu shots / immunizations and multiple sclerosis" reflected "not sure" responses of 38.81% and 42.51%, respectively. Finally, when asked how they would prefer to receive information about flu shots and immunizations in the future, leading responses were from their MS neurologist, general care practitioner, and through printed materials.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Healthy lifestyle while living with MS

(SELO2) Experimental Project Fashion Design and Social Inclusion: Multiple Women

Alice Estevo Dias,¹ Barbara Silva,² Barbara Gímenes Oliveira,² Caroline F. Mittermayer,² Giovana L. Pereira,² Grazielle M. Freitas,² Marcella A.M. Nogueira,² Rafaela L. Santos,² Rafaela F.B.N. Pereira,² Victoria V. Alvarenga,² Cristiane F. Mesquita²

¹Scientific Research, Brazilian Association of Multiple Sclerosis, Sao Paulo, Brazil; ²Design de Moda, Universidade Anhembi Morumbi, Sao Paulo, Brazil

Background: Multiple Women is an experimental project composed of an inclusive clothing collection for social occasions. Through social design guidelines, the work embraces textile mechanisms that aim to answer the physiological limitations of women with multiple sclerosis (MS). **Objectives:** To articulate fashion design and social inclusion in MS, with the purpose to develop garments with ergonomic adaptations. **Methods:** A group of 95 women with several types of MS, aged 20 to 40 with Expanded Disability Status Scale scores of 2.0 and 3.0, took part in structured interviews containing 12 questions about their daily requirements and clothing demands. A graphic and an infographic were elaborated as design visual tools to identify their current needs, the definition of the project parameters, and the planning of possible alternatives, along with a conceptual panel which associates the disease with textile features and panels of problems and solutions. **Results:** From the elucidation of the difficulties with zippers, buttons, closures, lashings, and thick fabrics, a collection of 6 drawn and 4 manufactured apparel was unleashed. The clothes present trespass, tailored zippers, and buttons and lightweight trim which collaborate with motor coordination. All of the clothes carry a QR Code that directs the user to an Instagram video that teaches how to wear the collection. **Conclusions:** The project proved to be appropriate to help quality of life and promote comfort, self-esteem, safety, and autonomy in the dressing process. The ergonomic alternatives show results that attend aesthetics and functionality demands that are scarce in the market. The challenge involves the creations in modeling and replacement of materials.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Complementary/alternative therapies in MS

(SELO3) An Assessment of the Feasibility of a Dyadic Physical Activity Intervention for Persons with Advanced Multiple Sclerosis and Their Family Caregivers: Work in Progress

Odessa J. McKenna,¹ Lara A. Pilutti,¹ Afolade Fakolade,¹ Marcia Finlayson,² Amy Latimer-Cheung,³ Mark S. Freedman,⁴ CMSC UOttawa Authors' Group

¹Interdisciplinary School of Health Sciences, University of Ottawa, Ottawa, ON, Canada; ²School of Rehabilitation Therapy, Queen's University, Kingston, ON, Canada; ³School of Kinesiology and Health Studies, Queen's University, Kingston, ON, Canada; ⁴University of Ottawa and the Ottawa Hospital Research Institute, Ottawa, ON, Canada

Background: Many people with MS (PwMS) and their caregivers (CGs) do not engage in sufficient physical activity to achieve important health benefits. Emerging evidence in other disease contexts (ie, Parkinson disease, Alzheimer disease) suggests that dyadic (ie, partnered) physical activity interventions may improve health and well-being for both care recipients and CGs. In multiple sclerosis (MS), physical activity interventions rarely focus on people with advanced disability or target PwMS-CG dyads. To address this important gap, we have developed a theory-based, manualized, dyadic intervention that incorporates evidence-based strategies for improving physical activity: Physical Activity Together for PwMS and Their CGs (PAT-MS). PAT-MS is a 12-week, teleconference-delivered intervention that includes education, guidance, and support from a trained activity coach, as well as behavior change techniques (eg, dyadic goal setting). **Objectives:** To evaluate the feasibility of PAT-MS for people with advanced MS and their CGs. **Methods:** A single-site, assessor-blinded, randomized controlled pilot feasibility trial, with 1:1 allocation into an immediate intervention or a delayed control condition. A target of 20 PwMS-CG dyads will be included. PwMS-CG dyads will receive 6 group teleconferencing sessions (~60 minutes) every other week for a period of 12 weeks. The group sessions will be interspersed with brief (~15 minutes) 1-on-1 support telephone calls in the weeks that the group sessions do not occur. Feasibility metrics will include process (eg, recruitment and retention rates), resource (eg, monetary costs and communication time), management (eg, time and accuracy of data collection/entry), and scientific assessment (eg, safety and participant experience). **Results:** Data collection is ongoing. Anticipated completion is March 2020. The main findings regarding intervention feasibility will be presented. **Conclusions:** PAT-MS is the first physical activity intervention to target both PwMS who have advanced disability and their CGs as active participants. The intervention presents a unique opportunity to increase physical activity behavior and improve the health outcomes of both PwMS and their CGs. The findings of this study will provide critical information on feasibility metrics that will inform and refine the design and delivery of subsequent stages of this research.

Supported by: None

Disclosure: Nothing to disclose

Keywords: MS and the caregiver/family, Physical activity

SYMPTOM MANAGEMENT

(SXM01) Effect of Nabiximols on Spasticity and Muscle Strength in Patients with Multiple Sclerosis Across 3 Randomized Controlled Trials

Francois Bethoux,¹ Kathryn Nichol,² Joanne Wagner,² Joris Berwaerts,² Daniel Checketts,³ Elizabeth Gardener,³ Karen Cartwright³

¹The Mellen Center for Multiple Sclerosis Treatment and Research, The Cleveland Clinic Foundation, Cleveland, OH; ²Greenwich Biosciences, Inc, Carlsbad, CA; ³GW Research Ltd, Cambridge, United Kingdom

Background: Spasticity is a common feature of multiple sclerosis (MS), especially in patients with longstanding illness. Medications that reduce spasticity may also reduce muscle strength, potentially impairing the ability to walk. Patients who do not respond adequately to conventional antispasticity medications need additional treatment options that improve spasticity without causing weakness. **Objectives:** Assess the relationship between changes in spasticity and muscle strength in lower extremities or mobility, using data from 3 randomized controlled trials (GWMS0106, GWSP0604, SAVANT) of nabiximols vs placebo in patients with spasticity due to MS inadequately controlled by antispasticity medications. **Methods:** Spasticity was evaluated using the Numerical Rating Scale (NRS) in all 3 trials, muscle strength using Motricity Index (MI) in GWMS0106 and GWSP0604, and mobility using timed 10-Meter Walk Test (10MWT) in GWSP0604 and SAVANT. Adjusted mean differences for change from baseline in outcome measures between nabiximols and placebo are summarized. Pearson correlation analysis was conducted to assess the association between change from baseline in spasticity and

change in strength or mobility for nabiximols and placebo groups separately. **Results:** This analysis included 184 patients from GWMS0106, 241 from GWSP0604, and 106 from SAVANT. The baseline mean (SD) Expanded Disability Status Scale score was 6.0 (1.42) in GWSP0604 and 5.9 (1.1) in SAVANT. In GWMS0106, nabiximols significantly improved mean NRS spasticity score (-0.52 points [95% CI: -1.029, -0.004]; $P = .048$), without significantly affecting the MI for legs (3.86 [-0.06, 7.78]; $P = .054$). In GWSP0604, nabiximols significantly improved mean NRS spasticity score from baseline vs placebo (-0.84 [-1.29, -0.40]; $P = .0002$), without significantly affecting the MI for legs (0.97 [-1.49, 3.42], $P = .439$) or the 10MWT results (-3.34 [-6.95, 0.26]; $P = .069$). In SAVANT, nabiximols significantly improved spasticity vs placebo (-1.9 [-2.73, -1.06]; $P < .0001$), without significantly affecting the 10MWT results (-1.71 [-3.84, 0.44]; $P = .11$). Pearson correlation coefficients were all under ± 0.30 (indicating negligible correlation) for the association between change in NRS and MI and for the association between change in NRS and 10MWT, except for the low positive correlation between NRS and 10MWT in the nabiximols group in SAVANT (0.326). **Conclusions:** The improvement in spasticity with nabiximols was not accompanied by the muscle weakness often observed with antispasticity medications or by a notable change in preferred walking speed.

Supported by: None

Disclosure: Francois Bethoux: Adamas Pharmaceuticals (contracted research); Biogen (speakers' bureau); GW Pharma (consulting fee); Springer Publishing (royalty). Kathryn Nichol: Greenwich Biosciences, Inc (salary). Joanne Wagner: Greenwich Biosciences/GW Pharmaceuticals (salary). Joanne Wagner, Karen Cartwright: GW Pharmaceuticals (ownership interest). Joris Beruwaerts, Elizabeth Gardener: GW Pharmaceuticals (salary). Daniel Checketts: GW Research Ltd (salary). Karen Cartwright: GW Pharmaceuticals (ownership interest/stockholder, salary).

Keywords: Complementary/alternative therapies in MS, Comprehensive care and MS, MS symptom management

(SXM02) MS Action Plan May Be Effective Tool Helping Patients with Acute Change in Multiple Sclerosis Symptoms

Gloria von Geldern,¹ Piper Paul,² Janet Piehl,³ Dennis Dietrich,⁴ Jan Shilling,⁵ Ted Brown,⁶ Kiren Kresa-Reahl,⁷ Annette Wundes,¹ Nicole Lauwers,⁸ Gary Stobbe,⁹ Kendra Yale,¹⁰ Piper Reynolds¹¹

¹Neurology, University of Washington, Seattle, WA; ²Nursing, Virginia Mason MS Center, Seattle, WA; ³Primary Care, UW Neighborhood Factoria Clinic, Bellevue, WA; ⁴Neurology, Advanced Neurology Specialists, Falls, MT; ⁵Kaiser Permanente, Seattle, WA; ⁶Evergreen Health MS Center, Kirkland, WA; ⁷Providence Multiple Sclerosis Center and Atara Biotherapeutics, Portland, OR; ⁸Neurology, St Vincent Neuroscience Center, Billings, MT; ⁹University of Washington, Seattle, WA; ¹⁰Swenson Healthcare, Tacoma, WA; ¹¹National Multiple Sclerosis Society, New York, NY

Background: The Greater Northwest Healthcare Provider Council of the National Multiple Sclerosis Society (NMSS) identified a need for patients with multiple sclerosis (MS) to be educated on what to do when they experience any new or worsening neurologic symptoms. Council members found that many of their patients with MS would check themselves into the emergency room (ER) or urgent care for symptoms that may have been better treated in the outpatient setting. **Objectives:** The objectives of the MS Action Plan are to 1) foster clear communication between patient and provider, 2) help patients avoid any undue trips to urgent care or ER, and 3) educate patients on what to look out for and when they need to contact their MS provider team. **Methods:** The Healthcare Provider Council started with a document in use at the Virginia Mason MS Center in Seattle and adapted it for all MS providers to use and share with their patients with MS to help them better understand their MS symptoms. The Virginia Mason document was adapted from the Asthma Action Plan, which is widely used throughout the country. The MS Action Plan was first distributed at the Greater Northwest Regional MS Summit annual professional education program to an audience of 80 MS care providers in March of 2019. Six months later, the Healthcare Provider Council created an online survey and distributed it to the Summit attendees to assess how the MS Action Plan is being used by providers and if they were finding it useful and effective. **Results:** Twenty providers responded to the survey. Of the respondents, 60% answered that they have used the MS Action Plan in their practice and 87% found it useful. When asked who provides the MS Action Plan to patients, 80% selected "provider" with "RN" being the second most commonly selected answer. The most common way the clinics are using the action plan is in hardcopy handed to patients, followed by placing the MS Action Plan in examination rooms for patients to take. **Conclusions:** The MS Action Plan could be an effective

tool in helping patients with MS better understand their MS symptoms, and feel confident in their decision making around when to be seen and when to wait. It has the potential for reducing the number of unnecessary ER visits and provides the patient with a clear plan of action should their symptoms change or worsen.

Supported by: None

Disclosure: Gloria von Geldern, Piper Paul, Janet Piehl, Jan Shilling, Nicole Lauwers, Kendra Yale, Piper Reynolds: Nothing to disclose. Dennis Dietrich: Adamas (contracted research); Biogen, Novartis (contracted research, speakers' bureau); Sanofi (speakers' bureau). Ted Brown: EMD Serono, Greenwich Pharmaceuticals (consulting fee); Merck (research grant); Novartis (speakers' bureau). Kiren Kresa-Reahl: Atara Biotherapeutics (salary from June 3, 2019, to present: they have a phase 1 MS trial but no medication on market); Biogen, Celgene, Genzyme, Novartis (speakers' bureau). Annette Wundes: AbbVie, Alkermes (contracted research); Biogen (consulting fee, contracted research). Gary Stobbe: Roche (contracted research).

Keywords: Comprehensive care and MS, MS and the caregiver/family, MS symptoms

(SXM03) Intrathecal Baclofen Therapy in Ambulatory and Nonambulatory Multiple Sclerosis Patients: A Single-Center Experience

Justin Abbatemarco,¹ Austin C. Griffin,² Noble Jones,^{3,4} Jennifer Hartman,⁵ Keith McKee,⁶ Zhini Wang,⁷ Sean Nagel,¹ Andre Machado,⁶ Francois Bethoux⁶

¹Neurological Institute, Cleveland Clinic, Cleveland, OH; ²Internal Medicine, The Cleveland Clinic Foundation, Cleveland, OH; ³Physical Medicine and Rehabilitation, Kessler Institute for Rehabilitation/Rutgers New Jersey Medical School, West Orange, NJ; ⁴Internal Medicine, Duke University, Durham, NC; ⁵Neurological Institute, Cleveland Clinic Mellen Center, Cleveland, OH; ⁶Neurological Institute, Cleveland Clinic Foundation, Cleveland, OH; ⁷Quantitative Health Sciences, Cleveland Clinic Foundation, Cleveland, OH

Background: Spasticity is a common cause of disability and diminished quality of life in patients with multiple sclerosis (MS). Intrathecal baclofen therapy (ITB) is an effective treatment option for patients with MS with severe spasticity that is refractory to oral drug administration, but there is limited evidence of its long-term efficacy and safety in ambulatory patients with MS. **Objectives:** This single-center, retrospective case series investigates the outcomes of ITB in ambulatory and nonambulatory patients with MS with medically intractable spasticity over a 5-year follow-up period. **Methods:** Data from the Mellen Center Intrathecal Baclofen Registry were analyzed retrospectively. All patients were diagnosed with MS and underwent an ITB test injection. Baseline demographics were collected along with outcome measures including Spasm Frequency Scale, Modified Ashworth scale (MAS), hip flexor strength, and walking speed on the Timed 25 Foot Walk. Group comparisons were done using 2-sample t test or Wilcoxon rank sum test, and logistic regression was used to assess the occurrence of complications. **Results:** 170 patients with MS underwent ITB infusion system implantation. The aggregate MAS score for the ambulatory cohort ($n = 87$) was significantly reduced from 13.5 ± 6.96 to 4.54 ± 4.18 at 5 years ($P < .001$) post ITB implantation. Similarly, spasm frequency (0-4 scale) was significantly reduced in ambulatory patients, from 1.71 ± 0.78 at baseline, to 0.77 ± 0.94 at 5 years ($P < .001$). The average ITB dose was lower for the ambulatory cohort compared to the nonambulatory cohort except at the 5-year follow-up visit. Among ambulatory patients at baseline, 56 (77.8%) were ambulatory at 1 year with no significant change in walking speed (baseline $0.45 \text{ m/s} \pm 0.30$ vs 1 year $0.38 \text{ m/s} \pm 0.39$ at 1 year, $P = .80$). At the 5-year follow-up point, 20 (41.7%) patients remained ambulatory with a walking speed of $0.21 \text{ m/s} \pm 0.37$ ($P < .001$). Longer disease duration (hazard ratio [HR] 1.04; 95% CI 1.01-1.07; $P = .018$), and lower hip flexor strength at baseline (HR 0.40; 95% CI 0.27-0.57; $P < .001$) were predictors for transition to nonambulatory status after ITB implantation. Complications were more common in the ambulatory ITB group ($n = 29$, 22.1%) compared to the nonambulatory group ($n = 10$, 8.0%) with an odds ratio of 3.30 (95% CI 2.17-5.02; $P = .017$). **Conclusions:** ITB is an effective therapy for reducing spasticity in ambulatory patients with MS without compromising walking speed in the short term, although we did observe a higher complication rate in this cohort. This study supports the use of ITB in carefully selected ambulatory patients with MS. Randomized, prospective studies are needed to provide more information on this important subject.

Supported by: None

Disclosure: Justin Abbatemarco, Austin C. Griffin, Noble Jones, Jennifer Hartman, Keith McKee, Zhini Wang, Sean Nagel: Nothing to disclose. Andre Machado: Abbott (consulting fee); Medtronic (fellowship support). Francois Bethoux: Adamas Pharmaceuticals (contracted research); Biogen (speakers' bureau); GW Pharma (consulting fee); Springer Publishing (royalty).

Keywords: Intrathecal baclofen, Management of activities of daily living in MS

(SXM05) Virtual Delivery of Mindfulness-Based Art Therapy (MBAT) to Improve Symptoms Among Adults with Multiple Sclerosis (MS)

Pamela K. Newland,¹ Rebecca D. Miller,² Eva Buch,³ Rebecca Lorenz⁴

¹Goldfarb School of Nursing at Barnes Jewish College, St Louis, MO; ²Art Therapy, St Mary of the Woods College, Terre Haute, IN; ³Goldfarb School of Nursing, St Louis, MO; ⁴University of Buffalo, Buffalo, NY

Background: Stress and fatigue in individuals with multiple sclerosis (MS) have been linked to a more severe course of the disease and weakened immune system. Mindfulness-based interventions (MBIs) have been found to reduce stress and fatigue and improve quality of life (QOL) in adults with MS. MBIs provided in an online format and led by a multidisciplinary team (nurse and art therapist) might strengthen overall effectiveness of MBIs. **Objectives:** The purpose of this pilot study was to compare the effect of virtually delivered meditation and mindfulness-based art therapy (MBAT) on the level of symptoms in a sample of adults with MS. We first beta-tested the protocol in 2 case subjects and collected feedback to identify areas needing improvement. We then plan to pilot the modified intervention to test feasibility, acceptability, and preliminary efficacy among a different sample of adults with MS. **Methods:** Art therapy and psychoeducation interventions were conducted by an MS nurse and an art therapist. Subjects engaged in mindfulness expressive arts interventions following a MBAT protocol on TEAMS, a video conferencing platform. Two facilitators led the interventions, as well as provided psychoeducational resources and feedback. We then interviewed the subjects to gather perceptions and feedback regarding the effectiveness of MBAT interventions in relieving stress and other symptoms. We also collected saliva cytokines, body temperature, and self-reported data on symptoms, physical function (standing balance, gait speed, minutes of daytime activity), and QOL. **Results:** At present, data analysis is ongoing. However, preliminary anecdotal feedback indicates that patients accept the MBAT interventions as accessible, easy to use, and helpful in reducing stress and fatigue. For example, 1 participant reported that creation of a particular art image enabled her to more effectively communicate the need for change with her spouse in a particular area of her life so that she could better manage her stress levels. Participants have also been responsive to psychoeducational resources given during sessions. **Conclusions:** Preliminary findings suggest that an MBAT intervention led by a nurse and art therapist in a virtual format may be an effective method for relieving stress and fatigue in adults with MS. A future larger study is warranted for this important intervention.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Management of activities of daily living in MS, Nursing management in MS

(SXM06) A Pilot Study of Mirabegron (Myrbetriq) and Behavioral Modification Including Pelvic Floor Exercise for Overactive Bladder in Multiple Sclerosis

Theodore R. Brown, Virginia I. Simnad

EvergreenHealth Multiple Sclerosis Center, EvergreenHealth, Kirkland WA, WA

Background: Urinary symptoms, including overactive bladder (OAB), are seen in up to 80% of patients with multiple sclerosis (MS). Most antispasmodics for OAB are anticholinergic, which may worsen cognition and constipation in MS. Mirabegron, approved for OAB in the general population, is a B3 adrenergic agonist, so it may be better tolerated in patients with MS. **Objectives:** To assess safety, tolerability, and efficacy of mirabegron in treating OAB in MS. **Methods:** Twenty-eight patients with MS and OAB were randomized 1:1 into placebo and treatment arms of this double-blind, placebo-controlled 10-week study. All patients received pelvic floor exercise training and watched a video about behavioral management of OAB. Patients in the control arm received placebo while the treatment arm received mirabegron (25 mg) with optional up-titration to 50 mg. Seventy-two-hour voiding diaries were used. The primary outcome measure was the change in OAB Symptom Composite Score (OAB-SCS), which assesses voiding frequency and urgency; higher scores mean worse symptoms. Secondary measures included number and volume of micturition, incontinence episodes, and patient assessments of OAB severity. **Results:** While both groups' scores were lower at final visit than at baseline, the final daily average OAB-SCS for the mirabegron group was 0.47 higher than that of the placebo group [95% CI = 0.047, 0.893, $P = .031$]. Thus, the mirabegron group had a worse primary outcome. On the other hand, for Subject Global Impression, the mirabegron group rated overall bladder control as significantly better relative to the placebo group [95% CI = 0.375, 2.381, $P = .009$]. Trends suggesting

treatment-related improvement in other secondary measures favored mirabegron on number of micturition and incontinence episodes per day, but these and other secondary outcomes did not reach statistical significance. Adverse events were limited and similar between groups, and there were no serious adverse events. Drug adherence was about 95%. **Conclusions:** Mirabegron was safe and well-tolerated in this MS population. Our mixed results do not demonstrate benefit from adding mirabegron to a program of behavior modification for OAB. Patients with MS may have neurologic differences from a general OAB population that reduce the responsiveness to beta-3 adrenergic agonists. A larger study population may elucidate the extent of the treatment effect on bladder function in patients with MS.

Supported by: None

Disclosure: Theodore R. Brown: AbbVie, Adamas, Astellas, Merck (contracted research); EMD Serono, Greenwich Pharma (consulting fee); Novartis (speakers' bureau). Virginia I. Simnad: Actelion, Biogen, Celgene, Novartis (contracted research).

Keywords: Bladder management, Comprehensive care and MS, Nursing management in MS

(SXM07) The Clinical Spectrum of Myelin Oligodendrocyte Glycoprotein (MOG) Antibody-Associated Demyelinating Disorders: Three Case Reports

Ameneh Zare, Aram Zabeti

Neurology and Rehabilitation Medicine, University of Cincinnati, Cincinnati, OH

Background: Myelin oligodendrocyte glycoprotein (MOG) antibody-associated disorders are distinct from multiple sclerosis (MS) and neuromyelitis optica spectrum disorder (NMOSD) with positive aquaporin-4-IgG (AQP4-Ab). The full clinical spectrum of this newly described condition is unknown. In this report we sought to describe clinical and imaging presentations of 3 MOG antibody-seropositive patients. **Objectives:** To present 3 cases of autoimmune MOG antibody-associated demyelination. **Methods:** Case 1: A 45-year-old woman presented with spasm and pain in lower back with radiation to both legs. Also, she had complaints about muscle spasm, fatigue, and tremor in right hand. Five years after the onset of first symptoms, she developed left retrobulbar optic neuritis (ON) which responded to intravenous (IV) steroids. Magnetic resonance imaging (MRI) confirmed left ON and mild cervical and thoracic spinal cord atrophy. A repeat MRI 24 months later showed a few nonspecific foci of T2/FLAIR signal hyperintensity in the subcortical white matter of the bilateral frontal lobes, which was not typical in appearance for MS. MS mimics work-up repeated and cell-based immunoassay revealed positivity for anti-MOG antibody with a titer of 1:100 and negativity for AQP4-Ab. She was started on mycophenolate mofetil. Case 2: A 57-year-old man who was previously diagnosed with relapsing MS presented with ON. He was treated with interferon beta-1a, glatiramer acetate, teriflunomide, and fingolimod. After 8 years from the onset of MS, the patient developed dysarthria, vertigo, and dysphagia initially treated with IV steroids but did not have a good recovery, and repeat MRI showed progression of multiple areas of enhancement in the posterior fossa. So, he was admitted to the hospital and received PLEX, which significantly improved his symptoms. Cell-based immunoassay was positive for anti-MOG antibody with a titer 1:1000. Subsequently his treatment was switched to rituximab, and his symptoms remained stable ever since. Case 3: A 45-year-old woman with first presentation as paresthesia and progressive cognitive decline was diagnosed with relapsing-remitting MS based on MRI and presence of oligoclonal band in cerebrospinal fluid. She had 2 major attacks that presented as weakness and numbness in lower extremities and face, which dramatically responded to high-dose steroids. Her last attack presented as ON that partially improved with IV steroids. MRI indicated several enhancing lesions in brain as well as cervical and thoracic spine. Cell-based immunoassay revealed positivity for anti-MOG antibody with a titer 1:100 and negativity for AQP4-Ab. Treatment with ocrelizumab started afterward. **Results:** Three cases are presented. **Conclusions:** MOG-associated demyelinating disorders seem to represent a new disease entity. Reporting MOG seropositive cases helps in expanding our knowledge about its clinical and imaging presentations and disease course and the best available treatment options.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Myelin oligodendrocyte glycoprotein antibody, MOG antibody associated demyelinating disorder, Cell-based immunoassay, Natural history of MS

WHITAKER RESEARCH TRACK

The late Dr. John N. Whitaker was a world-famous researcher in multiple sclerosis (MS). His work inspired many scientists to enter the field of MS and develop their skills and talents. Each year, the Consortium of Multiple Sclerosis Centers (CMSC) honors Dr. Whitaker's memory by presenting one award for innovative research by a young investigator. This year the CMSC and the Foundation of the CMSC will present a prize to a scholar whose work exemplifies the mission of this visionary leader in MS.

(WHI01) Characterizing the Acute Exercise Response in Nonambulatory People with Progressive Multiple Sclerosis

Thomas Edwards,¹ Dilpreet Bajwa,² Lisa A.S. Walker,^{3,4,5} Lara A. Pilutti⁶

¹School of Human Kinetics, University of Ottawa, Ottawa, ON, Canada; ²Interdisciplinary Health Sciences, University of Ottawa, Ottawa, ON, Canada; ³Psychology, Carleton University, Ottawa, ON, Canada; ⁴Psychology, University of Ottawa Brain and Mind Research Institute, Ottawa, ON, Canada; ⁵Neuroscience, Ottawa Hospital Research Institute, Ottawa, ON, Canada; ⁶Interdisciplinary School of Health Sciences, University of Ottawa, Ottawa, ON, Canada

Background: While the benefits of exercise training among people with multiple sclerosis (MS) who have mild-to-moderate disability (Expanded Disability Status Scale [EDSS] score = 1.0-6.5) have been established, the role of exercise for persons with MS who are nonambulatory (EDSS \geq 7.0) is largely unknown. This evidence is limited, in part, due to the need for specialized exercise modalities to deliver exercise for persons who are nonambulatory. An investigation of the physiological and symptomatic response associated with adapted exercise is needed as a first step toward understanding the potential benefits of exercise training for this population. **Objectives:** To characterize and compare the acute physiological and symptomatic response associated with 3 adapted exercise modalities in persons with MS who are nonambulatory. **Methods:** Ten participants (mean age, 62.5 ± 10.3 ; all progressive MS) with EDSS score 7.0-8.0 were recruited. Participants completed 1 baseline testing session to determine peak cardiorespiratory fitness (VO_{2peak}). Participants then completed 3 submaximal exercise sessions on adapted exercise modalities (arm ergometer, recumbent stepper, and functional electrical stimulation [FES] cycle). Physiological variables including oxygen consumption (VO_2) and heart rate were recorded continuously during exercise. Symptomatic outcomes including pain and fatigue were measured at 3 time points: before exercise, immediately postexercise, and 30 minutes postexercise. **Results:** All participants completed the FES cycling and recumbent stepper submaximal exercise sessions. Two participants could not complete the arm ergometer exercise, citing pain as the reason for cessation. All modalities elicited a VO_2 response exceeding 40% of VO_{2peak} and a heart rate response exceeding 70% of heart rate. This exercise intensity corresponds with moderate-vigorous physical activity, an intensity associated with cardiorespiratory fitness benefits. There was a significant increase in pain immediately after arm ergometer exercise compared to FES cycling ($P < .05$) and recumbent stepper exercise ($P < .05$). **Conclusions:** All adapted exercise modalities tested appear to be viable approaches for improving cardiorespiratory fitness in nonambulatory people with MS. However, symptomatic response should be considered when prescribing and developing exercise interventions. These findings will help to inform and optimize exercise prescription for people with MS who are nonambulatory.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Exercise training, Management of activities of daily living in MS

(WHI03) Youth with Multiple Sclerosis Have Low Fitness Levels

Samantha Stephens,¹ Jane E. Schneiderman,² Tara Berenbaum,¹ Marcia Finlayson,³ Robert W. Motl,⁴ E. Ann Yeh⁵

¹Department of Neurosciences and Mental Health, The Hospital for Sick Children, Toronto, ON, Canada; ²Clinical Research Services, Research Institute, Hospital for Sick Children, Toronto, ON, Canada; ³School of Rehabilitation Therapy, Queen's University, Kingston, ON, Canada; ⁴Physical Therapy, University of Alabama at Birmingham, Birmingham, AL; ⁵Department of Paediatrics, Division of Neurology, The Hospital for Sick Children, Toronto, ON, Canada

Background: We have demonstrated an association between higher levels of physical activity and lower levels of disease activity, depression, and fatigue in youth with multiple sclerosis (MS). Fitness is a separate construct which may be dependent on a multitude of factors, including

genetics and physical activity. The relationship between fitness, physical activity, disease activity, and symptoms in youth with MS is unknown. **Objectives:** 1) To determine if measures of cardiovascular fitness and strength differ between youth with MS and healthy controls. 2) To explore the relationship between fitness, physical activity, fatigue, and depression in youth with MS. **Methods:** Youth (age <18 years) with MS ($n = 19$) and controls ($n = 21$) completed a battery of tests that included a cycle ergometer test to determine aerobic capacity (VO_{2peak}), and NIH Toolbox protocols for a 2-minute walk test and grip strength, to determine endurance and muscular strength, respectively. We administered questionnaires and clinical and physiological assessments including PedsQL (fatigue), CES-DC (depression), Godin Leisure Time Exercise Questionnaire, height/weight/body mass index (BMI), and physical activity (accelerometry [Actigraph GT-3X]). Differences between MS and control groups were determined using student t tests. Pearson R and Spearman Rho were used to examine the relationship between fitness, physical activity, fatigue, and depression in MS (JASP version 0.10.2). **Results:** MS and control groups did not differ in sex or age (MS 16 years, control 16.5 years, $P = .09$; M:F MS 3:1, HC 2:1, $P = .585$). Youth with MS had a median Expanded Disability Status Scale (EDSS) score of 1.25 (interquartile range 0.5), disease duration of 1.7 (SD 2.0) years, and total relapses of 3.0, and higher BMI than controls ($T = -5.8$, $P < .001$). VO_{2peak} and 2-minute walk test differed in MS vs controls ($t = 4.6$, $P < .001$, Cohen $d = 1.5$) and ($t = 2.6$, $P = .02$, Cohen $d = 0.9$), respectively. Controls demonstrated higher average maximum workloads (170 W) ($U = 221$, $P = .002$, Cohen $d = 0.5$) and peak heart rate of 186 (no difference) bpm compared to 136 W and 172 bpm in MS. Measures of grip strength, rating of perceived exertion, physical activity, fatigue, and depression did not differ between the groups. VO_{2peak} was negatively correlated with MS relapses (Spearman $\rho = -0.5$, $\hat{P} = .04$ and grip strength with disability [EDSS] $r = -0.6$, $P = .03$). No significant correlations were found between indices of fitness with physical activity, fatigue, or depression in MS. **Conclusions:** Youth with MS demonstrate low levels of fitness, endurance, and muscular power compared to healthy controls. Lower levels of fitness and strength were also associated with increased disability and disease activity in MS, but not depression or fatigue. Further assessment of the relationship between physical fitness, depression, and fatigue is warranted due to the small sample size of the current study. Interventions that target aerobic exercise training are necessary for improving fitness in youth with MS, and this may decrease risk of future comorbidities.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Management of activities of daily living in MS, Physical fitness

(WHI04) Dietary Patterns and Health-Related Quality of Life of Individuals with Multiple Sclerosis

Babita Bisht,¹ Laurie K. Mischley,² Tyler J. Titcomb,¹ Joshua Farahnik,² Linda G. Snetselaar,³ Terry L. Wahls¹

¹Department of Internal Medicine, University of Iowa, Iowa City, IA; ²Bastyr University Research Institute, Bastyr University, Kenmore, WA; ³Department of Epidemiology, University of Iowa, Iowa City, IA

Background: Individuals with multiple sclerosis (MS) look for dietary changes to improve their disease outcome. Information regarding what specific dietary changes are being implemented by individuals with MS and if these changes affect quality of life (QOL) can be useful in shaping future research. **Objectives:** 1) To assess prevalence of MS-specific diets (eg, Wahls diet, Swank diet, vegan diet) and dietary patterns of individuals with MS. 2) To investigate effects of intake of certain food groups on relapses and self-reported QOL measure. **Methods:** Individuals with MS participated in an online survey and completed questions regarding intake of specific diets, frequency of intake of specific foods, relapses, and Patient-Reported Outcome Measurement Information System (PROMIS) based on past 6 months. PROMIS Global scores (range, 10-50) were used to measure QOL where higher scores reflect higher QOL. We estimated daily servings of foods from food frequency questions. In this cross-sectional analysis, we included 977 participants who provided complete data. **Results:** Individuals with MS with mean age 47.7 (SD 11) years and average 10 (SD 8.9) years since diagnosis participated in this study. Specific diets for MS were followed by 72% of participants, suggesting that most individuals with MS are implementing dietary changes. Most prevalent diets were Wahls (26%), paleolithic (16%), and anti-inflammatory (13%) diets. Some participants (11%) reported following multiple diets as well. Only 12% reported having a relapse in past 6 months. Mann-Whitney U tests showed that individuals who did not have any relapse had higher median daily intake of alcohol (0.08 vs 0.05 ounce equivalent, $P = .016$) than individuals who had relapses in prior 6 months. PROMIS Global average scores were 33.4 (SD 6.1).

Median (interquartile range) daily servings intake of different foods were as follows: total fruits and vegetables, 2.2 (1.2-3.4); total dairy, 0.1 (0-0.5); total grains, 0.4 (0.1-1.4); total meat and fish, 3.7 (1.9-5.1); total alcohol, 0.1 (0-0.4); and total eggs, 0.1 (0-0.4). These results show low intake of dairy, grains, eggs, and alcohol among individuals with MS. Spearman correlation did not show any significant relationship between PROMIS Global scores and dosage of different food intake. However, Mann-Whitney *U* test showed that individuals who were not taking dairy had higher PROMIS scores than those who were (median 36 vs 33, $P < .001$). Additionally, individuals taking alcohol had higher PROMIS scores than those who were not taking any (median 34 vs 32, $P = .001$). These results suggest potential beneficial effects of avoiding dairy and consuming alcohol on QOL of individuals with MS. **Conclusions:** Most individu-

als with MS report making dietary changes to improve disease outcome. Dairy and alcohol may affect relapse rate and QOL of these individuals. Future studies should assess role of dietary changes as complimentary treatment for MS.

Supported by: *None*

Disclosure: *Babita Bisht, Laurie K. Mischley, Tyler J. Titcomb, Joshua Farabnik, Linda G. Snetselaar: Nothing to disclose. Terry L. Wahls: BioCeuticals, Genova Diagnostics, Institute for Functional Medicine, MCG Health LLC (consulting fee); Dr Terry Wahls LLC, FBB Biomed Inc, The Wahls Institute PLC, TZ Press LLC (ownership interest); Penguin Random House (royalty).*

Keywords: *Complementary/alternative therapies in MS, Diet*

Information for Readers

The *International Journal of MS Care* (IJMSC) (ISSN 1537-2073) is the official peer-reviewed publication of the Consortium of Multiple Sclerosis Centers (CMSC). It was also adopted as the official publication of two sister organizations: the International Organization of Multiple Sclerosis Nurses (IOMSN) and the International Organization of Multiple Sclerosis Rehabilitation Therapists (IOMSRT). It is published bimonthly in January/February, March/April, May/June, July/August, September/October, and November/December by the Delaware Media Group, PO Box 937, Glen Rock, NJ 07452, USA. It contains peer-reviewed articles in areas of interest to multiple sclerosis health care professionals.

SUBSCRIPTIONS

Print subscriptions to IJMSC are available. Please direct subscription inquiries and notice of change of address to: Ryan Francia, Consortium of MS Centers, 3 University Plaza Dr., Suite 116, Hackensack, NJ 07601, USA; fax 862-772-7275; e-mail rfrancia@mscare.org.

ADVERTISING

Matters regarding commercial advertising should be directed to: Joseph J. D'Onofrio, Delaware Media Group, PO Box 937, Glen Rock, NJ 07452, USA; e-mail jdonofrio@delmedgroup.com. IJMSC does not accept classified advertising.

Acceptance and publication of an advertisement does not imply endorsement or approval of the company, product, or service by IJMSC or the CMSC.

INDEXING AND ABSTRACTING

IJMSC is indexed in the Cumulative Index to Nursing and Allied Health Literature (CINAHL), Rehabilitation & Sports Medicine Source (EBSCO Publishing), and Scopus. All articles are archived in PubMed Central, with citations and abstracts retrievable in PubMed.

EDITORIAL CONTENT AND SUBMISSION PROCEDURE

Statements and opinions in this publication are solely those of the authors and contributors and do not necessarily reflect the views of the Editorial Board, the CMSC, the publisher, or sponsors.

Manuscripts and editorial correspondence should be submitted using the online submission system located at <http://ijmsc.msubmit.net>. Editorial copy must conform to the guidelines contained in the IJMSC Information for Authors. General correspondence may be sent to the Editorial Office by e-mail at IJMSC@mscare.org, or by mail to Charlene Belsole, Mellen Center for MS Treatment and Research, U10, Cleveland Clinic, 9500 Euclid Ave., Cleveland, OH 44119, USA; or fax 216-445-0331.

PEER REVIEW

Submissions are sent for peer review to content experts in the particular area of the submitted manuscript. No less than two reviewers are assigned to each manuscript using the single-blind model. Members of the Editorial Board may participate in the peer-review process. Conflict-of-interest disclosures of the Editorial Board members are on file in the Editorial Office.

IJMSC WEBSITE

IJMSC is available online at <http://ijmsc.org>. The website includes current and past issues, published supplements, information for authors and reviewers, and a listing of Editorial Board members.

COPYRIGHT AND PERMISSIONS

Copyright ©2020 by the Consortium of Multiple Sclerosis Centers. All rights reserved. None of the contents of this publication may be reproduced without written permission of the publisher.

Permission requests to reproduce material from IJMSC should be sent to the Editorial Office by e-mail at IJMSC@mscare.org or by mail to Charlene Belsole, Mellen Center for MS Treatment and Research, U10, Cleveland Clinic, 9500 Euclid Ave., Cleveland, OH 44119, USA. Letters requesting permission should include the issue and page number on which the material appears and the name of the publication in which the material will be reproduced.

PRINTING INFORMATION

Printed in the United States of America at Quad/Graphics, 1700 James Savage Rd., Midland, MI 48642.

LATE BREAKING

The Consortium of Multiple Sclerosis Centers and its educational partner, the Delaware Media Group, the publisher of the International Journal of MS Care, are pleased to provide you with a selection of late-breaking abstracts. As you know, at the live meeting, this work is usually displayed in a separate section during the Poster Session. This year, due to circumstances beyond our control, these abstracts will be presented to you in a printed and digital format without prior review. Our leadership has opted to do this to provide you with the broadest selection of material to enhance your knowledge about multiple sclerosis and related problems with the broadest possible perspective.

Real-World Evidence Assessment of Betaseron (interferon beta-1b) Adherence Following the Introduction of the BETACONNECT Autoinjector

Oisín Butler,¹ Simone Werner,¹ Katsiaryna Holl,¹ Ann-Kathrin Frenz,² Eva-Maria Wicklein,³ Sandy Yeo,⁴ Mark Rametta⁵

¹Real World Evidence, Medical Affairs, Bayer AG, Berlin, Germany; ²Medical Affairs Statistics, Bayer AG, Berlin, Germany; ³Bayer Pharma AG, Berlin, Germany; ⁴Bayer (South East Asia) Pte Ltd, Singapore, Singapore; ⁵Medical Affairs, Bayer HealthCare Pharmaceuticals Inc., Whippany, NJ

Background: Maintaining adherence to disease-modifying therapies (DMTs) is challenging for chronic conditions such as multiple sclerosis (MS), and poor adherence in MS has been associated with increased risk of disease activity and higher resource utilization. For DMTs requiring parenteral self-administration, such as interferons, autoinjector devices may help patients overcome injection-related factors interfering with treatment adherence. The BETACONNECT device is an electronic autoinjector for the injection of interferon beta-1b (Betaseron), a DMT used in relapsing-remitting MS (RRMS). **Objectives:** This retrospective analysis of a US claims database evaluated adherence, as indexed by medication possession ratio (MPR) and persistence, to 2 subcutaneous DMTs, Rebif (interferon beta-1a) and Betaseron (interferon beta-1b), during the period prior to and following the introduction of the BETACONNECT autoinjector for Betaseron in patients with MS. **Methods:** Data from MarketScan, a US claims database, for patients with a medical claim for Rebif or Betaseron either prior to the introduction of BETACONNECT (October 2013–September 2015) or post approval and uptake of BETACONNECT (October 2016–September 2018), were evaluated. Patients aged ≥ 18 years with ≥ 1 confirmed MS diagnosis in the 12-month period prior to the first relevant DMT prescription within the defined time periods were included in this analysis. Four cohorts were defined: incident Rebif or Betaseron users over the 24-month period prior to the introduction of BETACONNECT or over the 24-month period following the introduction and uptake of BETACONNECT. Within each time period, patient populations were propensity score matched on demographic and clinical characteristics. MPR and persistence to both DMTs are described for the period prior to and following the introduction of BETACONNECT. **Results:** MPR: In the pre-BETACONNECT period, the proportion of users with $\geq 80\%$ MPR was higher for Rebif (90%, 95% CI 87%–93%) than Betaseron (83%, 95% CI 76%–88%), while in the post-BETACONNECT period, the proportion of users with $\geq 80\%$ MPR was higher for Betaseron (92%, 95% CI 85%–95%) than Rebif (86%, 95% CI 81%–91%). Persistence: In the pre-BETACONNECT period, median persistence in days was higher for Rebif (199, 95% CI 167–235) than for Betaseron (152, 95% CI 105–231), while in the post-BETACONNECT period, persistence was higher for Betaseron (327, 95% CI 244–440) than for Rebif (229, 95% CI 184–304). **Conclusions:** Following the introduction of BETACONNECT, Betaseron users were more adherent, with improved persistence and with $>90\%$ of users meeting 80% MPR, a threshold commonly used to define good adherence.

Supported by: None

Disclosure: Oisín Butler, Simone Werner, Katsiaryna Holl, Ann-Kathrin Frenz, Eva-Maria Wicklein: Bayer AG (salary). Sandy Yeo: Bayer (South East Asia) Pte Ltd (salary). Mark Rametta: Bayer HealthCare Pharmaceuticals Inc (salary).

Keywords: Disease-modifying treatments in MS, Equipment in MS, Patient empowerment in MS treatment

A Unique Case of a Patient with Tuberous Sclerosis and Recent Diagnosis of Neuromyelitis Optica

Maziar Eslami Farsani,¹ Jacob Rube,² Shitiz K. Sriwastava,³ Evanthia Bernitsas⁴

¹Neurology, Wayne State School of Medicine, Detroit, MI; ²Neurology, Wayne State University, Detroit, MI; ³Neurology, West Virginia University, Morgantown, WV; ⁴Neurology, Wayne State University School of Medicine, Detroit, MI

Background: Tuberous sclerosis complex (TSC) is an inherited neurocutaneous disorder that is characterized by pleomorphic features involving many organ systems, including multiple benign hamartomas of the brain, eyes, heart, lung, liver, kidney, and skin. Neuromyelitis optica (NMO, previously known as Devic disease) and NMO spectrum disorders (NMOSDs) are inflammatory disorders of the central nervous system characterized by severe, immune-mediated demyelination and axonal damage predominantly targeting optic nerves and spinal cord. Traditionally considered a variant of multiple sclerosis, NMO is now recognized as a distinct clinical entity based on unique immunologic features. **Objectives:** To report a unique case of TSC and NMO. **Methods:** Case report. **Results:** A 30-year-old woman who carries the diagnosis of TSC suddenly started to have tingling in the legs and vomiting. The vomiting was intractable and continued for a week. Her tingling progressed up to her waist and into her abdomen. She also started having episodic spasms in her arms and legs. A month later she started having difficulty walking and tingling in her hands. She underwent magnetic resonance imaging (MRI) of spine that showed longitudinally extensive spinal cord lesions in her upper cervical and lower thoracic cord. Differential diagnosis at that time included NMO, transverse myelitis, and astrocytoma in the setting of tuberous sclerosis, but then NMO IgG antibody was found to be positive both in serum and cerebrospinal fluid (CSF). The patient received 5 days of steroids, and, as she still had symptoms, she was given plasma exchange for 5 sessions and then continue treatment plan with rituximab.

Conclusions: This is a unique case of a patient who was diagnosed with TSC with clinical (seizures) and brain MRI features (cortical tubers) of TSC in her childhood and now diagnosed with NMO at the age of 30 with clinical features: longitudinally extensive spinal cord lesions and positive NMO IgG antibody in serum and CSF. Further studies are needed to find more information regarding the co-occurrence of a genetic disorder like TSC and an immune-mediated disease like NMO.

Supported by: None

Disclosure: Nothing to disclose

Keywords: Genetics and MS, NMO, TS

Shorter Infusion Time of Ocrelizumab: Primary Results from the ENSEMBLE PLUS Study in Patients with Relapsing-Remitting Multiple Sclerosis

Amy Perrin Ross,¹ Thomas Berger,² Robert Bermeil,³ Mark S. Freedman,⁴ Trygve Holmøy,^{5,6} Joep Killestein,⁷ Regine Buffels,⁸ Monika Garas,⁸ Karen Kadner,⁹ Marianna Manfrini,⁹ John McNamara,⁹ Hans-Peter Hartung¹⁰

¹Loyola University Chicago, Chicago, IL; ²Department of Neurology, Medical University of Vienna, Vienna, Austria; ³Mellen Center for Multiple Sclerosis, Cleveland Clinic, Cleveland, OH; ⁴University of Ottawa and the Ottawa Hospital Research Institute, Ottawa, ON, Canada; ⁵Department of Neurology, Akershus University Hospital, Lørenskog, Norway; ⁶Institute of Clinical Medicine, University of Oslo, Oslo, Norway; ⁷Department of Neurology, VU University Medical Center, Amsterdam, Netherlands; ⁸F. Hoffmann-La Roche Ltd, Basel, Switzerland; ⁹John McNamara Consulting Limited, Cambridge, United Kingdom; ¹⁰Department of Neurology, UKD, Center of Neurology and Neuropsychiatry and LVR-Klinikum, Heinrich-Heine University Düsseldorf, Düsseldorf, Germany

Background: Ocrelizumab is an intravenously administered anti-CD20 antibody approved for relapsing and primary progressive multiple sclerosis (MS). Shortening the infusion to 2 hours may reduce the total site stay from 5.5–6 hours (approved infusion duration including mandatory premedication/observation) to 4 hours, which may reduce patient and site staff burden. **Objectives:** To investigate the safety profile of ocrelizumab when administered over a shorter infusion period, using primary results from ENSEMBLE PLUS. **Methods:** ENSEMBLE PLUS is a randomized, double-blind substudy to the single-arm ENSEMBLE study (trial registration: NCT03085810). In ENSEMBLE, patients with early-stage relapsing-remitting MS (18–55 years; treatment-naïve; disease duration ≤ 3 years; Expanded Disability Status Scale score 0.3–5) receive ocrelizumab 600-mg infusions every 24 weeks for 192 weeks. In ENSEMBLE PLUS, ocrelizumab (600 mg) administered over the approved infusion time (3.5 hours; conventional duration), is compared with a 2-hour infusion (shorter duration); the initial infusion (2 \times 300 mg) duration remains unaffected. The ENSEMBLE PLUS primary end point is the proportion of patients with infusion-related reactions (IRRs) following the first randomized infusion (frequency/severity assessed during and 24 hours postinfusion). **Results:** As of September 2019, 291 and 289 patients were randomized to the conventional and shorter infusion groups, respectively. Following the first randomized infusion, 67 patients (23.1%) in the conventional and 71 patients (24.6%) in the shorter infusion group had IRRs, from which 17.9% vs 31.0% were throat irritation and 25.4% vs 23.9% were fatigue, respectively. Most IRRs were mild or moderate; $>98\%$ of all IRRs resolved without sequelae in both groups. No IRRs were life threatening, serious, or fatal; 1 severe IRR occurred in both the conventional

(laryngeal inflammation [second randomized dose]) and shorter duration groups (fatigue [first randomized dose]). No IRR-related discontinuations occurred. During the first randomized dose, 14 (4.8%) and 25 (8.7%) patients in the conventional and shorter infusion groups had IRRs leading to infusion interruption/slowness, respectively. **Conclusions:** Primary analysis suggests that the frequency/severity of IRRs are comparable between conventional and shorter infusions. No new safety signals were detected.

Supported by: None

Disclosure: *Amy Perrin Ross:* Nothing to disclose. *Thomas Berger:* Alexion, Teva (participation in clinical trials in MS sponsored by); Almirall, Biologix, Bionorica, Genzyme, MedDay, Teva-ratiopharm, TG Pharmaceuticals, UCB (participated in meetings sponsored by and received honoraria [lectures, advisory boards, consultations] from pharmaceutical companies marketing treatments for MS); Bayer, Biogen, Merck, Novartis, Sanofi Genzyme (institution has received financial support in past 12 months by unrestricted research grants, participated in meetings sponsored by, and received honoraria [lectures, advisory boards, consultations] from pharmaceutical companies marketing treatments for MS, participation in clinical trials in MS sponsored by); Octapharma, Roche (participated in meetings sponsored by and received honoraria [lectures, advisory boards, consultations] from pharmaceutical companies marketing treatments for MS, participation in clinical trials in MS sponsored by); Teva ratiopharm (institution has received financial support in past 12 months by unrestricted research grants). *Robert Bernel:* Biogen, F. Hoffmann-La Roche Ltd, Genentech Inc, Genzyme, Novartis (consulting fee). *Mark S. Freedman:* Actelion, Bayer HealthCare, Biogen, Clene Nanomedicine, F. Hoffmann-La Roche Ltd, Merck Serono, Novartis, Sanofi-Aventis (honoraria consultation fees, member of a company advisory board, board of directors or other similar group); Celgene, Chugai, EMD Canada, Genzyme, Pendopharm, Teva Canada Innovation (honoraria/consultation fees); Genzyme Canada (research grant); MedDay (member of a company advisory board, board of directors, or other similar group); Sanofi Genzyme (speakers' bureau). *Trygve Holmøy:* Biogen, Merck, Roche, Sanofi Genzyme (honoraria/consultation fees). *Joep Killestein:* Biogen, F. Hoffmann-La Roche Ltd, Merck, Novartis, Sanofi Genzyme, Teva (contracted research). *Regine Buffels:* F. Hoffmann-La Roche (salary). *Karen Kaden:* F. Hoffmann-La Roche Ltd (salary). *Monika Garas, Marianna Manfrini:* F. Hoffmann-La Roche (ownership interest, salary). *John McNamara:* AstraZeneca, F. Hoffmann-La Roche Ltd (consulting fee); John McNamara Consulting (ownership interest, salary). *Hans-Peter Hartung:* Bayer, Biogen, GeNeuro, Merck, Novartis, Roche, Sanofi Genzyme (honoraria for serving on steering and data monitoring committees).

Keywords: Disease-modifying treatments in MS, Ocrelizumab

Effect of Ofatumumab on Serum Immunoglobulin Levels and Infection Risk in Relapsing Multiple Sclerosis Patients from the Phase 3 ASCLEPIOS I and II Trials

Jérôme de Seze,¹ Amit Bar Or,² Jorge Correale,³ Anne H. Cross,⁴ Ludwig Kappos,⁵ Krzysztof Selmaj,⁶ Heinz Wiendl,⁷ Cecile Kerloeguen,⁸ Alexandra Goodyear,⁹ Ratnakar Pingili,⁹ Roseanne Sullivan,⁹ Ayan Das Gupta,¹⁰ Valentine Jehl,⁸ Dieter A. Häring,⁸ Martin Merschhemke,⁸ Stephen L. Hauser¹¹

¹University Hospital of Strasbourg, Strasbourg, France; ²Center for Neuroinflammation and Experimental Therapeutics and Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ³Institute for Neurological Research Dr. Raul Carrea, Buenos Aires, Argentina; ⁴Washington University School of Medicine, St. Louis, MO; ⁵Neurologic Clinic and Policlinic, Departments of Medicine, Clinical Research, Biomedicine and Biomedical Engineering, University Hospital and University of Basel, Basel, Switzerland; ⁶Center for Neurology, Lodz, Poland; ⁷University of Münster, Münster, Germany; ⁸Novartis Pharma AG, Basel, Switzerland; ⁹Novartis Pharmaceuticals Corporation, East Hanover, NJ; ¹⁰Novartis Healthcare Pvt. Ltd., Hyderabad, India; ¹¹Department of Neurology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA

Background: Ofatumumab, the first fully human anti-CD20 monoclonal antibody, demonstrated superior efficacy vs teriflunomide in patients with relapsing multiple sclerosis (RMS) in the phase 3 ASCLEPIOS I/II trials. A decline in serum immunoglobulin (Ig) levels was observed with other anti-CD20 therapies. **Objectives:** To determine serum IgG and IgM levels and investigate associations between IgG/IgM levels and risk of infections in ofatumumab-treated patients. **Methods:** In the ASCLEPIOS trials, patients received subcutaneous ofatumumab 20 mg on days 1, 7, and 14, week 4, and every 4 weeks thereafter or once-daily oral teriflunomide 14 mg for up to 30 months (average follow-up duration: 18 months). Serum IgG/IgM levels were monitored at baseline, weeks 4 and 12, and every 12 weeks thereafter (ofatumumab, n = 946; teriflunomide, n = 936). A notable decline in IgG/IgM levels was defined as 50% of the lower limit of normal (LLN) at any time [IgG, 3.5 g/L; IgM, 0.2 g/L]. Outcomes included the proportion of patients with IgG/IgM levels <50%

LLN, and association between low IgG/IgM levels and incidence of infections. **Results:** At week 120, no patients reached IgG levels <50% LLN with ofatumumab (median IgG [g/L]: ASCLEPIOS I and II, 10.57 and 9.57, respectively) or teriflunomide (10.01 and 9.65). The proportion of patients who reached IgM levels <50% LLN was 2.1% (n = 20/944) with ofatumumab (median IgM [g/L]: 0.91 and 0.59) and 0.6% (n = 6/933) with teriflunomide (0.84 and 0.92) at week 120. Of these patients, 5 had infections with ofatumumab, mostly nonserious (grade 1/2 in severity), except one grade 3 recurrent urinary tract infection, but all infections were resolved. One patient on teriflunomide who had nasopharyngitis had not recovered at the time of last follow-up. **Conclusions:** A reduction in serum IgG levels <50% LLN was not observed with either treatment. IgM levels showed reductions with both ofatumumab and teriflunomide treatments; there was no apparent association with increased rate of serious/nonserious infections in patients with RMS.

Supported by: None

Disclosure: *Jérôme de Seze:* Alexion, Allergan, Almirall, Bayer, Biogen, Chugai, CSL Behring, F. Hoffmann-La Roche Ltd, Genzyme, LFB, Merck, Novartis, Teva (consulting fee). *Amit Bar Or:* Atara Biotherapeutics, Biogen, Celgene, Genentech/Roche, GlaxoSmithKline, Mapi, MedImmune, Merck, EMD Serono, Novartis, Sanofi Genzyme (consulting fee, grant support, speakers' bureau). *Jorge Correale:* Merck-Serono, Biogen, Novartis, Roche, Genzyme (board member, contracted research). *Anne H. Cross:* Academic CME (CE provider) (speaking, preparation of slides for CME talk); Biogen, EMD Serono, Celgene, Genentech/Roche, Novartis, TG Therapeutics (consulting fee); EMD Serono, Genentech/Roche, Novartis, TG Therapeutics (contracted research); Genentech/Roche, Rockpointe (Potomac Center for Medical Education) (CE provider) (speakers' bureau); Novartis (fees for non-CME/CE services received directly from commercial interest or its agent, scientific advisory board for ASCLEPIOS I and II); Projects in Knowledge (CE provider) (preparation of educational manuscripts, activities). *Ludwig Kappos:* Actelion, Minoryx, Receptos, Santhera (consulting fee, steering committee; advisory board); Allergan, Almirall, CSL Behring, Pfizer (support for educational activities); Bayer HealthCare (consulting fee, speakers' bureau, steering committee; advisory board; support for educational activities; license fees for Neurostatus products; grants); Biogen, Merck (consulting fee, steering committee; advisory board; support for educational activities; license fees for Neurostatus products; grants); Celgene, Genzyme (consulting fee, steering committee; advisory board; support for educational activities); INNO-Swiss, Roche Research Foundation (grants); Novartis, Roche (consulting fee, contracted research, speakers' bureau, steering committee; advisory board; support for educational activities; license fees for Neurostatus products; grants); Sanofi (consulting fee, speakers' bureau, steering committee; advisory board; support for educational activities); Teva (consulting fee, steering committee; advisory board; support for educational activities; license fees for Neurostatus products). *Krzysztof Selmaj:* Novartis, Biogen, F. Hoffmann-La Roche Ltd, Merck, Synthron, Celgene, Genzyme (honoraria for advisory boards). *Heinz Wiendl:* Bayer HealthCare, Biogen, Fresenius Medical Care, GlaxoSmithKline (consulting fee, honoraria). *Cecile Kerloeguen, Alexandra Goodyear, Ratnakar Pingili, Roseanne Sullivan, Ayan Das Gupta, Valentine Jehl, Dieter A. Häring, Martin Merschhemke:* Novartis (salary). *Stephen L. Hauser:* Alector, Annexon, Bionure, Molecular Stethoscope, Symbiotix (scientific advisory board member); Alector, Annexon, Bionure, Molecular Stethoscope, Symbiotix, Neuron (stock options received); F. Hoffmann-La Roche (travel support/reimbursement and writing support for anti-CD20 meetings and presentations); Neuron (board of trustees member); Novartis (travel support/reimbursement for anti-CD20 meetings and presentations).

Keywords: Disease-modifying treatments in MS, Immunology and MS, Safety of DMTs in MS

Ofatumumab Versus Teriflunomide in Relapsing Multiple Sclerosis: Analysis of No Evidence of Disease Activity (NEDA-3) from the ASCLEPIOS I and II Trials

Stephen L. Hauser,¹ Amit Bar-Or,² Jeffrey A. Cohen,³ Giancarlo Comi,⁴ Jorge Correale,⁵ Patricia K. Coyle,⁶ Anne H. Cross,⁷ Jérôme de Seze,⁸ Xavier Montalban,^{9,10} Krzysztof Selmaj,¹¹ Heinz Wiendl,¹² Roman Willi,¹³ Bingbing Li,¹⁴ Dieter A. Häring,¹³ Krishnan Ramanathan,¹³ Martin Merschhemke,¹³ Ludwig Kappos¹⁵

¹Department of Neurology, UCSF Weill Institute for Neurosciences, University of California San Francisco, San Francisco, CA; ²Center for Neuroinflammation and Experimental Therapeutics and Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ³Department of Neurology, Mellen MS Center, Neurological Institute, Cleveland Clinic, Cleveland, OH; ⁴University Vita-Salute San Raffaele, Milan, Italy; ⁵Institute for Neurological Research Dr. Raul Carrea, Buenos Aires, Argentina; ⁶Department of Neurology, Stony Brook University, Stony Brook, NY; ⁷Washington University School of Medicine, St. Louis, MO; ⁸University Hospital of Strasbourg, Strasbourg, France; ⁹St Michael's Hospital, University of Toronto, Toronto, ON, Canada; ¹⁰Department of Neurology-Neuroimmunology, Centre d'Esclerosi Múltiple de Catalunya

(Cemcat), Hospital Universitario Vall d'Hebron, Barcelona, Spain; ¹¹Center for Neurology, Lodz, Poland; ¹²University of Münster, Münster, Germany; ¹³Novartis Pharma AG, Basel, Switzerland; ¹⁴Novartis Pharmaceuticals Corporation, East Hanover, NJ; ¹⁵Neurologic Clinic and Policlinic, Departments of Medicine, Clinical Research, Biomedicine and Biomedical Engineering, University Hospital and University of Basel, Basel, Switzerland

Background: Ofatumumab, the first fully human anti-CD20 monoclonal antibody, demonstrated superior efficacy vs teriflunomide in the phase 3 ASCLEPIOS I/II relapsing multiple sclerosis (RMS) trials. No evidence of disease activity (NEDA-3), a comprehensive composite measure, is commonly used to determine the treatment outcome in RMS. **Objectives:** To investigate the effect of subcutaneous ofatumumab 20 mg (monthly) vs oral teriflunomide 14 mg (once daily) in achieving NEDA-3, and to separately assess the annualized relapse rate (ARR) and gadolinium-enhancing (Gd+) T1 lesion activity in the pooled ASCLEPIOS I/II trials. **Methods:** We pooled data from the ASCLEPIOS I (n = 927) and II (n = 955) trials. Outcomes included NEDA-3 (defined as a composite of no 6-month confirmed disability worsening [6mCDW], no confirmed multiple sclerosis relapse, no new/enlarging T2 lesions, and no Gd+ T1 lesions) and its individual components in a modified full analysis set (modified FAS; logistic regression model). ARR by time intervals and Gd+ T1 lesions in the FAS (negative binomial model for both) were also analyzed. **Results:** The odds of achieving NEDA-3 with ofatumumab vs teriflunomide were >3-fold higher at month (M) 0-12 (47.0% vs 24.5% of patients; odds ratio [95% CI]: 3.36 [2.67; 4.21], *P* < .001) and >8-fold higher at M12-24 (87.8% vs 48.2% of patients; 8.09 [6.26; 10.45], *P* < .001). Over 2 years, a higher proportion of ofatumumab-treated patients were free from 6mCDW (91.9% vs 88.9%), relapses (82.3% vs 69.2%), and lesion activity (54.1% vs 27.5%) compared with teriflunomide. Ofatumumab significantly reduced ARR compared with teriflunomide at all cumulative time intervals: M0-3 (*P* = .011), and all subsequent time intervals from M0-27 (*P* < .001). Ofatumumab significantly reduced the mean number of Gd+ T1 lesions per scan by 95.9% compared with teriflunomide (mean [95% CI]: 0.02 [0.01; 0.03] vs 0.50 [0.42; 0.59]; *P* < .001). **Conclusions:** Ofatumumab increased the probability of achieving NEDA-3 and demonstrated superior efficacy vs teriflunomide in patients with RMS.

Supported by: None

Disclosure: Stephen L. Hauser: Alector, Annexon, Bionure, Molecular Stethoscope, Symbiotix (scientific advisory board member, stock options received); F. Hoffmann-La Roche (travel support/reimbursement and writing support for anti-CD20 meetings and presentations); Neurona (board of trustees member, stock options previously received); Novartis (travel support/reimbursement for anti-CD20 meetings and presentations). Amit Bar-Or: Atava Biotherapeutics, Biogen, Celgene, EMD Serono, Genentech/Roche, GlaxoSmithKline, Mapi, MedImmune, Merck, Novartis, Receptos, Sanofi Genzyme (consulting fee, grant support, speakers' bureau). Jeffrey A. Cohen: Multiple Sclerosis Journal (co-editor); Adamas, Convelo, Mylan, Population Council (consulting fee). Giancarlo Comi: Almirall SpA, Biogen, Biogen Italia Srl, Celgene Group, EXCEMED, F. Hoffmann-La Roche, Forward Pharma, Genzyme Corporation, MedDay, Merck KGaA, Merck Serono SpA, Novartis, Roche SpA, Sanofi Genzyme, Teva Italia Srl, Teva Pharmaceutical Industries Ltd (consulting fee); Genzyme Europe (consulting fee, speakers' bureau). Jorge Correale: Biogen, Genzyme, Merck-Serono, Novartis, Roche (board member, contracted research). Patricia K. Coyle: Accordant, Alexion, Bayer, Biogen, Celgene, Sanofi Genzyme, Serono, TG Therapeutics (consulting fee); Actelion, Alkermes, MedDay, National Institute of Neurological Disorders and Stroke (contracted research); Genentech/Roche, Novartis (consulting fee, contracted research). Anne H. Cross: Academic CME (CE provider) (speaking, preparation of slides for CME talk); Biogen, EMD Serono, Celgene, Genentech/Roche, Novartis, TG Therapeutics (consulting fee); EMD Serono, Genentech/Roche, Novartis, TG Therapeutics (contracted research); Genentech/Roche, Rockpointe (Potomac Center for Medical Education) (CE provider) (speakers' bureau); Novartis (fees for non-CME/CE services received directly from commercial interest or its agent, scientific advisory board for ASCLEPIOS I and II); Projects in Knowledge (CE provider) (preparation of educational manuscripts, activities). Jérôme de Seze: Alexion, Allergan, Almirall, Bayer, Biogen, Chugai, CSL Behring, F. Hoffmann-La Roche Ltd, Genzyme, LFB, Merck, Novartis, Teva (consulting fee, expert on advisory boards). Xavier Montalban: AbbVie, Biogen, MedDay, Merck, Novartis, Roche, Sanofi Genzyme, Teva Pharmaceutical (contracted research); Actelion, Alexion, Biogen, Celgene, EMD Serono, Genzyme, Immunic, MedDay, Merck, Nervgen, Novartis, Roche, Sanofi Genzyme, Teva Pharmaceutical, TG Therapeutics (consulting fee). Krzysztof Selmaj: Novartis, Biogen, F. Hoffmann-La Roche Ltd, Merck, Synthon, Celgene, Genzyme (honoraria for advisory boards). Heinz Wiendl: Bayer HealthCare, Biogen, Fresenius Medical Care, GlaxoSmithKline (consulting fee, honoraria). Roman Willi Bingel: Li, Dieter A. Häring, Krishnan Ramanathan, Martin Merschbemeke; Novartis (salary). Ludwig Kappos: Actelion (advisory board, steering committee); Bayer

HealthCare (advisory board, grants, license fees for Neurostatus products, speakers' bureau, steering committee, support for educational activities); Biogen (advisory board, grants, license fees for Neurostatus products, steering committee, support for educational activities); Celgene, Genzyme (advisory board, steering committee, support for educational activities); CSL Behring, Desitin, Pfizer, Allergan, Almirall (support for educational activities); INNO-Swiss, Roche Research Foundation, Swiss MS Society, Swiss National Research Foundation (grants); Merck (advisory board, grants, license fees for Neurostatus products, steering committee, support for educational activities); Minoryx, Receptos, Sanofi, Santhera (advisory board, steering committee); Novartis, Roche (advisory board, contracted research, grants, license fees for Neurostatus products, speakers' bureau, steering committee, support for educational activities); Sanofi (speakers' bureau, support for educational activities); Teva (advisory board, license fees for Neurostatus products, steering committee, support for educational activities).

Keywords: Comprehensive care and MS, Disease-modifying treatments in MS, Immunology and MS

Real-World Findings of Usability and Usefulness of Multiple Sclerosis Progression Discussion Tool: A Physician-Completed Digital Tool to Evaluate Early Signs of Disease Progression

Tjalf Ziemssen,¹ Gavin Giovannoni,² Enrique Alvarez,³ Virender Bhan,⁴ Carrie M. Hersh,⁵ Olaf Hoffmann,⁶ Celia Oreja-Guevara,⁷ René Robles Cedeño,⁸ Maria Trojano,⁹ Patrick Vermersch,¹⁰ Pamela Dobay,¹¹ Mudeer Khwaja,¹² Bianca Stadler,¹² Thomas Hach,¹² Daniela Piani-Meier,¹² Jason Burton¹³

¹Department of Neurology, University Clinic Carl-Gustav Carus, Dresden, Germany; ²Queen Mary University of London, London, United Kingdom; ³Geriatric, Research, Education, and Clinical Center, VA Eastern Colorado Healthcare System, Aurora, CO; ⁴University of British Columbia, Vancouver, BC, Canada; ⁵Lou Ruvo Center for Brain Health, Cleveland Clinic, Las Vegas, NV; ⁶Department of Neurology, Alexianer St. Josef Hospital, Potsdam, Germany; ⁷Hospital Clinico San Carlos, Madrid, Spain; ⁸Neuroimmunology and Multiple Sclerosis Unit, Girona, Spain; ⁹Department of Basic Medical Sciences, Neuroscience and Sense Organs, University of Bari, Bari, Italy; ¹⁰Department of Neurology, University of Lille, CHU Lille, Lille, France; ¹¹Real World Evidence Solutions, IQVIA Technology and Services, Basel, Switzerland; ¹²Novartis Pharma AG, Basel, Switzerland; ¹³Consultant Neurologist, Australian Neuro-muscular Research Institute, Perth, Australia

Background: MSProDiscuss is a validated physician-completed tool based on a set of well-validated questions that include information on multiple sclerosis (MS) relapses, symptoms, and impacts experienced by the patient within the past 6 months. The tool's traffic light system-linked output is meant as an aid for discussing the signs of MS disease progression. The tool is available online at www.msprodiscuss.com. **Objectives:** To report physician findings on usability and usefulness testing of the MSProDiscuss tool while discussing disease progression with patients in the real-world setting. **Methods:** An online qualitative survey was undertaken in 34 countries. Health care practitioners (HCPs) completed individual questionnaires (i) after using MSProDiscuss during face-to-face patient consultations and a final questionnaire (f) to capture the overall experience on the tool. The HCPs also provided general feedback and recommendations for improving the tool. **Results:** In total, 301 HCPs (including 23 MS nurses and 6 neurology nurse practitioners) tested the tool in 6974 patients with MS. The time taken to complete MSProDiscuss during a regular consultation was 1-4 minutes in 97% (i) to 98% (f) of the time. In 94% (i) to 97% (f) of cases, HCPs agreed that patients were able to comprehend the questions from the tool. HCPs were willing to use the tool again in the same patient 91% (i) of the time. MSProDiscuss was useful in discussing MS symptoms and its impact on daily activities (88% i / 92% f) and cognitive function (79% both i and f) and in discussing progression in general (88% i / 90% f). While completing the final questionnaire, 95% of HCPs agreed that the questions were similar to those asked in regular consultation. MSProDiscuss was also found helpful for understanding the impact of MS symptoms on daily activities (91%) and cognitive function (80%). Overall, 92% of the HCPs would recommend MSProDiscuss to a colleague. Regarding integration of MSProDiscuss into their clinical practice, 92% of HCPs think that it is feasible and 86% are willing to integrate. Key recommendations were to allow for longitudinal follow-up, expand on cognitive assessments, and provide a patient-completed version; these are considered in the updated version of MSProDiscuss. **Conclusions:** The findings from this real-world study suggest that MSProDiscuss is a usable and useful tool to facilitate physician-patient discussion on disease progression in daily clinical practice by capturing structured disease history.

Supported by: None

Disclosure: Tjalf Ziemssen: Almirall, Bayer, British American Tobacco, Biogen, Celgene, Sanofi Genzyme, Merck, Novartis, Roche, Vitaccess, Teva (speakers'

bureau). Gavin Giovannoni: AbbVie, Actelion, Atara Bio, Biogen, Celgene, Sanofi Genzyme, Genentech, GlaxoSmithKline, Merck-Serono, Novartis, Roche, Teva (consulting fee); Biogen, Roche, Merck, Merck-Serono, Novartis, Sanofi Genzyme, Takeda (contracted research); Elsevier (editor compensation on Mult Scler Relat Disord). Enrique Alvarez: Actelion, Biogen, Celgene, EMD Serono, Genentech, Genzyme, Novartis, Teva, TG Therapeutics (consulting fee); Biogen, Genentech, Novartis, Rocky Mountain MS Center (contracted research). Vivender Bhan, Jason Burton: Novartis (consulting fee). Carrie M. Hersh: Biogen, Novartis, EMD Serono, Genzyme, Genentech (consulting fee); Biogen, Novartis, Genentech (contracted research); Genzyme (speakers' bureau). Olaf Hoffmann: Alexion, Novartis, Roche, Sanofi, Teva, Bayer, Biogen, Boehringer Ingelheim, Celgene, Daiichi-Sankyo, Merck (consulting fee, contracted research, travel). Celia Oreja-Guevara: Alexion Pharmaceuticals, Merck, Novartis, Roche, Sanofi (consulting fee, contracted research). René Robles Cedeño: Biogen, Roche, Novartis, Merck, Sanofi Genzyme, Teva (consulting fee). Maria Trojano: Biogen, Merck, Roche, Novartis (consulting fee, speakers' bureau). Patrick Vermersch: Biogen, Novartis, Teva, Celgene, Roche, Sanofi (consulting fee); Biogen, Novartis, Teva, Celgene, Roche, Sanofi (registration, travel and accommodations for meetings); Roche, Sanofi (contracted research). Pamela Dobay: IQVIA Technology and Services AG (salary). Mudeer Khwaja, Bianca Stadler, Thomas Hach, Daniela Piani-Meier: Novartis (salary).

Keywords: Comprehensive care and MS, Symptoms of disease progression

Efficacy and Safety of the Bruton's Tyrosine Kinase Inhibitor (BTKI) Evobrutinib in Relapsing Multiple Sclerosis over 108 Weeks: Open-Label Extension to a Phase 2 Study

Xavier Montalban,^{1,2} Douglas L Arnold,^{3,4} Martin S. Weber,⁵ Ivan Staikov,⁶ Karolina Piasecka-Stryczynska,⁷ Emily C. Martin,⁸ Matthew Mandel,⁹ Victor Ona,⁹ Fernando Dangond,⁹ Jerry S. Wolinsky¹⁰

¹University of Toronto, St. Michael's Hospital, Toronto, ON, Canada; ²Vall d'Hebron University Hospital, Barcelona, Spain; ³NeuroRx Research, Montreal, QC, Canada; ⁴Montreal Neurological Institute, McGill University, Montreal, QC, Canada; ⁵Institute of Neuropathology and Department of Neurology, University Medical Center, Göttingen, Germany; ⁶Department of Neurology, Acibadem City Clinic Tokuda Hospital, Sofia, Bulgaria; ⁷Department of Neurology and Cerebrovascular Diseases, Poznan University of Medical Sciences, Poznan, Poland; ⁸EMD Serono Research & Development Institute, Inc. (a business of Merck KGaA, Darmstadt, Germany), Billerica, MA; ⁹Global Clinical Development Center, EMD Serono, Inc. (a business of Merck KGaA, Darmstadt, Germany), Billerica, MA; ¹⁰McGovern Medical School, UTHealth, Houston, TX

Background: In a phase 2 randomized controlled trial (RCT; trial registration: NCT02975349) in patients with relapsing multiple sclerosis (MS), evobrutinib 75 mg twice daily reduced total T1 gadolinium-enhancing lesions (primary end point) and annualized relapse rate (ARR) over 24 weeks vs placebo, with efficacy maintained through week 48. **Objectives:** To assess long-term efficacy and safety in the open-label extension (OLE) of the study. **Methods:** In the 48-week double-blind period, patients received evobrutinib 25 mg or 75 mg once daily, evobrutinib 75 mg twice daily, open-label dimethyl fumarate (240 mg twice daily), or placebo for the first 24 weeks; all arms continued with the original treatment assignment until 48 weeks, except placebo patients who were switched to evobrutinib 25 mg once daily. At week 48, all patients could enter the OLE, where treatment was initially evobrutinib 75 mg once daily (for approximately 48 weeks, median) before switching to 75 mg twice daily. The OLE assessed long-term efficacy (0-108 weeks) and safety (60-week OLE) of evobrutinib. **Results:** Of 267 randomized patients, 213 (79.8%) completed 108 weeks of treatment (48-week main study, 60-week OLE). For patients who received evobrutinib 75 mg twice daily in the double-blind period, the ARR (95% CI) was 0.11 (0.04-0.25) at week 48 and 0.12 (0.06-0.22) for the 108-week period. Evobrutinib was generally well tolerated, with the safety profile maintained during the 60-week OLE. Transient elevated liver aminotransferases, reported in the 48-week double-blind period, were not observed in the OLE. **Conclusions:** Efficacy and safety were maintained long-term. Two phase 3 RCTs evaluating the efficacy and safety of evobrutinib in patients with relapsing MS commence in 2020.

Sponsored by: EMD Serono, Inc, Rockland, MA, USA, a business of Merck KGaA, Darmstadt, Germany

Disclosure: Xavier Montalban: Actelion, Alexion, Bayer, Biogen, Celgene, EMD Serono, EXCEMED, Genzyme, Immunic, MedDay, Merck, MS International Federation, Mylan, Nervgen, National MS Society, Novartis, Roche, Sanofi Genzyme, Teva Pharmaceutical, TG Therapeutics (consulting fee). Douglas L. Arnold: Acorda, Adelphi, Alkermes, Biogen, Celgene, Frequency Therapeutics, Genentech, Genzyme, F. Hoffmann-La Roche, Immune Tolerance Network, Immunotec, MedDay Pharmaceuticals, Merck Serono, Novartis, Pfizer, Receptos,

Sanofi-Aventis (consulting fee); NeuroRx Research (ownership interest). Martin S. Weber: Bayer, Biogen, F. Hoffmann-La Roche, Genzyme, Merck Serono, Novartis, Teva (consulting fee). Ivan Staikov: Adapt, Bayer, Boehringer Ingelheim, Ewopharma-Biogen, F. Hoffmann-La Roche, Gedeon Richter, Merck Serono, Mylan, Penumbra, Pfizer, Polpharma, Sanofi Genzyme, Shire, Teva (consulting fee). Karolina Piasecka-Stryczynska: Biogen, F. Hoffmann-La Roche, Merck Serono, Sanofi-Aventis, Teva (consulting fee). Emily C. Martin, Matthew Mandel, Victor Ona, Fernando Dangond: EMD Serono, Inc (a business of Merck KGaA, Darmstadt, Germany) (salary). Jerry S. Wolinsky: Acorda Therapeutics, Actelion, Alkermes, Brainstorm Cell Therapeutics, Celgene, EMD Serono, GeNeuro, GW Pharma Ltd, MedDay Pharmaceuticals, NervGen Pharma Corp, Novartis, Otsuka, PTC Therapeutics, Roche/Genentech, Sanofi Genzyme (consulting fee); Millipore Corporation (paid to UTHealth), University of Texas (royalty).

Keywords: Disease-modifying treatments in MS

Natalizumab-Treated Patients with Relapsing-Remitting Multiple Sclerosis Report Better "Feel-Good" Outcomes on Key Physical, Emotional, and Cognitive Domains Compared to Other Disease-Modifying Therapies

John Foley,¹ Regina Berkovich,² Mark Gudesblatt,³ Carl de Moor,⁴ Stephanie Licata,⁴ Pei-Ran Ho,⁵ Karthik Bodhinathan,⁴ Robin Avila,⁴ Lily Lee⁶

¹Rocky Mountain MS Clinic, Salt Lake City, UT; ²Neurology Program, Los Angeles County and University of Southern California Medical Center, Los Angeles, CA; ³South Shore Neurologic Associates, Patchogue, NY; ⁴Biogen, Cambridge, MA; ⁵Biogen (at time of this study), Cambridge, MA; ⁶Biogen (at time of this analysis), Cambridge, MA

Background: Natalizumab is an efficacious therapy for patients with relapsing-remitting multiple sclerosis (RRMS). Some patients have reported a "feel-good" experience on natalizumab. Prior qualitative interviews of natalizumab-treated patients have suggested that the feel-good experience may be associated with increased energy and improved emotional and cognitive functioning. **Objectives:** To describe survey results assessing patients self-reporting a "feel-good" experience while receiving natalizumab or other disease-modifying therapies (DMTs). **Methods:** Surveys were administered to adult patients with RRMS through MyMSTeam (part of the MyHealthTeam application). Patients were asked about their current DMT use and its "feel-good" effect assessed by self-reported improvements in physical, emotional, or cognitive domains. Natalizumab vs other-DMT patient responses were corrected for multiple measures and compared using *t* test and log-rank tests. **Results:** Patients receiving natalizumab (n = 95) or other-DMT (n = 252) were included. Time since RRMS diagnosis was <6 years in 29% (natalizumab) and 35% (other-DMT) and >15 years in 29% (natalizumab) and 27% (other-DMT). Significantly higher percentages of natalizumab than other-DMT patients reported that they "feel good" on their DMT (63% vs 45%; *P* = .001). Physical benefits were reported by 78% of natalizumab and 67% of other-DMT patients (*P* = .017), with significantly higher rates of improved energy (23% vs 12%; *P* = .011) and coordination (22% vs 12%; *P* = .017) for natalizumab vs other-DMT. Comparison of patients on natalizumab vs other-DMT indicated significantly higher scores on emotional components involving improved motivation (25% vs 13%; *P* = .007) and happiness (24% vs 11%; *P* = .004); and improvement on cognitive component involving organizing one's thoughts (24% vs 14%; *P* = .021). Among natalizumab patients, 59 out of 95 (61%) scored high (either 4 or 5 for "agree"/"strongly agree") on feel-good effect, and also reported significantly increased energy, balance, coordination, motivation, happiness, and cognitive benefits (each *P* < .001). **Conclusions:** These real-world patient-centric survey results suggest that natalizumab is associated with a feel-good experience compared with other-DMTs. Increased physical, emotional, and cognitive functioning were more common in patients receiving natalizumab than in patients receiving other DMTs, consistent with qualitative interviews. These results are limited by the subjective nature of the survey responses.

Supported by: Biogen

Disclosure: John Foley: Biogen, EMD Serono, Genzyme, Genentech-Roche (consulting fee); Biogen, Genentech-Roche, Novartis, Genzyme, Mallinckrodt, Adamas (research support). Regina Berkovich: Alexion, Bayer, Biogen, Celgene, Genentech, Novartis, Sanofi (consulting fee). Mark Gudesblatt: Biogen, EMD Serono, Novartis, Sanofi Genzyme, Teva (consulting fee, research support). Carl de Moor, Stephanie Licata, Karthik Bodhinathan, Robin Avila: Biogen (employee of and owns stock and/or stock options in). Pei-Ran Ho, Lily Lee: Biogen (former employee of and may own stock and/or stock options in).

Keywords: Disease-modifying treatments in MS, Patient-reported outcomes

Author Index

A			
Aaen, Greg	1	Carlson, Aaron	28, 51, 68
Abasiyanik, Zuhail	44	Cartwright, Karen	79
Abbatemarco, Justin	80	Casper, T. Charles	1
Abu-ALHawa, Maha	55	Casserly, Courtney S.	6
Aburashed, Rany A.	39	Cederberg, Katie L.J.	74
Achiron, Anat	41	Chahin, Salim	14
Adeyemi, Ayo	18	Chapman, Katherine	58
Afsar, Salman	43	Chapman, Sandra	8
Agresta, Thomas P.	70	Checketts, Daniel	79
Aktas, Orhan	2, 21	Chen, Hailu	17, 34
Alarieh, Rola F.	38	Chen, Kun	1
Aldridge, Julie	5, 18, 23, 25, 33	Chester, Jennifer	13
Alexandri, Nektaria	50	Chinchilla, Dennis	32
Allen, Kerstin	20	Chinea, Angel	12, 45
Almen, Ruth	8	Chitnis, Tanuja	1
Alroughani, Raed	4, 39, 43	Chong, Suyinn	50
Altincatal, Arman	18, 41	Choudhry, Zia	43
Alvarenga, Victoria V.	79	Chrisant, Sarah	45, 46
Alvarez, Enrique	17, 74, 86	Christiansen, Cory	74
Amezcuca, Lilyana	31	Chung, Luke	39
Anadani, Nidhiben	13	Ciftci, Beyza	3
Anania, Veronica G.	77, 78	Cimbora, Daniel	2, 21
Antonovich, Natasha M.	53	Cinc, Edith	19
Aquillano, Lisa	32	Ciotti, John R.	14
Ararat, Kerime	56	Cofield, Stacey S.	8
Armstrong, Roisin	20, 26, 30, 40	Cohan, Stanley L.	16, 22, 24, 41
Arndt, Nancy	17	Cohen, Evan T.	71
Arnold, Douglas L.	87	Cohen, Jeffrey A.	4, 23, 29, 38, 44, 85
Arroyo, Rafael	63	Coleman, Lynette	59, 61
Aslan, Taha	40	Colhoun, Samantha R.	66
Atkins, Harold	55	Colic, Marija	55
Augustyn, Joan	60	Comi, Giancarlo	4, 25, 29, 33, 42, 43, 85
Aungst, Angela	23	Conway, Devon	3
Avila, Mirla	50	Cook, Stuart	36, 42
Avila, Robin L.	17, 20, 87	Correale, Jorge	4, 30, 85
Aydemir, Aida	30	Corsi, Gina	69
Ayer, Mavis G.	59, 66	Costello, Fiona	3
B		Costello, Kathleen	3, 57, 59
Baba, Cavid	40	Courtman, David	55
Babenko-Mould, Yolanda	63	Coyle, Patricia K.	4, 16, 25, 32, 38, 85
Bagger, Morten	16	Cree, Bruce A.C.	2, 16, 17, 19, 21, 24, 29, 31, 37, 44
Bahmanyar, Shahram	34	Crispino, Alexis	78
Bailey, Mary	67	Cross, Anne H.	4, 14, 38, 85
Baird, Jessica F.	6, 8, 9, 70	Cunningham, Rebecca	76
Bajwa, Dilpreet	82	Curko, Nina A.	55
Baker, Darren P.	39, 41	Cutter, Gary	2, 8, 15, 21, 24
Baker, Matthew	77	D	
Balabanov, Roumen	65	Dahlke, Frank	19
Balcer, Laura	65	Daizadeh, Nadia	4, 43, 63
Bamman, Marcas	9	Dalrymple, Jessica	62
Bando, Mauricio O.	56	Damian, Doris	37, 42
Banks, Amy	69	Damiri, Peter	3
Banwell, Brenda	3	Dangond, Fernando	5, 25, 28, 87
Bardsley, Belinda	19	Danon, Uri	23
Barkhof, Frederik	3	Da Silva, Talita D.	56
Barksdale, Heather	71, 72	Dastgheyb, Neda	55
Barlow, Laura	3	Davis, Bryan	62
Bar-Or, Amit	4, 16, 17, 19, 31, 37, 38, 44, 85	Dawes, Helen	72
Barron, Gerard	2, 21	Daykin, Nicola	66
Barstow, Elizabeth	11	DelMastro, Heather M.	67, 70, 72
Bass, Ann D.	4, 22, 43	DeLuca, Gabriele C.	70, 72
Bautista, Jonathan	48	DeLuca, John	7, 29
Beauvais, John	51		
Beckstead, Jason W.	73		
Benabou, Reina	56		
Benedict, Ralph H.B.	24, 37		
Bennett, Jeffrey L.	2, 21		
Benson, Leslie	1		
Berard, Jason A.	55		
Beren, Ian A.	19		
Berenbaum, Tara	3, 47, 82		
Berger, Thomas	84		
Berkovich, Regina	4, 87		
Bermel, Robert	84		
Bernitsas, Evanthia	84		
Berthele, Achim	20, 26, 30, 34, 40, 63		
Bertolotto, Antonio	63		
Berwaerts, Joris	7, 79		
Bethoux, Francois	79, 80		
Bet-Shlimon, Sargon	15		
Bhan, Virender	86		
Bisht, Babita	82		
Bleich Kimelman, Nadav	23		
Block, Allison N.	11		
Bolling, Jamie	28, 51, 68		
Bomprezzi, Roberto	56		
Bonafede, Machaon	38		
Borgert, Brooke	57		
Borowski, Jenna	22		
Boschert, Ursula	27		
Bosco, Jenna	22		
Boster, Aaron L.	4, 63		
Bowman, Mary H.	7		
Bowman, Sean	8		
Boyd, A. Suzanne	56		
Boyko, Alexey	30, 50		
Brale, Tiffany J.	74		
Brandstadter, Rachel	58		
Brandt, Alexander	47		
Bridges, Gail	24		
Broadley, Simon	43		
Brook, Richard A.	19		
Brown, Brandon	17		
Brown, Cynthia	9		
Brown, Ted/Theodore R.	80, 81		
Bruen, Denise R.	1		
Brys, Arne	47		
Buch, Eva	81		
Bueno, Ericka M.	63		
Buffels, Regine	84		
Buhse, Marijean	7, 48, 59, 67		
Bumstead, Barbara	7, 48, 67		
Burkill, Sarah	34		
Burnham, Alexander	68, 69		
Burton, Jason	86		
Butler, Oisín	84		
C			
Calabresi, Peter A.	54		
Calkwood, Jonathan	29		
Campagnolo, Denise	65		
Canzonieri, Ana	48		
Canzonieri, Ana Maria	56		
Carera, Karen	10		
Carey, Stephanie L.	73		
Carlisle, Nicola	14, 36		

DeMartino, Julie.....	27	Frenz, Ann-Kathrin.....	84	Haynes, John.....	19	
de Moor, Carl.....	87	Friedman, Coleen.....	59	Hellwig, Kerstin.....	1	
De Moraes, Ibis Ariana P.....	56	Fritz, Nora E.....	76	Hemingway, Cheryl.....	77	
Demory Beckler, Michelle.....	54	Fry, Donna.....	69	Henderson, Rochelle.....	24	
Dergalust, Sunita.....	35	Fujihara, Kazuo.....	2, 20, 21, 26, 30, 34, 40	Henke, Christian.....	19	
de Seze, Jérôme.....	4, 77, 85	Fujita, Kenji P.....	20, 26	Henry, Jaimie.....	59	
De Stefano, Nicola.....	18, 33	Fuller, Ryan.....	39, 49	Heriot, Elise.....	19	
Dias, Alice E.....	10, 48, 56, 69, 79	G			Herndon, Nicole.....	68
Dias, Marcia Baptista.....	10	Gabriele, Simone.....	66	Hersh, Carrie M.....	3, 8, 19, 86	
Diaz, Anne L.....	46	Galazka, Andrew.....	18, 23, 25, 33, 42	Hillert, Jan.....	34	
Diaz, Astrid.....	12, 45	Galetta, Steven.....	65	Hillman, Lynda R.....	58	
Dibble, Leland E.....	76	Garas, Monika.....	84	Hillner, Jennifer R.....	52	
Dietrich, Dennis.....	80	Garcia-Alonso, Belen.....	30	Ho, Pei-Ran.....	87	
Dive, Dominique.....	39	Gardener, Elizabeth.....	79	Hodges, Wayne.....	71	
Dobay, Pamela.....	86	Garmon, Eric.....	57	Hoffman, Leah.....	36	
Doniger, Glen.....	7, 53	Gawryluk, Jodie R.....	60	Hoffman, Paul M.....	71	
Donnelly, Catherine.....	52, 53	Gaythorpe, Jackie.....	52	Hoffmann, Olaf.....	86	
Dons, Benjamin.....	70	Geils, Hannah M.....	26, 27	Holl, Katsiaryna.....	84	
Dorholt, Mary.....	24	Geissbuehler, Yvonne.....	34	Holmøy, Trygve.....	84	
Doyle, Lisa.....	69	Geremakis, Caroline.....	17, 20	Hooshmand, Sam I.....	14, 36	
Drappa, Jorn.....	2, 21	Gilbertson, Peter.....	46	Hopkins, Sarah.....	13	
Drost, Taylor.....	67	Giovannoni, Gavin.....	5, 18, 19, 23, 25, 27, 30, 33, 36, 42, 67, 86	Hotermans, Christophe.....	65	
Duddy, Martin.....	61	Giri, Shivraman.....	3	Houtchens, Maria K.....	1, 60	
Due, Bryan.....	77	Glanz, Bonnie Ilene.....	60	Howard, Jennifer L.....	63	
Dunlop, Desiree.....	17	Golan, Daniel.....	7, 48, 53, 67	Hua, Le H.....	3, 8, 17, 31	
Dux, Moira.....	62	Gold, Ralf.....	19	Huang, Deren.....	22	
E			Goldenberg, Anna.....	3	Huang, Min H.....	69
Edan, Gilles.....	30	Goodman, Andrew.....	61	Hunter, Samuel F.....	39, 63	
Edwards, Erin M.....	76	Goodyear, Alexandra.....	4, 16, 23, 85	Hutchinson, Brian.....	6, 11, 68, 75	
Edwards, Thomas.....	9, 82	Gorman, Mark.....	1	Hyvert, Yann.....	18, 33	
Edwards, Zoe.....	11	Grader-Beck, Thomas.....	54	I		
Elam, Chelsea.....	44	Graves, Jennifer S.....	1, 55	Iftar, Lee S.....	55	
Erdman, Lauren.....	3	Green, Ari.....	2, 21	Ingram, Michele F.....	35	
Esfahani, Neda Zarghami.....	14	Greenberg, Benjamin M.....	58, 77	Inshasi, Jihad Said.....	28	
F			Grenningloh, Roland.....	28	Ionete, Carolina.....	63
Fabian, Michelle.....	35	Griffin, Andrea.....	35	Issard, Delphine.....	33	
Fafard, Lori.....	7, 48, 67	Griffin, Austin C.....	80	Ito, Hajime.....	77, 78	
Fakolade, Afolasade.....	46, 79	Gromisch, Elizabeth S.....	36, 51, 70, 72	J		
Farahnik, Joshua.....	82	Gudesblatt, Mark.....	7, 48, 53, 67, 87	Jack, Dominic.....	5, 25, 37, 42	
Farrell, John W., III.....	9	Guikema, Benjamin.....	40	Jackson, Kimberley.....	63	
Farsani, Maziar Eslami.....	84	Gupta, Ayan Das.....	38, 85	Jacobs, Alan.....	4	
Feasel, Corey D.....	70	Gyang, Tirisham V.....	28, 51, 57, 68	Jacobs, Dina.....	57	
Fei, Maria.....	66	H			Jacques, Francois.....	40
Feltham, Tara.....	3	Hach, Thomas.....	86	Jaenicke, Kaitlyn.....	67	
Fernandes, Dyana, Sr.....	48	Haering, Dieter A.....	4, 16, 38	Jaitly, Nina.....	44	
Fernandes, Ivone R.....	49, 65	Hakkarainen, Katja.....	34	Jasinska, Elzbieta.....	34	
Ferrero, Julianna M.....	56	Haladay, Douglas E.....	73	Jehl, Valentine.....	38, 85	
Fifolt, Matthew.....	11	Halper, June.....	3, 45, 46, 59	Johnson, Elizabeth.....	57	
Figueroa-Amaro, Joanie.....	12	Hameed, Natasha.....	67	Johnston, Karissa.....	63	
Fink, Mary Kay.....	34	Hamill, Brenda.....	59, 66	Jones, Angela.....	46	
Finlayson, Marcia.....	46, 52, 53, 79, 82	Han, May H.....	44	Jones, Catherine D.....	70	
Fitzgerald, Kathryn C.....	54	Hanssen, Andrea.....	35	Jones, Cynthia C.....	1	
Fleming, Renee.....	75	Harder, Lana.....	58	Jones, Daniel.....	23, 25	
Florio, Jordanne.....	66	Hardigan, Patrick.....	54	Jones, Noble.....	80	
Foley, John.....	87	Häring, Dieter A.....	85	K		
Fonseca, Rodrigo N.....	49	Harris, Colleen.....	52	Kaczmarek, Olivia.....	7, 48, 53, 67	
Foreman, Bo.....	76	Harris, Haleigh.....	3	Kadas, Ella M.....	47	
Fox, Edward J.....	16, 22, 23, 77	Harris, Yolanda.....	1	Kadner, Karen.....	84	
Fox, Robert J.....	15, 19, 31, 37, 38, 65	Hartigan, Clare T.....	12, 49	Kakariekka, Algirdas.....	4	
Franco, Adakson M.....	65	Hartman, Jennifer.....	80	Kalatskaya, Irina.....	27	
Fratto, Timothy.....	7, 48, 53, 67	Hartung, Hans-Peter.....	2, 21, 43, 84	Kantor, Daniel.....	7	
Freedman, Mark S.....	10, 17, 18, 30, 33, 46, 55, 79, 84	Harty, Gerard.....	37	Kapadia, Shivani.....	17, 34	
Freire-Lill, Doris.....	61	Harwood, Meagan.....	63	Kaplan, Jeffrey.....	77	
Freitas, Grazielle M.....	79	Haselkorn, Jodie K.....	10, 51, 58	Kaplan, Tyler.....	13	
French, Simon.....	53	Hauser, Stephen L.....	4, 38, 85	Kappos, Ludwig.....	4, 19, 29, 33, 37, 38, 85	

Karhonen, Pasi.....	34	Lill, Doris.....	59	McMurtrie, Melanie.....	19	
Karlsson, Goeril.....	19	Limmroth, Volker.....	4	McNamara, John.....	84	
Karpatkin, Herbert.....	71, 73	Liserio, Elaine.....	57	McNiff, Ryan.....	28, 51	
Karukote, Amputch.....	50	Liston, Dorion.....	55	Mealy, Maureen A.....	2, 21	
Katz, Amos.....	17	Liu, Hongjuan.....	29	Meca-Lallana, José.....	32	
Katz, Eliezer.....	2, 21	Liu, Jeremy.....	35	Mehta, Fal S.....	64	
Katz, Joshua D.....	26, 27	Liu, Yilan.....	8	Mehta, Rina.....	38	
Kavaklioglu, Can.....	3	Livingston, Terrie.....	31	Mehta, Tapan.....	75	
Kazmierski, Margaret.....	61	Lo, Albert C.....	70	Melo, Elisa.....	49	
Kearney, Lynda.....	59	Long, Lauren.....	39, 49	Melville, Patricia.....	59	
Keller, Birgit.....	25, 36	Longoni, Giulia.....	3, 47	Mendoza, Jason P.....	15, 17, 20	
Kera, Elizabeth.....	55	Lopez, Angel.....	12	Meng, Xiangyi.....	16, 17, 24, 31, 37, 44	
Kerloeguen, Cecile.....	4, 23, 38, 85	Lorenz, Rebecca.....	81	Merschhemke, Martin.....	4, 16, 38, 85	
Kesselman, Marc.....	54	Lotze, Timothy.....	1	Mesquita, Cristiane F.....	79	
Khatri, Bhupendra O.....	28, 32, 41, 63	Lovett-Racke, Amy.....	22	Messer, Jordan.....	17	
Khurana, Seema.....	45	Lublin, Alex L.....	28, 32	Messier, Randall S.....	67	
Khwaja, Mudeer.....	86	Lublin, Fred D.....	16, 21, 31	Miller, Aaron E.....	28	
Kielhorn, Adrian.....	63	Ludwig, Inga.....	16	Miller, Larisa.....	30, 40	
Killestein, Joep.....	84	Luhring, Sarah Kiefer.....	69	Miller, Laura K.....	74	
Kim, Ho Jin.....	2, 4, 20, 21, 26, 30, 34, 39, 40, 43	Lycke, Jan.....	4, 39	Miller, Rebecca D.....	81	
Kinkel, Revere P.....	55	M			Miller, Tamara A.....	40, 77
Klement, Riho.....	34	Mabbott, Donald.....	47	Miller, Vanessa.....	76	
Kline, Paul.....	74	Macdonell, Richard A.L.....	4, 19, 39	Miner, Annalise.....	55	
Klingelschmitt, Gaelle.....	78	Macdougall, Arlene.....	6	Misasi, Elizabeth.....	60	
Kolb, Stephen.....	15	Machado, Andre.....	80	Mischley, Laurie K.....	82	
Kolodny, Scott.....	44	Madani, Soraya.....	2, 21	Mitchell, Lisa M.....	53	
Korjagina, Marta.....	34	Maddox, Denise.....	58	Mittermayer, Caroline F.....	79	
Kou, Xiujing.....	77, 78	Maes, Micki.....	3	Mok, Koby.....	22	
Kraus, Alison.....	68	Magee, Wendy L.....	75	Mokliatchouk, Oksana.....	17	
Krehbiel, Allison.....	61	Mager, Douglas.....	24	Montague, Amanda.....	65	
Kresa-Reahl, Kiren.....	80	Mah, Jean.....	3	Montalban, Xavier.....	4, 28, 30, 85, 87	
Krieger, Stephen.....	58	Mahlanza, Tatenda.....	60	Montgomery, Scott.....	34	
Kropshofer, Harald.....	16	Mahmud, Razeen M.....	8	Moodley, Manikum.....	1	
Krug, Robert.....	72	Makin, Charles.....	17, 18, 20	Morris, David M.....	7	
Krupp, Lauren B.....	1, 29	Malone, Tiffany.....	6, 11, 68	Morrow, Sarah A.....	3, 6, 50	
Krysko, Kristen M.....	1	Maloni, Heidi W.....	59	Motl, Robert W.....	1, 6, 8, 9, 11, 70, 74, 82	
Kubala Havrdova, Eva.....	43	Manago, Mark M.....	74	Mowry, Ellen M.....	54	
Kumar, Achint.....	34, 41	Mancione, Linda.....	23	Mozzicato, Susan.....	40	
Kutz, Christen.....	20	Mandel, Matthew.....	18, 23, 87	Murphy, Kristina.....	32	
L			Manfrini, Marianna.....	84	Murphy, Sara McCurdy.....	45, 46
Lacey, Colleen.....	60	Manieri, Maria Claudia.....	60	N		
Lake, Janice.....	52	Mao-Draayer, Yang.....	22, 43, 44	Nagel, Sean.....	80	
Lancia, Samantha.....	15	Mar, Soe.....	1	Naismith, Robert T.....	17, 29	
Langdon, Dawn.....	7	Marcillat, Carole.....	77	Nakamura, Kunio.....	43	
Lathi, Ellen S.....	26, 27	Marignier, Romain.....	2, 21	Nakashima, Ichiro.....	20, 26, 30, 34, 40	
Latimer-Cheung, Amy E.....	46, 79	Mark, Victor W.....	7, 8	Narula, Sona.....	57	
Laudon-Meyer, Eva.....	34	Marom, Ehud.....	23	Navas, Carlos.....	43	
Lauwers, Nicole.....	80	Marques, Natali M.....	49	Naylor, Maria L.....	18, 41	
Lava, Christine.....	35	Marrie, Ruth Ann.....	15, 55	Naylor, Melissa.....	60	
Lava, Neil.....	32	Marriott, James.....	55	Nelles, Gereon.....	18, 41	
Lawson, Diane.....	34	Martin, Emily C.....	28, 87	Ness, Jayne.....	1	
Lazarus, K.-J.....	19	Martinez, Viviana.....	12, 45	Neto, Lindsay O.....	36, 70	
Lebson, Lori.....	5, 19	Mason, Keely.....	57	Newland, Pamela K.....	81	
Lee, Lily.....	65, 87	Massa, Marcelo.....	56	Newsome, Scott D.....	3	
Lee-Wilk, Terry.....	62	Mathison, Brianna.....	74	Nguyen, Jackie T.....	54	
LeGrand, Marie.....	50	Mayer, Lori.....	23	Nichol, Kathryn.....	79	
Leipertz, Steven L.....	51	Maynard, Michelle.....	14, 36	Nicholas, Jacqueline.....	40	
Leist, Thomas P.....	5, 23, 25, 27, 30, 42	Mayo, Chantel D.....	60	Nichols, Jamie.....	58	
Lemieux, Caroline.....	18, 25, 33	McCcoll, Mary Ann.....	52, 53	Niemeier, Janet.....	8	
Lennon-Chrimes, Sian.....	77, 78	McCreary, Morgan.....	58	Nogueira, Marcella A.M.....	79	
Leppert, David.....	4, 55	McGuigan, Christopher.....	1	Nolting, Axel.....	5, 23, 25, 42	
Levy, Michael.....	20, 26, 30, 34, 40, 63	McHugh, Brittany.....	71	O		
Lewin, James B.....	15, 22	McKay, Staci.....	7	Obeidat, Ahmed Z.....	11, 14, 36	
Li, Bingbing.....	4, 85	McKee, Keith.....	80	Oh, Jiwon.....	3, 5, 28, 41	
Li, David.....	3, 45, 46	McKenna, Odessa J.....	46, 79	Okuda, Darin T.....	48	
Licata, Stephanie.....	87	McLaughlin, Whitney.....	62	Oliveira, Barbara Gimenes.....	79	

Oliveira, Evanda Soares	10	Ramanathan, Krishnan	4, 38, 85	Sabidó, Meritxell	34
Oliver, Brant J.	64, 65	Rametta, Mark	84	Safar, Laura T.	60
Olzmann, Amy	62	Rammohan, Kottil	23, 25, 30, 36, 37, 45, 46	Sagici, Ozge	40, 44
O'Maley, Tim	54	Rano, Thomas	29	Saiz, Albert	28, 41, 77
Ona, Victor	87	Ratchford, John N.	2, 21	Salter, Amber	15
O'Neill, Jessica	66	Raulf, Friedrich	55	Salveti, Marco	18, 41
Oreja-Guevara, Celia	30, 34, 40, 86	Read, Paula	65	Sandroff, Brian M.	8, 70
Osborne, Meaghan	50	Reder, Anthony T.	17	Santos, Maria L.	49
Ozakbas, Serkan	40, 44	Rees, William	2, 21	Santos, Priscila S.	48
Ozdogar, Tuba	44	Reginald, Arun	3, 47	Santos, Rafaela L.	79
Ozog, Mark	39	Reich, Daniel S.	3	Sanwald Ducray, Patricia	77, 78
P					
Pace, Amy	26, 30, 34, 40	Rennie, Nicola	23	Sasaki, Jeffer E.	8
Pachariyanon, Pavida	50	Rensel, Mary	1	Saslow, Lori	3
Pagnotta, Patricia A.	36	Repovic, Pavle	10, 22, 34	Sayre, Lacey	6, 68
Palace, Jacqueline	20, 26, 30, 34, 40, 63	Reynolds, Piper	80	Schafer, John	6
Palomino, Miryam	55	Rhein, Juliana	48	Schick, Regina	42
Pan, Jie	50	Rhodes, Rachel	71, 73	Schipping, Sven	4
Pandey, Krupa	55	Rhys, Katherine	61	Schmidt, Hollie	65
Pardo, Gabriel	23, 25, 33	Richardson, Emma V.	1, 11	Schneider, Beth	51, 64
Parikh, Dhavan	14	Rieckmann, Peter	25	Schneiderman, Jane E.	82
Park, Julie	38	Riegler, Lindsay J.	69	Schobel, Virginia R.	43
Park, Min Su	32	Rimmer, James	75	Schorr, Emily M.	58
Paschoal, Maria A.	65	Rinker, John R.	9, 10	Schorr, Rebecca	45
Paton, Gillian R.	13	Riolo, Jon V.	29	Schreiner, Teri	1
Patton, Lisa	45, 46	Riser, Emily	75	Schuetz, Morgan L.	74
Paul, Friedemann	2, 3, 21	Riva, Sean	73	Scott, Thomas F.	45
Paul, Piper	80	Robb, Jessica F.	61	Selmaj, Krzysztof W.	4, 43, 85
Paz Soldan, M. Mateo	10	Robertson, Carly	60	Sena, Ana C.	49, 65
Pekmezi, Dorothy	75	Robertson, Derrick	17, 22, 23, 73	Sergott, Robert C.	53
Peng, Xiaomei	1	Robinson, Jennifer	43	Sharma, Alicia	62
Pereira, Giovana L.	79	Robles Cedeño, René	86	Sharrack, Basil	4
Pereira, Rafaella F.B.N.	79	Rodriguez, Lourdes	71, 73	Shaw, Jamie	28
Perez, Gina S.	13	Rodriguez, Moses	1	She, Dewi	2, 21
Perumal, Jai	65	Rodriguez, Rosangelis	71, 73	Sheffield, James K.	29
Peters, Jessica	71	Rog, David	1, 41	Shewchuk, Jason	3
Peters, Sharon	52	Rogers, Lydia	46	Shilling, Jan	80
Peterson, Daniel S.	76	Rohrbacker, Nicholas J.	19	Shinohara, Russell	3
Peterson, Patty	35	Rolfé, Alex	27	Shubin, Richard	22
Petrin, Julie	52, 53	Roman, Cortnee	20, 34	Sieber, Aryn	10
Pham, Tim	38	Romba, Meghan C.	14	Silber Baumann, Hanna	77, 78
Phillips, Amy L.	19	Romero, Rebecca	14	Silva, Barbara	79
Piani-Meier, Daniela	86	Rose, John	1	Silva, Diego	29
Piasecka-Stryczynska, Karolina	87	Rosehart, Heather	6	Silva, Roger P.	56
Pick, Anton	72	Rosenthal, Jacqueline	10	Silveira, Stephanie L.	6
Picone, Mary Ann	74, 78	Ross, Amy Perrin	84	Sima, Diana M.	47, 48
Piehl, Janet	80	Rossier, Ciara	23	Simnad, Virginia I.	81
Pille, Alicia	60	Rouyrre, Nicolas	19	Simsek, Yasemin	40
Pilutti, Lara A.	9, 46, 79, 82	Rovira, Alex	43	Singer, Barry A.	4, 18, 43, 44
Pingili, Ratnakar	4, 38, 85	Roy, Sanjeev	33, 36	Sivakolundu, Dinesh	48
Pittock, Sean J.	2, 20, 21, 26, 30, 34, 40	Royston, Minying	63	Sloan, Alicia	61
Plumb, Patricia	58	Rube, Jacob	84	Smeets, Dirk	47
Pol-Patil, Jeta	60	Rubeo, Anna	71, 73	Smith-Carrier, Tracy	63
Pontaga, Maija	50	Rubi, Cristina M.	12, 45	Smith, Kara	56
Poole, Elizabeth M.	4, 28, 32, 39, 40, 41, 63	Rubnov, Shai	23	Smith, Michael A.	2
Popper, Laura	23	Rudell, Elaine	35	Snetselaar, Linda G.	45, 82
Power, Sean	22	Ruiz, Jennifer A.	36, 67, 70, 72	Soelberg-Sorensen, Per	27, 36
Pozzilli, Carlo	39	Rusch, Carley T.	68	Sohn, Gyeongmo	50
Pulcini, Serena	66	Rusk, Jennifer L.	52	Sollero, Carlos Eduardo Vervloet ...	28, 51, 57, 68
Q					
Quarterman, Patrick	3	Russo, Kaitlin	73	Solomon, Andrew J.	10
Quezada, Tim	45	Russo, Liliana	48	Sonke, Jill	57
R					
Rai, Wijdan	15	Rutatangwa, Alice	1	Spain, Rebecca I.	10
Rajendra, Savanur Deepak	75	Rutledge, Danette	65	Sprenger, Till	28
S					
Saadatpour, Leila	14	Rypma, Bart	48	Srinivasan, Jared	7, 48, 53, 67
Sabatella, Guido	40, 63				

Stefoski, Dusan30
 Stephens, Samantha.....82
 Stephenson, Jeannie B.73
 Stevens, Sandon73
 Stillman, Jessica.....62
 Stobbe, Gary80
 Stokmaier, Daniela77
 Stoneman, Dee.....23
 Stribling, India C.26, 27
 Stuve, Olaf29
 Su, Ray.....17, 20, 22
 Su, Wendy.....16, 17, 24, 31, 37, 75
 Sugden, Joseph.....72
 Suh, Hyojin.....35
 Sullivan, Cynthia.....7, 48, 67
 Sullivan, Roseanne85
 Suzart-Woischnik, Kiliana34
 Syed, Sana25, 28, 33, 42
 Sylvester, Elke.....42
 Szafran, Eric.....57

T

Talente, Bari.....65
 Tanvir, Imran30
 Taub, Edward.....7
 Tchilibon, Susanna.....23
 Terzi, Murat20, 26, 30, 34, 40
 Tester, Nicole68
 Thakolwiboon, Smathorn50
 Thirumalai, Mohanraj.....75
 Thomas, Florian P.24, 55
 Thompson, Heidi.....50
 Thomson, Alison.....66
 Thrower, Ben10
 Tiel-Wilck, Klaus41
 Tilbery, Charles P.....49
 Tintoré, Mar4
 Titcomb, Tyler J.45, 82
 Tomic, Davorka4
 Totolyan, Natalia.....20, 26, 30, 34, 40
 Touil Allaoui, Ismahane.....55
 Traboulsee, Anthony3, 4
 Tracy, Tracy Flemming75
 Travis, Lori H.....32
 Trojano, Maria86
 Tsang, William A.55
 Tsao, Nicole18
 Turner, Aaron P.51, 70

U

Uswatte, Gitendra7

V

Vale, Vanessa D.49
 van de Ven, Kim.....3
 Van Hecke, Wim.....3, 48
 Van Liew, Charles.....76

Van Vlierberghe, Eline.....47, 48
 Van Wijmeersch, Bart.....63
 Vargas, Diana32
 Vercruyssen, Sophie.....47
 Verdun di Cantogno, Elisabetta.....30
 Verheijen, Melissa.....47
 Veri, Shelby.....56
 Verkoniemi-Ahola, Auli.....34
 Vermersch, Patrick.....4, 19, 25, 32, 36, 37, 39, 43, 86

Vernon, Karen.....59
 Verrinder, Amy.....3
 Vicente, Ivonne.....12, 45
 Vieira, Teresa Cristina.....10
 Vignos, Megan C.17, 20, 41
 Viswanathan, Shanthi.....20, 26, 30, 34, 40
 Volpenhein, Kelley69
 von Buedingen, H.-Christian77, 78
 von Geldern, Gloria14, 15, 80
 Vording, Nancy.....6
 Vucic, Steve.....41
 Vukusic, Sandra.....41

W

Wade, Peter36
 Wagner, Joanne7, 79
 Wahls, Terry L.....45, 82
 Walker, Bryan.....5, 25
 Walker, Lisa A.S.55, 82
 Walker, Tracy.....49
 Wallace, Trent.....45, 46
 Wallin, Mitchell T.3, 10, 53
 Walsh, Karen.....65
 Walters, Arthur S.....74
 Waltz, Michael1
 Wan, Michael J.47
 Wang, Kai-Chen20, 26, 30, 34, 40
 Wang, Liangwei.....2
 Wang, Yujie.....54
 Wang, Zhini.....80
 Ward, Jordan6
 Waslo, Carin S.10
 Waubant, Emmanuelle1
 Weber, Kristina77
 Weber, Martin S.....28, 87
 Weckbecker, Gisbert.....55
 Wei, Olivia.....36
 Weigel, Megan75
 Weinshenker, Brian G.2, 21, 77
 Weinstock-Guttman, Bianca1
 Weiss, Jamie L.....24
 Weiss, Michael S.22
 Wells, Christopher.....12
 Werner, Kristine.....10
 Werner, Simone.....84
 West, Kathryn.....48
 West, Timothy20

Wicklein, Eva-Maria84
 Wiendl, Heinz.....4, 16, 39, 85
 Wilken, Jeffrey7, 48, 53, 67
 Wilkinson, Carmel.....59, 61
 Willi, Roman4, 85
 Williams, Justin D.....61
 Williams, Mitzi J.....31
 Williamson, Eric35
 Willingham, Emily.....65
 Wilson, Kathleen.....38
 Wilson, Whitney.....57
 Wingerchuk, Dean2, 20, 21, 26, 30, 34, 40
 Wingerden, Janneke41
 Wisdom, Peggy J.13
 Witek, Stephanie.....39
 Wolinsky, Jerry S.3, 87
 Womble, Brent M.7
 Wong, Schiffo L.....37
 Wray, Sibyl.....18, 22, 34, 40, 41, 43
 Wu, Naomi35
 Wuerfel, Jens28
 Wundes, Annette.....14, 15, 17, 80

X

Xiong, Kuangnan65

Y

Yadav, Sunil K.47
 Yale, Kendra80
 Yamamura, Takashi77
 Yamout, Bassem30
 Yang, Ling.....29
 Yeh, E. Ann3, 47, 82
 Yeo, Sandy.....84
 Yigit, Pinar.....40, 44
 Young, Angela59
 Young, Hui-Ju75
 Yountz, Marcus20, 26, 30, 34, 40
 Yu, Winnie.....73

Z

Zabeti, Aram81
 Zachariah, Jikku.....45
 Zaffaroni, Mauro.....41
 Zalesak, Martin.....23
 Zare, Ameneh.....81
 Zarif, Myassar.....7, 48, 67
 Zhao, Enxu77
 Zheng, Chao45
 Zhou, Jia31
 Ziehn, Marina.....23, 44
 Ziemssen, Tjalf.....4, 86
 Zimmerman, Vanessa57
 Zivadinov, Robert.....29
 Zuppichini, Mark D.48

MS EMPOWERMENT SERIES

SHARED DECISION MAKING IN MULTIPLE SCLEROSIS

MS EMPOWERMENT SERIES
SHARED DECISION MAKING IN MULTIPLE SCLEROSIS



YOU ARE NOT ALONE
*Resources and Coping Tips
Care Partners of People
With MS*

 An educational series for people with multiple sclerosis (MS) developed in conjunction with the International Organization of MS Nurses (IOMSN).

Provided through an educational grant from Celgene.

MS EMPOWERMENT SERIES
SHARED DECISION MAKING IN MULTIPLE SCLEROSIS




FATIGUE IN MULTIPLE SCLEROSIS
*What Causes It?
How Do I Manage It?*


 An educational series for people with multiple sclerosis (MS) developed in conjunction with the International Organization of MS Nurses (IOMSN).

Provided through educational grants from Biogen, Celgene, and Sanofi Genzyme.

MS EMPOWERMENT SERIES
SHARED DECISION MAKING IN MULTIPLE SCLEROSIS




SEXUALITY AND MULTIPLE SCLEROSIS
*How to Overcome Emotional
Physical Barriers to Intimacy
Where to Find Help*


 An educational series for people with multiple sclerosis (MS) developed in conjunction with the International Organization of MS Nurses (IOMSN).

Provided through educational grants from Biogen, Celgene, and Sanofi Genzyme.

MS EMPOWERMENT SERIES
SHARED DECISION MAKING IN MULTIPLE SCLEROSIS



SLEEP MATTERS
*How Multiple Sclerosis
Can Affect Sleep*

 An educational series for people with multiple sclerosis (MS) developed in conjunction with the International Organization of MS Nurses (IOMSN).

Provided through educational grants from Biogen, Celgene, and Sanofi Genzyme.

Download at:

<http://iomsn.org/publications>