



Short report

Police interference with methadone treatment in Bangkok, Thailand



Kanna Hayashi ^{a,b,*}, Lianping Ti ^{a,c}, Karyn Kaplan ^d, Paisan Suwannawong ^d,
Evan Wood ^{a,b}, Thomas Kerr ^{a,b}

^a British Columbia Centre for Excellence in HIV/AIDS, 608 – 1081 Burrard Street, Vancouver, BC V6Z 1Y6, Canada

^b Faculty of Medicine, University of British Columbia, 317-2194 Health Sciences Mall, Vancouver, BC V6T 1Z3, Canada

^c School of Population and Public Health, University of British Columbia, 2206 East Mall, Vancouver, BC V6T 1Z3, Canada

^d Thai AIDS Treatment Action Group, 18/89 Vipawadee Rd., soi 40 Chatuchak, Bangkok 10900, Thailand

Introduction

Methadone maintenance therapy is an evidence-based pharmacotherapy to treat opioid dependence (World Health Organization [WHO], United Nations Office on Drugs and Crime, & Joint United Nations Programme on HIV/AIDS, 2012). The WHO and other United Nations agencies recommend it as part of essential services for the prevention, treatment and care of human immunodeficiency virus (HIV) infection among people who inject drugs (IDU) as it has been demonstrated to reduce HIV risk behaviour and improve access and adherence to HIV treatment among this population (WHO et al., 2012). Despite the proven therapeutic benefits, methadone is not always readily accessible to those in need in many settings (Wolfe, Carrieri, & Shepard, 2010).

In Thailand, a country contending with a longstanding dual epidemic of opioid use and HIV among IDU (Assanangkornchai et al., 2008; National AIDS Prevention and Alleviation Committee [NAPAC], 2010; Reid & Costigan, 2002), methadone was approved for opioid substitution therapy in 2000, although having been provided as tapered detoxification regimens rather than as a maintenance therapy (Tyndall, 2011). In 2008, methadone was added to the universal healthcare coverage scheme (NAPAC, 2010). In Bangkok, methadone treatment is primarily provided by the Bangkok Metropolitan Authority (BMA) through its 17 public health centers, two hospitals and one stand-alone clinic (Tyndall, 2011).

The Thai government has for many years implemented aggressive drug prohibition approaches (Hayashi, Small, Csete, Hattirat, & Kerr, 2013a). A recent qualitative study of IDU in Bangkok has highlighted that the police often conduct surveillance of people accessing methadone clinics; harass methadone patients with extortion and urine drug testing; and arrest patients who test positive for an illicit drug (Hayashi et al., 2013a). In response, some

interviewees reported feeling discouraged from continuing their methadone treatment (Hayashi et al., 2013a). While these reports are consistent with international literature indicating that aggressive policing can impede IDU's access to healthcare (Kerr, Small, & Wood, 2005), few studies have quantitatively evaluated the impact of policing around health service sites. Therefore, we sought to examine the prevalence and correlates of noticing police presence around methadone clinics among methadone-treated IDU in Bangkok.

Methods

Data were derived from the Mitsampan Community Research Project (MSCRP), a collaborative research effort involving the Mitsampan Harm Reduction Center (MSHRC; a drug user-run drop-in centre in Bangkok, Thailand), Thai AIDS Treatment Action Group (Bangkok, Thailand), Chulalongkorn University (Bangkok, Thailand), and the British Columbia Centre for Excellence in HIV/AIDS/University of British Columbia (Vancouver, Canada). This serial cross-sectional study aims to investigate drug-using behaviour, healthcare access, and other drug-related harms among IDU in Bangkok. Between July and October 2011, 440 IDU in Bangkok were surveyed. Potential participants were recruited through peer outreach efforts and word-of-mouth, and were invited to attend the MSHRC or O-Zone House (another drop-in centre in Bangkok) to enrol in the study. Recruitment criteria included adults residing in Bangkok or in adjacent provinces who had injected drug(s) in the past six months. The recruitment methods and the sample characteristics have been described in detail previously (Hayashi et al., 2012). All participants provided informed consent and completed an interviewer-administered questionnaire eliciting a range of information, including demographic characteristics, drug use patterns, and experiences with drug law enforcement and accessing healthcare. Upon completion of the questionnaire, participants received a stipend of 350 Thai Baht (approximately US\$12). The study was approved by the research ethics boards at Chulalongkorn University and the University of British Columbia.

* Corresponding author at: BC Centre for Excellence in HIV/AIDS, 608 – 1081 Burrard Street, Vancouver, BC V6Z 1Y6, Canada. Tel.: +1 604 682 2344x63210; fax: +1 604 806 9044.

E-mail addresses: khayashi@cfenet.ubc.ca, kanna.hayashi@gmail.com (K. Hayashi).

Participants who reported having received methadone treatment during the previous six months and who had complete data were included in this study. The primary outcome of interest was noticing police presence near methadone clinics during the previous six months, defined as answering "Yes" to a question: "In the last six months, have you noticed police presence where you obtain methadone?" Explanatory variables included: younger age; male gender; daily heroin injection; daily midazolam (a short-acting benzodiazepine) injection; daily methamphetamine injection; binge drug use; rushed injection; syringe sharing; HIV serostatus (positive vs. negative or unknown); non-fatal overdose; and avoiding healthcare. Drug-using behaviours referred to the previous six months. Consistent with our previous MSCR study (Kerr et al., 2014), avoiding healthcare was defined as answering "Yes" to a question: "Do you sometimes avoid healthcare because you are a drug user?"

To examine bivariate associations between each explanatory variable and the outcome, we used the Pearson χ^2 test. Fisher's exact test was used when one or more of the cells contained values less than or equal to five. Next, we fit a multivariate logistic regression model including all variables that were significantly associated with the outcome at the $p < 0.05$ level in bivariate analyses. All p -values were two-sided.

In a sub-analysis, we also asked participants if they have been harassed by police after receiving methadone, if they had stopped methadone treatment anytime during the past six months, and the reasons for stopping treatment (participants could provide more than one response). We used the Pearson χ^2 test to determine if there was an association between methadone discontinuation and noticing police presence near methadone clinics.

Results

In total, 190 IDU who accessed methadone treatment during the previous six months participated in this study, including 29 (15.3%) women. The median age was 38 years (interquartile range: 34–48). In total, 109 (57.4%) individuals reported having noticed police presence near methadone clinics in the previous six months. Of these, 42 (38.5%) reported having been harassed by police after receiving methadone.

Table 1 shows the results of bivariate and multivariate logistic regression analysis. As shown, in the multivariate analysis, factors independently and positively associated with the outcome included daily midazolam injection (adjusted odds ratio [AOR]: 2.33; 95% confidence interval [CI]: 1.26–4.32) and avoiding healthcare (AOR: 2.20; 95% CI: 1.02–4.74).

In the sub-analysis, 39 (20.5%) of the entire sample reported having stopped methadone treatment during the previous six months. There was no significant difference in the proportion of methadone discontinuation between those who did and did not notice police presence near methadone clinics ($p = 0.62$). Among 190 individuals who noticed police presence, 21 (19.3%) reported methadone discontinuation, and the primary reasons for stopping treatment were: incarceration (33.3%); difficulty accessing clinics (19.0%); and did not want to take methadone any more (14.3%). In contrast, 18 (22.2%) of 81 participants who did not notice police presence reported methadone discontinuation, and the primary reasons included: did not want to take methadone any more (42.1%); incarceration (22.2%); and too many side effects (11.1%).

Discussion

We found that more than half of a sample of methadone-treated IDU in Bangkok noticed police presence near methadone clinics in the previous six months. Patients who injected midazolam daily

and avoided accessing healthcare were more likely to have witnessed police officers near clinics. Approximately 40% of those who had noticed police presence also experienced police harassment and 20% stopped methadone treatment primarily due to incarceration and difficulty accessing clinics.

While rates of methadone discontinuation did not differ between those who did and did not notice police presence, the primary reasons for stopping methadone appeared to differ between the two groups. Specifically, difficulty-accessing clinics was referred to only among those who noticed police presence. Further, incarceration appeared to be more commonly reported among this group, which may suggest that patients who saw police officers near clinics may have been subsequently arrested by these police officers, detained, incarcerated and taken off methadone. These findings corroborate previous qualitative studies indicating that Thai police often conduct targeted surveillance and arrests of drug offenders around methadone clinics in Bangkok (Hayashi et al., 2013a). It has also been reported that drug policing in this setting generally involves various forms of police misconduct and violence (Hayashi et al., 2013a). Given the aggressiveness of policing practices in this setting, it may be that policing activities around methadone clinics have created a climate of fear among methadone patients and deterred some patients from accessing treatment. Such impacts of policing have also been indicated in the international literature (Meng & Burris, 2013).

The independent association between daily midazolam injection and noticing police presence near clinics may indicate a potential consequence of compromised access to methadone treatment. A previous study in this setting has shown that the majority of midazolam injectors use midazolam as a legal substitute for heroin due to the low price and its potential to alleviate heroin withdrawal symptoms (Hayashi et al., 2013b). In light of these reports, our findings may suggest that methadone patients who noticed police presence may have skipped some methadone doses to avoid a future encounter with police near clinics, and they may have resorted to injecting midazolam to diminish the symptoms of opioid withdrawal. High rates of midazolam misuse are concerning as there is no proven pharmacotherapy to treat benzodiazepine dependence (WHO-SEARO, 2008), and frequent midazolam injection is linked to serious venous disease and injuries (Hayashi et al., 2013b). An alternative explanation for our findings is that midazolam injectors are more likely to generate police attention. Future research should seek to unpack the association between policing and midazolam injecting among Thai IDU accessing methadone treatment.

We also found that IDU who reported avoiding healthcare were more likely to report witnessing police officers near clinics. In Bangkok, methadone clinics are typically housed within the BMA public health centers and hospitals that provide general healthcare services (Tyndall, 2011). Therefore, it may be that policing around methadone clinics may not only interfere with methadone treatment but also impede methadone patients' access to healthcare more generally. Alternatively, individuals who actively avoid healthcare due to their drug use may be more cautious about police presence.

Collectively, these findings raise concern that policing around methadone clinics may be undermining the potential benefits of including methadone treatment in the universal healthcare scheme. Efforts to ensure that policing practices do not interfere with the essential health service for IDU are urgently needed in this setting. A recent report has indicated that harm reduction services, including methadone treatment, are poorly understood by the police and are not part of the police curriculum in Thailand, although a plan for a pilot harm reduction training program for police officers is underway (Macdonald & Nacapew, 2013). There is a clear need for the prompt implementation,

Table 1

Bivariate and multivariate logistic regression analysis of factors associated with noticing police presence near methadone clinics in the previous six months among methadone-treated IDU in Bangkok, Thailand ($n = 190$).

Characteristic	Odds ratio (OR)	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
<i>Demographic</i>			
Younger age (<38 years vs. ≥38 years)	0.78 (0.44–1.38)		
Gender (Male vs. female)	0.79 (0.35–1.79)		
<i>Drug use behaviour</i>			
Heroin injection* (Daily vs. < daily)	2.31 (0.87–6.15)		
Midazolam injection* (Daily vs. < daily)	2.15 (1.19–3.89)		2.33 (1.26–4.32)
Methamphetamine injection* (Daily vs. < daily)	1.25 (0.23–8.28)		
Binge drug use* (Yes vs. no)	1.49 (0.81–2.75)		
Rushed injection* (Yes vs. no)	2.08 (1.16–3.75)		1.75 (0.94–3.25)
Syringe sharing* (Yes vs. no)	1.50 (0.73–3.10)		
<i>Health problems and healthcare access</i>			
HIV serostatus (Positive vs. negative or unknown)	2.10 (0.97–4.53)		
Non-fatal overdose* (Yes vs. No)	2.26 (0.18–120.32)		
Avoiding healthcare (Yes vs. No)	2.27 (1.11–4.67)		2.20 (1.02–4.74)

IDU, people who inject drugs; CI, confidence interval.

* Activities in the previous six months.

evaluation and scale-up of such a training program. However, international literature indicates that effective harmonization of police and public health activities has proven challenging, and therefore more work needs to be done in this area (Meng & Burris, 2013).

This study has several limitations. First, we cannot infer causation from this observational study. Further, the cross-sectional study design did not allow us to assess temporal relationships between the outcome and independent variables. Second, the self-reported data may have been affected by some reporting biases, including socially desirable responding and recall bias. However, we note that this type of data has been commonly utilized in observational studies involving IDU and has been found to be valid (Darke, 1998). Third, as the study sample was not randomly selected, generalizability of our findings may be limited. Lastly, we recognize that our dependent variable does not directly measure intensive policing around methadone sites. However, given that targeted policing around these sites has been well documented (Hayashi et al., 2013a), and considering the high prevalence (40%) of experiencing police harassment after receiving methadone, we feel that the variable adequately functions as a proxy measure of intensive policing in this particular study. That said, future research should seek to refine the type of measure of police influence.

In summary, we found that more than half of methadone-treated IDU in Bangkok noticed police presence near methadone clinics in the previous six months. Daily midazolam injectors and individuals who reported avoiding healthcare were more likely to have witnessed police officers around methadone clinics. Approximately one-fifth of those who had noticed police presence stopped methadone treatment, most commonly due to incarceration and difficulty accessing clinics. These findings raise concern that intensive policing in the vicinity of methadone clinics may be interfering with methadone treatment in this setting. Efforts to align law enforcement and public health activities are urgently needed.

Acknowledgements

We would particularly like to thank the staff and volunteers at the Mitsampan Harm Reduction Center, Thai AIDS Treatment Action Group and O-Zone House for their support and Dr. Niyada Kiatying-Angsulee of the Social Research Institute, Chulalongkorn University, for her assistance with developing this project. We also thank Tricia Collingham, Deborah Graham, Caitlin Johnston, Calvin Lai and Peter Vann for their research and administrative assistance, and Prempreeda Pramoj Na Ayutthaya, Arphatsaporn Chaimongkon and Sattara Hattirat for their assistance with data collection. The study was supported by Michael Smith Foundation for Health Research. This research was also undertaken, in part, thanks to funding from the Canada Research Chairs program through a Tier 1 Canada Research Chair in Inner City Medicine which supports E.W.

Conflicts of interest

None declared.

References

- Assanangkornchai, S., Aramrattana, A., Perngparn, U., Kanato, M., Kanika, N., & Na Ayudhya, A. S. (2008). Current situation of substance-related problems in Thailand. *Journal of the Psychiatric Association of Thailand*, 53(Suppl. 1), 24S–36S.
- Darke, S. (1998). Self-report among injecting drug users: A review. *Drug and Alcohol Dependence*, 51(3), 253–263, discussion 267–8.
- Hayashi, K., Fairbairn, N., Suwannawong, P., Kaplan, K., Wood, E., & Kerr, T. (2012). Collective empowerment while creating knowledge: A description of a community-based participatory research project with drug users in Bangkok, Thailand. *Substance Use & Misuse*, 47(5), 502–510.
- Hayashi, K., Small, W., Csete, J., Hattirat, S., & Kerr, T. (2013). Experiences with policing among people who inject drugs in Bangkok, Thailand: A qualitative study. *PLOS Medicine*, 10(12), e1001570.
- Hayashi, K., Suwannawong, P., Ti, L., Kaplan, K., Wood, E., & Kerr, T. (2013). High rates of midazolam injection and associated harms in Bangkok, Thailand. *Addiction*, 108(5), 944–952.
- Kerr, T., Hayashi, K., Ti, L., Kaplan, K., Suwannawong, P., & Wood, E. (2014). The impact of compulsory drug detention exposure on the avoidance of healthcare

- among injection drug users in Thailand. *International Journal of Drug Policy*, 25(1), 171–174.
- Kerr, T., Small, W., & Wood, E. (2005). The public health and social impacts of drug market enforcement: A review of the evidence. *International Journal of Drug Policy*, 16(4), 210–220.
- Macdonald, V., & Nacapew, S. (2013, November 18). *Drug control and harm reduction in Thailand*. London: International Drug Policy Consortium. Retrieved from <http://idpc.net/publications/2013/11/idpc-briefing-paper-drug-control-and-harm-reduction-in-thailand> (31.03.14)
- Meng, J., & Burris, S. (2013). The role of the Chinese police in methadone maintenance therapy: A literature review. *International Journal of Drug Policy*, 24(6), e25–e34.
- National AIDS Prevention and Alleviation Committee. (2010). *UNGASS country progress report Thailand: Reporting period: January 2008–December 2009*. Retrieved from http://data.unaids.org/pub/Report/2010/thailand_2010-country_progress_report_en.pdf (31.03.14)
- Reid, G., & Costigan, G. (2002). *Revisiting the hidden epidemic: A situation assessment of drug use in Asia in the context of HIV/AIDS*. Fairfield: The Center for Harm Reduction, The Burnet Institute.
- Tyndall, M. (2011). *Harm reduction policies and interventions for injection drug users in Thailand*. Bangkok: World Bank.
- World Health Organization, United Nations Office on Drugs and Crime, & Joint United Nations Programme on HIV/AIDS. (2012). *WHO, UNODC, UNAIDS technical guide for countries to set targets for universal access to HIV prevention, treatment and care for injecting drug users: 2012 revision. World Bank East Asia and Pacific Economic Update 2010*. Geneva: World Health Organization.
- WHO-SEARO. (2008). *Operational guidelines for the management of opioid dependence in the South-East Asia region*. New Delhi: World Health Organization, Regional Office for South-East Asia (WHO-SEARO).
- Wolfe, D., Carrieri, M. P., & Shepard, D. (2010). Treatment and care for injecting drug users with HIV infection: A review of barriers and ways forward. *Lancet*, 376(9738), 355–366.