

## Knee ACL Reconstruction with Meniscus Repair Rehabilitation Protocol

By James L Chen, MD

Phase	Time	Exercises/Activities/RESTRICTIONS	Goals
Early Post operation	0-2 weeks	<b>Brace at all times</b> <b>Crutches for 4 weeks</b> <b>Non Weight Bearing for 4 weeks</b> Passive/Active ROM Straight leg raise Quadriceps isometrics Patella mobilizations	Pain and swelling control
Phase I	2-6 weeks	<b>Brace/crutches</b> <b>Crutches for 4 weeks</b> <b>Non Weight Bearing for 4 weeks</b> Patella mobilizations Active/passive/AAROM Straight leg raise Wall slides/leg press (0-80°) Quad isometrics Hip/CORE/ankle strengthening Step-up program Proprioception program Modalities OK Home exercise program	ROM to 120° Healing of ACL Quadriceps control Good patella mobility
Phase II	6-12 wks	AAROM exercises Leg press—0-90° arc Step-up/Step-down Lunges Squats Proprioception/balance program Hip/CORE program Straight leg raise Stair master/elliptical Modalities OK Home exercise program	ROM to normal Normal patellar mobility Restore normal gait Ascent stairs with good control Return to normal ADL
Phase III	12-20 wks	Progress squat program Advance quad/hamstring program Proprioception training Agility exercises Elliptical Lower extremity stretching Modalities Home exercise program	Descend stairs with good control Improve flexibility  Running at 4 months
Phase IV	>20 wks	Continue LE strengthening/flexibility Forward running (per MD) Brace for sports (per MD)	Sport specific activity Return to competitive activities at 6-8 months

## **Post Operative Instructions**

### **ON THE DAY OF SURGERY:**

**Ice:** Use ice or the cold therapy unit over the leg for 15 minutes every 1 to 2 hours to decrease pain and inflammation.

**Pain medication:** You will be given a prescription for a narcotic for pain control. Begin taking the pain medication as prescribed as soon as you have pain. Do not wait until you are in severe pain before taking them as it takes 30 to 45 minutes to work. You should be able to stop taking the pain medicine within two to three weeks. Avoid taking Tylenol while you are taking the narcotic as they both contain acetaminophen which can be dangerous to your liver in high amounts.

**Eating:** Begin eating with liquids and light foods (jello, soup, bread). Progress to your normal diet slowly over the 24 hours following surgery. The narcotics can make you nauseous, so take them with food whenever possible.

### **DAY 1 AND UNTIL POST-OP VISIT:**

**Dressings:** Keep the dressing on the surgical site until your first post-operative visit— usually 5-7 days after your surgery. This decreases the risk of post-operative infection. Do not put ointment on your incisions.

**Bathing:** You may sponge bathe following the surgery. You can also shower with a plastic garbage bag around the leg, but you must keep the dressing absolutely dry. Do not bathe in a tub. Once the sutures are removed you may shower or bathe and get the incision wet. Do not swim in a pool or the ocean until instructed by your surgeon.

**Crutches:** Continue to use the crutches for approximately 4 weeks or until instructed by your doctor.

**Driving:** Do not drive until instructed to do so.

**Therapy:** Contact a physical therapist prior to surgery to begin formal physical therapy approximately 5-10 days following your surgery.

It is normal to experience some swelling in the knee and leg. To decrease this, keep your leg elevated above the level of your heart as much as possible.

Avoid putting a pillow behind your knee as this can cause problems getting your knee straight.


**Please call the office (415) 900-3000 if you have questions or any of the following:**

- Fever greater than 101.5 or 38.5 degrees
- Continuing drainage from the wound
- Unrelenting pain
- Excessive nausea/vomiting due to pain medication
- ANY other worrisome condition


**Post Operative Instructions: Knee Surgery**

**EXERCISES: START POST-OPERATIVE DAY 1:**


1. **Straight Leg Raises.** Lift your leg about twenty-four (24) inches off the ground and hold it for 10 seconds. Do a minimum of three (3) sets of ten (10) repetitions. Make sure you do this with your brace locked in full extension.




2. **Quad Sets.** With this exercise, you tighten your thigh muscles and hold for five (5) seconds. Do a minimum of three (3) sets of ten (10) repetitions. When you tighten your thigh muscles, it will feel like your knee is being pushed into the ground. Make sure you do this with your brace locked in full extension.



3. **Move your kneecap (patella) side to side.** Move the kneecap toward the inner side of your leg and hold it for five (5) seconds. Do three (3) sets of ten (10) repetitions in each direction. This will be harder to do until the bandages are removed.




4. **Heel Slides.** (bending the knee) slide your heel toward your buttock. This may be assisted by using a towel to pull your foot.



5. **Bridging Exercises.** Roll a towel and put it under your heel with nothing under your knee. Keep this position for 5 to 10 minutes. You can assist this by resting your hand on your thigh, near the knee.



6. **Calf Pumps.** Move both ankles up and down, at least 10 times an hour until you are up and around regularly to encourage blood flow in the lower legs and reduce the risk of DVT (blood clot).



7. **CPM.** A CPM (continuous passive motion) machine may be ordered by your surgeon. This machine helps gain your motion a little faster than if you did not have it, and may help with stiffness. Use this machine at least six (6) hours a day (you may want to do this at night)- if so, you will probably need sleeping pills such a Benadryl to help you for the first few nights. Start at zero (0) to thirty (30) degrees, and increase it five (5) degrees every twelve (12) hours as tolerated. You may return the machine when your knee bends more than ninety (90) degrees with the therapist or the machine is at its full range zero (0) to one hundred twenty (120) degrees for two (2) to three (3) days. Unlock your brace to use the CPM (**do not remove the brace**).