

LIMIT COMPUTABILITY AND ULTRAFILTERS

URI ANDREWS, MINGZHONG CAI, DAVID DIAMONDSTONE, NOAH SCHWEBER

ABSTRACT. We study a class of operators on Turing degrees arising naturally from ultrafilters. Suppose \mathcal{U} is a nonprincipal ultrafilter on ω . We can then view a sequence of sets $A = (A_i)_{i \in \omega}$ as an “approximation” of a set B produced by amalgamating the A_i via \mathcal{U} : we set $\lim_{\mathcal{U}}(A) = \{x : \{i : x \in A_i\} \in \mathcal{U}\}$. This can be extended to the Turing degrees, by defining $\delta_{\mathcal{U}}(\mathbf{a}) = \{\lim_{\mathcal{U}}(A) : A = (A_i)_{i \in \omega} \in \mathbf{a}\}$. The $\delta_{\mathcal{U}}$ — which we call “ultrafilter jumps” — resemble classical limit computability in certain ways. In particular, $\delta_{\mathcal{U}}(\mathbf{a})$ is always a Turing ideal containing $\Delta_2^0(\mathbf{a})$. However, they are also closely tied to Scott sets: $\delta_{\mathcal{U}}(\mathbf{a})$ is always a Scott set containing \mathbf{a}' . (This yields an alternate proof of the standard result in reverse mathematics that Weak Konig’s Lemma is strictly weaker than arithmetic comprehension.)

Our main result is that the converse also holds: if \mathcal{S} is a countable Scott set containing \mathbf{a}' , then there is some ultrafilter \mathcal{U} with $\delta_{\mathcal{U}}(\mathbf{a}) = \mathcal{S}$. We then turn to the problem of controlling the action of an ultrafilter jump $\delta_{\mathcal{U}}$ on two degrees simultaneously, and for example show that there are nontrivial degrees which are “low” for some ultrafilter jump. Finally, we study the structure on the set of ultrafilters arising from the construction $\mathcal{U} \mapsto \delta_{\mathcal{U}}$; in particular, we introduce a natural preordering on this set and show that it is connected with the classical Rudin-Keisler ordering of ultrafilters. We end by presenting two directions for further research.

CONTENTS

1. Introduction	1
2. Basic Properties of $\delta_{\mathcal{U}}$	4
3. Building Scott sets	6
4. Lowness notions	8
5. Comparing ultrafilters	12
6. Further directions	14
6.1. Filter jumps	14
6.2. Ultrafilter jumps of Turing ideals	15
References	16

1. INTRODUCTION

If X is a set of natural numbers, we can view X as a countable array of sets in a natural way:

Definition 1.1. *Let $X \subseteq \omega$. Then we let $X_i = \{j : \langle i, j \rangle \in X\}$ and $X^j = \{i : \langle i, j \rangle \in X\}$ be the i th column and j th row of X , respectively.*

We can then consider the eventual behavior of each row of X . In particular, in case every row of X is finite or cofinite — that is, if $\lim_s(X^j(s))$ exists for every j — then we can define the *limit* of X as

$$\lim(X) = \{j : \lim_s(X^j(s)) = 1\} = \{j : X^j \text{ cofinite}\}.$$

If $Z = \lim(Y)$ and $Y \leq_T X$, we say Z is *limit computable* relative to X .

If Z is limit computable relative to X then $Z \leq_T X'$. Conversely, Shoenfield's limit lemma shows that this is exact: if $A \leq_T X'$, then $A = \lim(Y)$ for some $Y \equiv_T X$. While this is only one of many characterizations of the jump, limit computability is of particular interest because it suggests a wide class of generalizations: given any notion of "generalized limit," we can consider the collection of sets which are *generalized limit computable* relative to a given X . These in turn yield generalized jump operators, taking a set X to the collection of sets which are generalized limit computable relative to X .

In this paper, we investigate limit computability along (nonprincipal) ultrafilters. For each ultrafilter \mathcal{U} , we introduce a function $\delta_{\mathcal{U}}$ taking each Turing degree \mathbf{a} to the collection of sets " \mathcal{U} -limit computable" in members of \mathbf{a} . Besides establishing its basic properties, we characterize the possible values of $\delta_{\mathcal{U}}(\mathbf{a})$, define a notion of "lowness for ultrafilters" and study the question of characterizing these degrees, and examine the ordering on ultrafilters induced by the construction $\mathcal{U} \mapsto \delta_{\mathcal{U}}$.

We recall the definition of an ultrafilter:

Definition 1.2. A collection of sets $\mathcal{A} \subseteq \mathcal{P}(\omega)$ is an *ultrafilter* if it satisfies the following properties:

- (1) $\omega \in \mathcal{A}$, $\emptyset \notin \mathcal{A}$.
- (2) If $X \in \mathcal{A}$ and $X \subseteq Y \subseteq \omega$, then $Y \in \mathcal{A}$.
- (3) If $X, Y \in \mathcal{A}$, then $X \cap Y \in \mathcal{A}$.
- (4) For every $X \subseteq \omega$, $X \in \mathcal{A}$ or $(\omega - X) \in \mathcal{A}$.
- (5) \mathcal{A} contains no finite set.

Although the existence of ultrafilters is not provable in ZF alone, it follows from the axiom of choice that there are $2^{2^{\aleph_0}}$ -many ultrafilters on ω , the maximum number possible.

Remark 1.3. The final axiom in our list is usually not included; ultrafilters satisfying it are called *nonprincipal* ultrafilters. In this paper we shall only be interested in nonprincipal ultrafilters, so for brevity we include it here. Additionally, we will only ever consider ultrafilters on ω .

Using the fourth ultrafilter axiom, we can take the limit along any ultrafilter of any sequence $(X_i)_{i \in \omega}$ of sets, and by 1.2(5) this notion of limit agrees with the classical one when each X^j is finite or cofinite. Taking limits along an ultrafilter then yields the notion of limit computability along an ultrafilter, which in turn yields a class of operators on Turing degrees.

Formally, we proceed as follows. We begin by defining the limit along an ultrafilter of an array of sets:

Definition 1.4. For a sequence of sets $X = (X_i)_{i \in \omega}$ and an ultrafilter \mathcal{U} , we let

$$\lim_{\mathcal{U}}((X_i)_{i \in \omega}) = \{j : \{i : j \in X_i\} \in \mathcal{U}\} = \{j : X^j \in \mathcal{U}\}.$$

Note that, as in the case of classical limit computability, each column X_i functions as an approximation to the limit set $\lim_{\mathcal{U}}(X)$, and dually each row X^j determines the j th bit of $\lim_{\mathcal{U}}(X)$.

We can now define the maps, $\delta_{\mathcal{U}}$:

Definition 1.5. Fix an ultrafilter \mathcal{U} . For \mathbf{a} a degree, we let

$$\delta_{\mathcal{U}}(\mathbf{a}) = \{\lim_{\mathcal{U}}((X_i)_{i \in \omega}) : (X_i)_{i \in \omega} = X \in \mathbf{a}\}.$$

It is the maps $\delta_{\mathcal{U}}$ and their images, especially $\delta_{\mathcal{U}}(\mathbf{0})$, which are the subject of this article. We call maps of the form $\delta_{\mathcal{U}}$ *ultrafilter jumps*.

We begin by establishing basic closure properties of sets of the form $\delta_{\mathcal{U}}(\mathbf{a})$; this culminates in the following characterization, which is our main result. Recall that a *Scott set* is a collection of subsets of ω closed under Turing reducibility and join, and which contains an element encoding an infinite tree $T \subseteq 2^{<\omega}$ only if it also contains an infinite path through T .

Theorem 1.6. *For \mathbf{a} a Turing degree, the following are equivalent:*

- \mathfrak{S} is a countable Scott set containing \mathbf{a}' .
- There is some ultrafilter \mathcal{U} such that $\delta_{\mathcal{U}}(\mathbf{a}) = \mathfrak{S}$.

Next, we look at how a single ultrafilter jump can behave with respect to different degrees. We call a degree \mathbf{a} *u-low* if there is some ultrafilter \mathcal{U} such that $\delta_{\mathcal{U}}(\mathbf{a}) = \delta_{\mathcal{U}}(\mathbf{0})$. Using techniques similar to those in the proof of the main theorem, we show the following:

Theorem 1.7. *If \mathbf{a} is bounded by a 2-generic or is computably traceable, then \mathbf{a} is u-low. Conversely, any degree which computes a DNR_2 -real or is high is not u-low.*

We then turn our attention to the structure on the class of all ultrafilters provided by the construction $\mathcal{U} \mapsto \delta_{\mathcal{U}}$. Our main result is that the partial order induced by this construction is related to a classical reducibility notion on ultrafilters:

Definition 1.8. For ultrafilters \mathcal{U}, \mathcal{V} , we write “ $\mathcal{U} \leq \mathcal{V}$ ” if $\delta_{\mathcal{U}}(\mathbf{a}) \subseteq \delta_{\mathcal{V}}(\mathbf{a})$ for all degrees \mathbf{a} on some cone, and “ $\mathcal{U} \equiv \mathcal{V}$ ” if $\mathcal{U} \leq \mathcal{V}$ and $\mathcal{V} \leq \mathcal{U}$.

Theorem 1.9. *The partial order on ultrafilters induced by \leq is a quotient of the Rudin-Keisler ordering of ultrafilters on ω .*

We also show that the operation of composition of ultrafilter jumps is captured by a binary operation on ultrafilters:

Theorem 1.10. *There is a binary operation $*$ such that for every pair of ultrafilters \mathcal{U} and \mathcal{V} , we have*

$$\delta_{\mathcal{U}} \circ \delta_{\mathcal{V}} = \delta_{\mathcal{U} * \mathcal{V}}.$$

This operation is immediately seen to be compatible with the ordering, \leq , so that we have the structure of a partially ordered semigroup.

Finally, we end by presenting two directions for further research. Additionally, throughout this paper we raise a number of questions arising from the theorems above, which remain open.

Throughout this paper, we will need the following pair of basic combinatorial facts:

Definition 1.11. A collection $\{X_i : i \in I\}$ of sets is *free* if every finite Boolean combination of sets X_i for distinct i is infinite. In particular, each X_i and its complement must be infinite and the X_i must be distinct.

Fact 1.12. *Suppose $\{X_i : i \in I\}$ is free, and $J \subseteq I$. Then there is an ultrafilter \mathcal{U} with $\{i \in I : X_i \in \mathcal{U}\} = J$.*

Fact 1.13. *We can effectively find large free sets. Specifically, there is a total Φ_e such that*

$$\{\Phi_e^X : X \subseteq \omega\}$$

is a free set.

Fact 1.12 is immediate. To prove 1.13, construct a computable function $\iota : 2^{<\omega} \rightarrow 2^{<\omega}$ which

- builds reals along paths: $\sigma \prec \tau \iff \iota(\sigma) \prec \iota(\tau)$, and

- forces all Boolean combinations to be large: for every $I \subseteq 2^n$, the set

$$\{j : \forall \sigma \in 2^n (\iota(\sigma)(j) = 1 \iff \iota(\sigma) \in I)\}$$

has size at least n .

We then let $\Phi_e^X = \iota(X)$. (Note that, in fact, we have $\Phi_e^X \equiv_T X$.)

Our notation and terminology are standard. For background on computability theory and set theory, we refer to [DH10] and [Jec03], respectively. As far as facts about ultrafilters are concerned, this paper is self-contained; however, the book “The theory of ultrafilters” by Comfort and Negreponis [CN74] contains a wealth of information on the topic, and we refer to it from time to time.

Finally, a word of reassurance: since ultrafilters usually arise in the context of set theory, it is reasonable to worry that answers to questions about the maps $\delta_{\mathcal{U}}$ may be independent of ZFC. However, since the action of $\delta_{\mathcal{U}}$ on a degree \mathbf{a} is determined by countably much information about \mathcal{U} , most relevant questions are at worst Π_2^1 , and hence set-theoretically absolute (see chapter 25 of [Jec03]). Indeed, with two exceptions set theory will not be a serious concern in this article. The exceptions are proposition 5.2 — where we examine properties of the partial order \leq on ultrafilters — and section 6.2, where we mention a set-theoretic direction for further research.

2. BASIC PROPERTIES OF $\delta_{\mathcal{U}}$

In the previous section, we motivated the study of the functions $\delta_{\mathcal{U}}$ by drawing a comparison with the Turing jump. We begin this section by elaborating on that analogy. The following lemma shows that each function $\delta_{\mathcal{U}}$ dominates the Turing jump in a completely uniform way:

Lemma 2.1. *There are Turing functionals $\Phi_{e_0}, \Phi_{e_1}, \Phi_{e_2}$ witnessing the following (for every ultrafilter \mathcal{U}):*

- (1) $\delta_{\mathcal{U}}$ grows at least as fast as the Turing jump: for every $Y = \lim f(x, s)$, we have $\lim_{\mathcal{U}}(\Phi_{e_0}^f) = Y$.
- (2) $\delta_{\mathcal{U}}$ strictly dominates the Turing jump: for every set X , we have $\lim_{\mathcal{U}}(\Phi_{e_1}^X) \notin \Delta_2^0(X)$.
- (3) For every set X , we have $\lim_{\mathcal{U}}(\Phi_{e_2}^X) \not\leq_T \lim_{\mathcal{U}}(X)$. In particular, $\delta_{\mathcal{U}}(\text{deg}(X))$ has no top element.

Proof. (1) follows from the relativized limit lemma. Suppose f is a total X -computable function such that

$$\forall x, \lim_{s \rightarrow \infty} f(x, s) \downarrow = Y(x).$$

Let Φ_{e_0} be defined by

$$\Phi_{e_0}^f(\langle i, j \rangle) = f(j, i).$$

Then since \mathcal{U} contains all cofinite sets we have $\lim_{\mathcal{U}}(\Phi_{e_0}^f) = Y$.

For (2), say that a set Z has the *limit property* if for all j , $\lim_{i \rightarrow \infty} Z(\langle i, j \rangle)$ exists. To prove (2) we need only construct a $Z \leq_T X$ such that for all nonprincipal \mathcal{U} and all $\hat{Z} \leq_T X$ with the limit property, we have $\lim_{\mathcal{U}}(Z) \neq \lim_{\mathcal{U}}(\hat{Z})$. To do this, we proceed as follows. For $e, s \in \omega$, let

$$n_{e,s} = \max\{i : \Phi_e^X(\langle i, e \rangle)[s] \downarrow\}, \quad v_{e,s} = \Phi_e^X(\langle n_{e,s}, e \rangle)$$

(with the convention that $v_{e,s} = 0$ if $n_{e,s}$ is undefined). Now let Z be defined by

$$Z(\langle k, e \rangle) = 1 - v_{e,k},$$

and note that $Z \leq_T X$. The proof of (2) is completed by noting that whenever Φ_e^X is the characteristic function of a set with the limit property, then

$$\lim_{k \rightarrow \infty} Z(\langle k, e \rangle) \downarrow = 1 - \lim_{k \rightarrow \infty} \Phi_e^X(\langle k, e \rangle),$$

so $\lim_{\mathcal{U}}(Z)(e) = 1 - \lim_{\mathcal{U}}(\Phi_e^X)(e)$, and hence $\lim_{\mathcal{U}}(Z)$ is not Δ_2^0 . This construction, moreover, is effective, so we get the desired index e_1 .

The proof of (3) is similar to that of (2). \square

Lemma 2.1(3) shows that $\delta_{\mathcal{U}}$ is never a function from degrees to degrees. This raises the problem of classifying the possible images of $\delta_{\mathcal{U}}(\mathbf{a})$.

Lemma 2.2. $\delta_{\mathcal{U}}(\mathbf{a})$ is a Turing ideal, that is, it is closed under \oplus and \leq_T .

Proof. Showing closure under \oplus is easy: $\lim_{\mathcal{U}}(\{A_i\}_{i \in \omega}) \oplus \lim_{\mathcal{U}}(\{B_i\}_{i \in \omega}) = \lim_{\mathcal{U}}(\{A_i \oplus B_i\}_{i \in \omega})$. To show that $\delta_{\mathcal{U}}(\mathbf{a})$ is closed under \leq_T , fix $A = (A_i)_{i \in \omega}$ and suppose $\Phi_e^{\lim_{\mathcal{U}}(A)} = B$. Then let

$$C_i = \{j : \Phi_e^{A_i}(j)[i] \downarrow = 1\}$$

and let $C = (C_i)_{i \in \omega}$. We claim that $\lim_{\mathcal{U}}(C) = B$. To see this, fix $k \in \omega$. There is some initial segment $\sigma \prec \lim_{\mathcal{U}}(A)$ such that $\Phi_e^\sigma(k) \downarrow$; since ultrafilters are closed under finite intersections, for \mathcal{U} -many i we have $\sigma \prec A_i$, and for cofinitely many i we have $i > |\sigma|$. Together, these facts imply that for \mathcal{U} -many i we have $\sigma \prec C_i$, and so $\sigma \prec \lim_{\mathcal{U}}(C)$. \square

In fact, an even stronger closure property is satisfied:

Proposition 2.3. For every ultrafilter \mathcal{U} and degree \mathbf{a} , $\delta_{\mathcal{U}}(\mathbf{a})$ is a Scott set. In fact, as in Lemma 2.1 this is uniform: there is a single $e \in \omega$ such that for all X and \mathcal{U} , we have

$$\lim_{\mathcal{U}}(X) \text{ is an infinite subtree of } 2^{<\omega} \quad \Rightarrow \quad \lim_{\mathcal{U}}(\Phi_e^X) \text{ is a path through } \lim_{\mathcal{U}}(X).$$

Proof. The intuition behind this proof is that a tree T in $\delta_{\mathcal{U}}(\mathbf{a})$ must be “named” by a sequence of trees $(X_i)_{i \in \omega}$ in \mathbf{a} , which — if T is to be infinite — must have arbitrarily long paths. By producing a sequence of increasingly long paths through this sequence of trees, we produce a sequence of sets in \mathbf{a} which \mathcal{U} sends to an infinite path through the named tree. Note that this is intuitively the same argument as for closure under Turing reducibility.

The details are as follows. Suppose $X = (X_i)_{i \in \omega} \in \mathbf{a}$ is such that $T = \lim_{\mathcal{U}}(X)$ is an infinite subtree of $2^{<\omega}$. First, we can assume without loss of generality that each column X_i is also a tree (i.e., downwards closed). To see this, let Y_i be the downwards-closed part of X_i , and let $Y = (Y_i)_{i \in \omega} \in \mathbf{a}$. Since $Y \subseteq X$ we have $\lim_{\mathcal{U}}(Y) \subseteq \lim_{\mathcal{U}}(X) = T$, and the former is clearly a tree; so any path we build through $\lim_{\mathcal{U}}(Y)$ will also be a path through T . (In fact, it is not hard to see that we have $\lim_{\mathcal{U}}(Y) = \lim_{\mathcal{U}}(X)$.)

So assume X is a sequence of trees. Then X computes a sequence $P = (f_i)_{i \in \omega}$ of sets $f_i \subseteq X_i$ such that f_i is a finite path through X_i of maximal length $\leq i$ (the “ $\leq i$ ” is required to make this search effective). We claim that $\lim_{\mathcal{U}}(P)$ is an infinite path through T .

Clearly $\lim_{\mathcal{U}}(P) \subseteq T$, is closed downwards, and is a path in T (that is, any two elements are comparable); so it is enough to show that $\lim_{\mathcal{U}}(P)$ is infinite. Towards a contradiction, suppose $\sigma \in \lim_{\mathcal{U}}(P)$ of length n is terminal. Then since ultrafilters are closed under finite intersections, we have that for \mathcal{U} -many i , $f_i \supseteq \sigma$. Moreover, by definition of P , for all but n -many i , we have

$$|f_i| \leq n \iff ht(X_i) \leq n.$$

Together these imply that for \mathcal{U} -many i , X_i has height at most n , and so $\lim_{\mathcal{U}}(X)$ has height at most n as well, which is a contradiction.

By examining the argument above, it is clear that this is a uniform construction — that is, the construction of P is uniformly computable in X and does not depend on \mathcal{U} . \square

Remark 2.4. Proposition 2.3 yields an alternate proof of the classical result in reverse mathematics that the theory WKL_0 is strictly weaker than the theory ACA_0 (see chapter VIII of [Sim99]), as follows: letting \mathcal{U} be an ultrafilter such that the set $\{e : W_e \in \mathcal{U}\}$ is Δ_4^0 ensures that $\delta_{\mathcal{U}}(\mathbf{0})$ consists entirely of Δ_4^0 sets, and so is not arithmetically closed. This is genuinely different from the standard proof, which follows from iterating the Low Basis Theorem. In particular, neither lowness nor iterated forcing are used in the proof of Proposition 2.3.

In the following section, we will show that the converse of Proposition 2.3 holds: given any countable Scott set \mathfrak{S} containing $\mathbf{0}'$, there is a nonprincipal ultrafilter \mathcal{U} such that $\delta_{\mathcal{U}}(\mathbf{0}) = \mathfrak{S}$, and more generally if $\mathbf{a}' \in \mathfrak{S}$ then we can find a \mathcal{U} with $\delta_{\mathcal{U}}(\mathbf{a}) = \mathfrak{S}$.

3. BUILDING SCOTT SETS

We now completely characterize the possible images of ultrafilter jumps by proving the converse of Proposition 2.3. This does not provide a characterization of the maps $\delta_{\mathcal{U}}$, however, since we only determine the possible *local* behaviors of those maps, and indeed a characterization of the possible $\delta_{\mathcal{U}}$ s seems well out of reach. However, in the next section we do make progress towards this goal, by studying what sorts of simultaneous behaviors can be realized by ultrafilter jumps.

Theorem 3.1. *Let \mathbf{a} be a degree, and let \mathcal{I} be a countable Scott set containing \mathbf{a}' . Then $\mathcal{I} = \delta_{\mathcal{U}}(\mathbf{a})$ for some nonprincipal ultrafilter \mathcal{U} .*

Proof. Call a pair (A, B) with $A \in \mathbf{a}$ and $B \in \mathcal{I}$ an *axiom*; informally, we interpret (A, B) as meaning “ A is mapped to B by $\lim_{\mathcal{U}}$.” Precisely, for \mathcal{C} a set of axioms, say that an ultrafilter \mathcal{U} *satisfies* \mathcal{C} if $\lim_{\mathcal{U}}(A) = B$ whenever $(A, B) \in \mathcal{C}$. Since every family of sets, all of whose finite intersections are infinite, can be extended to a nonprincipal ultrafilter, satisfiability has a purely *combinatorial* definition: if $\mathcal{C} = \{(A_i, B_i) : i \in I\}$ is a set of axioms, we say \mathcal{C} is *consistent* if for every $F \subseteq I$ finite and $n \in \omega$, the intersection

$$\left[\bigcap_{j \in F, m < n, B_j(m)=1} (A_j)^m \right] \cap \left[\bigcap_{j \in F, m < n, B_j(m)=0} \overline{(A_j)^m} \right]$$

is infinite. Equivalently, \mathcal{C} is consistent if and only if there is a nonprincipal ultrafilter satisfying \mathcal{C} .

Fix enumerations $\mathcal{I} = \{Y_i : i \in \omega\}$ and $\mathbf{a} = \{X_i : i \in \omega\}$; we will build fragments of the desired ultrafilter in stages. We will build a consistent set of axioms \mathcal{C} such that (i) for every $A \in \mathbf{a}$ there is some $B \in \mathcal{I}$ with $(A, B) \in \mathcal{C}$, and (ii) for every $B \in \mathcal{I}$ there is some $A \in \mathbf{a}$ such that $(A, B) \in \mathcal{C}$. Then since \mathcal{C} is consistent, we can take any ultrafilter \mathcal{U} satisfying \mathcal{C} . For this \mathcal{U} , we have then ensured that $\mathcal{I} = \delta_{\mathcal{U}}(\mathbf{a})$.

We handle (i) at even stages, and (ii) at odd stages:

- In (i), in deciding where to map a set $A \in \mathbf{a}$ we run the risk of contradicting already-enumerated axioms $(A_i, B_i)_{i < k}$ — for example, if the fifth rows of A and A_0 are identical, then the fifth bit of B must be $B_0(5)$. To find a $B \in \mathcal{I}$ to which it is “safe” to map A , it turns out to be equivalent to find a path through a certain infinite binary tree computable in the (finitely many) axioms built so far.
- In (ii), an apparent difficulty is posed by the fact that \mathbf{a} cannot “see” the commitment we have already made, since right components of axioms lie outside \mathbf{a} ; however, this turns out not to matter. Suppose $B \in \mathcal{I}$ and $(A_i, B_i)_{i < k}$ is consistent; then if A is “sufficiently different” from the A_i s, the set $(A_i, B_i)_{i < k} \cup \{(A, B)\}$ is also consistent. So in deciding what should be mapped to B , we ignore B entirely, and simply choose some A which is sufficiently different from the sets we’ve enumerated on the left so far.

Formally, we proceed as follows:

Even stage. Suppose that we have $\mathcal{C}_{2s} = \{(A_i, B_i) : i < 2s\}$, and that \mathcal{C}_{2s} is consistent, and consider the set $X_s \in \mathbf{a}$. We will find a $B \in \mathcal{I}$ such that $\mathcal{C}_{2s} \cup \{(X_s, B)\}$ is consistent. Let $D = \{(A_i)^j : i < 2s, B_i(j) = 1\} \cup \{\mathbb{N} - (A_i)^j : i < 2s, B_i(j) = 0\}$; intuitively, D is the collection of sets we've guaranteed are in the ultrafilter so far. Write $D = \{D_k : k \in \omega\}$, and note that this can be done effectively in $A_0 \oplus \bigoplus_{i < s} B_i := \hat{B} \in \mathcal{I}$. Say that $\sigma \in 2^{<\omega}$ is temporarily consistent if $|\sigma| = n$ and $\forall m < n$,

- $\sigma(m) = 1 \Rightarrow |(X_s)^m \cap (\bigcap_{j < n} D_j)| \geq n$, and
- $\sigma(m) = 0 \Rightarrow |(\mathbb{N} - (X_s)^m) \cap (\bigcap_{j < n} D_j)| \geq n$;

note that $\hat{B} \oplus X'_s$ can uniformly decide whether a $\sigma \in 2^{<\omega}$ is temporarily consistent. Let $T \subseteq 2^{<\omega}$ be the tree of temporarily consistent nodes; since \mathcal{C}_{2s} is consistent by induction, T is infinite, and since \mathcal{I} is a Scott ideal containing X'_s and \hat{B} , there is some $B \in \mathcal{I}$ whose characteristic function is a path through T . Then $\mathcal{C}_{2s} \cup \{(X_s, B)\}$ is consistent, so let $\mathcal{C}_{2s+1} = \mathcal{C}_{2s} \cup \{(X_s, B)\}$.

Odd stage. Suppose that we have a consistent set of axioms $\mathcal{C}_{2s+1} = \{(A_i, B_i) : i < 2s+1\}$, and consider the set $Y_s \in \mathcal{I}$; we need to find some $A \in \mathbf{a}$ such that $\mathcal{C}_{2s+1} \cup \{(A, Y_s)\}$ is consistent. Our main difficulty is that the condition \mathcal{C}_{2s+1} we have built so far is not \mathbf{a} -computable — in \mathbf{a} , we can only see $\{A_i : i < 2s+1\}$ — so in order to guarantee consistency we will need to ensure that the axiom (A, Y) is consistent with any possible consistent set of axioms with left coordinates from among the A_i ($i < 2s+1$). To do this, we use a modification of Fact 1.13:

Definition 3.2. *A set X is free over a family of sets $Z = \{Z_i : i \in \omega\}$ if every Boolean combination of finitely many elements of Z , which is infinite, has infinite intersection with both X and $\omega - X$.*

Lemma 3.3. *We can find free sets in a uniformly effective manner. Specifically, there is an e so that for all $Z = \{Z_i : i \in \omega\}$, $X = \Phi_e^Z$ is free over $\{Z_i : i \in \omega\}$.*

Proof. We need to build X such that for every set B which can be written as a Boolean combination of finitely many elements of Z , either B is finite or both $B \cap X$ and $B \cap \bar{X}$ are infinite. Let $(B_i)_{i \in \omega}$ be an enumeration of all Boolean combinations of elements of Z , with each combination occurring infinitely often. Further, make $(B_i)_{i \in \omega}$ be so that $B_{2i} = B_{2i+1}$ for all i . Note that such an enumeration can be chosen recursively in Z . At the beginning of stage 0, say that all i await attention. At stage s , suppose we have defined a string $p_s \in 2^{<\omega}$ with length n_s . Say that j requires attention if $j < s$, and at the beginning of stage s , j awaits attention, and $n_s \in B_j$. Let i be the least number which requires attention, and let $p_{s+1} = p_s \widehat{\langle 1 \rangle}$ if i is even and $p_{s+1} = p_s \widehat{\langle 0 \rangle}$ if i is odd. From now on, say that i is satisfied, meaning it never again will await attention, and move on to stage $s+1$. Letting $X = \bigcup p_s$, we have actively ensured that if B is a boolean combination of finitely many elements of Z , then both $B \cap X$ and $B \cap \bar{X}$ are infinite. \square

To finish the description of the odd stages for Theorem 3.1, we iterate Lemma 3.3 to build an $X \in \mathbf{a}$ such that for each $k \in \omega$, X^k is free over $\{(A_i)^j : i < 2s+1, j \in \omega\} \cup \{X^j : j < k\}$; we then take $\mathcal{C}_{2s+2} = \mathcal{C}_{2s+1} \cup \{(X, Y_s)\}$. \square

Having completely classified the sets of the form $\delta_{\mathcal{U}}(\mathbf{a})$ in terms of \mathbf{a} , we now face the question of classifying ultrafilter jumps themselves:

Question 1. What conditions on a function $f: \{\text{Turing degrees}\} \rightarrow \{\text{Scott sets}\}$ ensure that $f = \delta_{\mathcal{U}}$ for some \mathcal{U} ?

This is, of course, a massively broad question. One interesting special case is the following:

Question 2. Is there an ultrafilter \mathcal{U} such that $\delta_{\mathcal{U}}(\mathbf{a})$ is always arithmetically closed?

This is partly motivated by Remark 2.4, which suggests that there may be further interaction between the study of the maps $\delta_{\mathcal{U}}$ and reverse mathematics.

Currently it is not clear how to approach this type of problem, largely because constructing ultrafilter jumps “to order” is quite difficult. We make some technical progress in this direction, however, in the following section, in which we study what *simultaneous* behaviors can be realized by ultrafilter jumps.

4. LOWNESS NOTIONS

Theorem 3.1 allows us to control the value of $\delta_{\mathcal{U}}(\mathbf{a})$ for a fixed degree \mathbf{a} ; however, it says nothing about what *simultaneous* behaviors can occur.

First of all, it is obvious that if $\mathbf{b} \geq \mathbf{a}$, then $\delta_{\mathcal{U}}(\mathbf{b}) \supset \delta_{\mathcal{U}}(\mathbf{a})$, and so one particularly interesting question is the following: for what degrees \mathbf{a} is there an ultrafilter \mathcal{U} such that $\delta_{\mathcal{U}}(\mathbf{a}) = \delta_{\mathcal{U}}(\mathbf{0})$? We will call such a degree *u-low*, and we will call a set *u-low* if it belongs to a *u-low* degree.

It is easy to see that $\mathbf{0}'$ is not *u-low*: By a standard diagonalization argument, $\mathbf{0}'$ computes an array $A = (A_i)_{i \in \omega}$ such that the e th row of the e th computable array has no agreement with A_e . More precisely:

Proposition 4.1. *If \mathbf{a} contains a DNR_2 , then \mathbf{a} is not *u-low*.*

Proof. Such a degree \mathbf{a} contains a set A such that for every e , if Φ_e is total then

$$\Phi_e(\langle i, e \rangle) \neq A(\langle i, e \rangle).$$

It follows that we can never have $\lim_{\mathcal{U}}(A) = \lim_{\mathcal{U}}(C)$ for any computable array C . \square

As an aside, note that this rules out the most natural possible positive answer to Question 2:

Corollary 4.2. *No ultrafilter jump $\delta_{\mathcal{U}}$ is the “arithmetic closure” operator. That is, for every \mathcal{U} there is some \mathbf{a} such that $\delta_{\mathcal{U}}(\mathbf{a}) \neq \text{ARITH}(\mathbf{a})$.*

However, this does not rule out the existence of ultrafilters which are arithmetically closed in pathological ways, so Question 2 remains open.

In addition, high degrees are not *u-low*. Recall that a degree \mathbf{a} is high if $\mathbf{a}' \geq \mathbf{0}''$.

Proposition 4.3. *If \mathbf{a} is high, then \mathbf{a} is not *u-low*.*

Proof. By Martin’s Lemma, such a degree \mathbf{a} computes a *dominant* function f which dominates every computable function. Using f we can compute a set A such that

$$\Phi_e(\langle i, e \rangle) \neq A(\langle i, e \rangle)$$

is true cofinitely often for each $e \in \text{Tot}$, i.e., for each e such that Φ_e is total. So as in the DNR_2 case we are done. \square

In light of Propositions 4.1 and 4.3, it is reasonable to ask whether *any* nonzero degree is *u-low*. In fact, many degrees are *u-low*, including every 2-generic and every computably traceable degree. We begin with a combinatorial lemma:

Lemma 4.4. *Suppose $\{A_i : i \in \omega\}$ and $\{X_i : i \in \omega\}$ are collections of sets of natural numbers. Then the following are equivalent:*

- (1) *There is an ultrafilter \mathcal{U} such that for all i , $\lim_{\mathcal{U}}(A_i) = \lim_{\mathcal{U}}(X_i)$. (Note that here we think of each A_i and X_i as an array of sets, so they will each have their own rows $(A_i)^j, (X_i)^j$ and columns $(A_i)_k, (X_i)_k$.)*
- (2) *For every n, k , there is some $m > k$ such that for every $i, j < n$, we have*

$$(A_i)^j(m) = (X_i)^j(m).$$

If $\mathcal{C} = \{(A_i, X_i) : i \in \omega\}$ is a collection of pairs of sets such that the above conditions hold, we call \mathcal{C} a *consistent system*; note that this is a different sense of consistency that that used in Theorem 3.1.

Proof. Let $\mathcal{D} = \{D_{i,j} : i, j \in \omega\}$ where $D_{i,j} = \{x : (A_i)^j(x) = (X_i)^j(x)\}$ is the set on which the j th rows of A_i and X_i agree. Condition (1) holds if and only if \mathcal{D} is contained in an ultrafilter, i.e., \mathcal{D} has the finite intersection property, which is equivalent to (2). \square

Theorem 4.5. *Every degree bounded by a 2-generic is u-low.*

Recall that a set X with characteristic function f is 2-generic if (when viewed as a filter in the poset $2^{<\omega}$) it *meets or avoids* every Σ_2^0 subset of $2^{<\omega}$: if $A \subseteq 2^{<\omega}$ is Σ_2^0 and $f|n \notin A$ for every n , then $\exists \tau \prec f(\forall \sigma \succ \tau, \sigma \notin A)$.

Proof. Fix G 2-generic; we will construct a \mathcal{U} such that $\delta_{\mathcal{U}}(\deg(G)) = \delta_{\mathcal{U}}(\mathbf{0})$. Let

$$\text{Tot}_G = \{e_0 < e_1 < \dots\} = \{e : \Phi_e^G \text{ is total}\}.$$

For $i \in \omega$, let t_i be the first condition in G such that $t_i \Vdash \text{“}\Phi_{e_j}^G \text{ is total”}$ for every $j \leq i$; note that such conditions exist since G is 2-generic. This is the only point in the proof where full 2-genericity is required. (We do not need to take the *least* such conditions, but we do need the t_i s to be successively stronger conditions: $t_0 \geq t_1 \geq \dots$.) Let $\mathbb{P} = \{p_j : j \in \omega\}$ be a listing of Cohen conditions.

We will construct recursive sets X_i such that there is an ultrafilter which maps X_i and $\Phi_{e_i}^G$ to the same set. These X_i will be defined column-by-column, with each column making an increasingly strong guess as to the corresponding column of $\Phi_{e_i}^G$. The complexity of the construction comes from the fact that these guesses must be made effectively, and also must cohere with each other; this second requirement is the reason for having X_i take into account the Φ_{e_j} with $j < i$ in the construction below.

Construction 4.6. We define the recursive sets X_i ($i \in \omega$) as follows:

- (1) For $p_k \not\leq t_i$, the k th column of X_i is empty: $\{\langle k, j \rangle : j \in \omega\} \cap X = \emptyset$.
- (2) For $p_k \leq t_i$, let $\mathfrak{p}_{i,k} = ((\sigma_j : j \leq i), q)$ be the lexicographically least pair such that $q \leq p_k$ and

$$\forall j \leq i, m < i \quad \Phi_{e_j}^q(\langle k, m \rangle) = \sigma_j(m);$$

note that such a pair must always exist since $p_k \leq t_i$. Then let the k th column of X_i be $\sigma_i \frown 0^\infty$.

We claim that there is an ultrafilter \mathcal{U} such that $\lim_{\mathcal{U}}(\Phi_{e_i}^G) = \lim_{\mathcal{U}}(X_i)$ for every i . By Lemma 4.4, it is enough to show that the pair of sequences

$$\{\Phi_{e_i}^G : i \in \omega\}, \{X_i : i \in \omega\}$$

satisfies the property 4.4(1).

To show this, fix n, k , and consider the set of conditions

$$E_{n,k} = \{p \in \mathbb{P} : (\exists m > k)(\forall i, j < n) ((\Phi_{e_i}^p)^j(m) \downarrow = (X_i)^j(m))\}.$$

We will use genericity of G to show that G meets the Σ_1^0 -set $E_{n,k}$.

It suffices to show that $E_{n,k}$ is dense below t_n — the 2-genericity of G , together with the fact that $E_{n,k}$ is Σ_1^0 , means that G must then meet $E_{n,k}$. Towards this, we fix some condition $p \leq t_n$. There must be some m such that $m > k$ and $p_m \leq p$. Since $p_m \leq p \leq t_n$, the m th column of X_n was constructed in Case (2) of Construction 4.6. Let $\mathbf{p}_{n,k} = ((\sigma_j : j \leq i), q)$ be as defined in the construction of X_n , and consider the condition q ; we claim $q \in E_{n,k}$.

To see this, note that by the construction of the m th column of X_i and the fact that $m > k$, we have, for every $i, j < n$,

$$(\Phi_{e_i}^q)^j(m) = (X_i)^j(m).$$

Thus the set $E_{n,k}$ is dense below t_n , and thus G meets it, verifying condition (2) of Lemma 4.4. \square

We ask the analogous question for measure:

Question 3. Are sufficiently random reals u -low?

By a similar argument to Theorem 4.5, we can show that another important computability-theoretic property implies u -lowness:

Theorem 4.7. *Every computably traceable degree is u -low.*

Recall that a degree \mathbf{a} is computably traceable if for every $h \in \mathbf{a}$, there is a computable j such that $h(n) \in D_{j(n)}$ and $|D_{j(n)}| \leq 2^n$ for every n , where D_e is the canonical finite set coded by e . Note that since there are computably traceable degrees which are not 2-generic and vice versa, Theorems 4.7 and 4.5 compliment each other.

Proof. Let $\{A_i : i \in \omega\}$ be a list of all sets of degree $\leq_T \mathbf{a}$. As in the proof of Theorem 4.5, we will construct a collection $\{X_i : i \in \omega\}$ of recursive sets such that there is some ultrafilter \mathcal{U} satisfying $\lim_{\mathcal{U}}(X_i) = \lim_{\mathcal{U}}(A_i)$ for every i . This ultrafilter will then satisfy $\delta_{\mathcal{U}}(\mathbf{a}) = \delta_{\mathcal{U}}(\mathbf{0})$.

To construct the X_i , we work in stages. Each X_i will have associated with it three functions: the *interval* function f_i , the *block* function g_i , and the *guessing* function h_i . We view A_i and X_i as arrays in the usual way, so that $A_i, X_i \subseteq \omega^2$. In order to construct X_i , we partition the full array ω^2 into “blocks,” and partition the n th block into 2^n -many “intervals,” and define X_i on each interval separately.

The functions g_i and f_i tell us how to perform this construction: $g_i(n)$ is the number of columns in the n th block, and $f_i(m)$ is the number of columns in the m th interval. (Recall that each block will be partitioned into exponentially-many intervals.) Now we let h_i be a computable function such that for every k , $D_{h_i(k)}$ is a finite set of size 2^k listing the possible behaviors of A_i on the (finitely many) values in the k th block and above the diagonal $\{\langle s, s \rangle : s \in \omega\}$; the existence of such an h_i is guaranteed by the assumption that \mathbf{a} is computably traceable. We then define X_i so that X_i agrees with A_i on at least one interval in each block, by predicting A_i 's behavior on the t th interval using the t th element of $D_{h_i(k)}$.

This describes the process for building a single X_i . To ensure that agreements between the X_i s and the A_i s occur across i 's, we make intervals of X_{i+1} correspond to blocks of A_i ; this guarantees that the collection of pairs $\{(A_i, X_i) : i \in \omega\}$ forms a consistent system (see Lemma 4.4).

Precisely, the construction is the following:

- At stage 0 we have $f_0 : x \mapsto 1$ and $g_0 : x \mapsto 2^x$.

- At stage $i+1$, blocks from stage i become intervals and the new blocks are exponentially large collections of intervals. That is, $f_{i+1} = g_i$, $g_{i+1}(0) = f_{i+1}(0)$, and

$$g_{i+1}(n+1) = \sum_{j=2^{n+1}-1}^{2^{n+2}-1} f_{i+1}(j).$$

The h_i are then computable maps such that for every x , (the canonical code for) the finite set

$$A_i \upharpoonright \{ \langle m, n \rangle : m \leq n \text{ and } \sum_{t=0}^{x-1} g_i(t) \leq n < \sum_{t=0}^x g_i(t) \}$$

is an element of $D_{h_i(x)} = \{s_1^{i,x} < s_2^{i,x} < \dots < s_{2^x}^{i,x}\}$.

We then let X_i be defined by copying the set coded by $s_m^{i,x}$ on the m th interval in the x th block. It is easy to see that for every j , there is at least one interval in the j th block such that X_i and A_i agree on the first j -many rows. Since our construction nests blocks at level i inside intervals at level $i+1$, it now follows from Lemma 4.4 that the family $\{(A_i, X_i) : i \in \omega\}$ is consistent. \square

Another collection of u -low degrees are the low c.e. degrees¹.

Theorem 4.8. *Every low c.e. degree is u -low.*

Proof. Let A be a low c.e. set. We will use Robinson low guessing to certify computations from A in order to ensure condition (2) of Lemma 4.4.

For a low c.e. set A , we fix a Δ_2^0 approximation to A' . Then to “certify” a computation $\Phi_e^{A_s}(x) = m$ (i.e., we want to certify that the s -stage approximation A_s to A is correct below the use of this computation), we place the use σ of this computation into a c.e. set V that we construct, and we “certify” the computation if our Δ_2^0 -approximation to A' then declares that there is some $\tau \in V$ so that $\tau \prec A$. Note that we either see the computation certified or A changes below the use of the computation, and we know not to trust the computation. If the computation is certified and then A changes below the use of the computation then the Δ_2^0 -approximation to the statement $(\exists \tau \in V)(\tau \prec A)$ must change back to false. We then use the same set V to “certify” the next computation. In this way, it is impossible for each one of the infinitely many computations certified via V to each be incorrect, as the Δ_2^0 -approximation to A' must settle down. We will in fact construct a uniform sequence $(V_i)_{i \in \omega}$ of c.e. sets to give us infinitely many correct certified computations, since each V_i only gives us one.²

We fix $\text{Tot}_A = \{e_0 < e_1 < \dots\}$, and we construct a sequence $\{X_i : i \in \omega\}$ so that the family $\{(\Phi_{e_i}^A, X_i) : i \in \omega\}$ is consistent.

If $n > i$, we let the i th column of n be empty. At stage s , we define the s th column of each X_n with $n \leq s$: We wait to simultaneously certify all the computations $\Phi_{e_n}^{A_s}(\langle m, j \rangle)$ for all $n, m, j \leq s$. That is, for the least i so that the Δ_2^0 -approximation declares the statement $(\exists \tau \in V_i)(\tau \prec A)$ false, we enter the greatest use from A of any of these computations into V_i and wait until either A changes below this use or the Δ_2^0 -approximation to A' changes to declare this statement true. Once we see the latter outcome (which must happen at some point, since all these computations do in fact converge), we define each $X_n(\langle s, j \rangle)$ for $j \leq s$ to be the now-certified computation $\Phi_{e_n}^{A_s}(\langle s, j \rangle)$. For $j > s$, we let $X_n(\langle s, j \rangle) = 0$.

¹We thank the anonymous referee for suggesting this result and its proof.

²A further summary of Robinson’s guessing method can be found in Soare [Soa87, detailed hint to exercise XI.3.5], summarizing the original argument in Robinson [Rob66]. An alternate approach, via cost functions, can be found, e.g., in Nies [Nie06, proof of Theorem 5.1].

Since each V_i must eventually give us one correct certified computation, this gives us some s so that each X_n for $n \leq s$ simultaneously correctly guesses at $\Phi_{e_n}^A$ for each of the first s rows. By condition (2) of Lemma 4.4, this guarantees that the family $\{(\Phi_{e_i}^A, X_i) : i \in \omega\}$ is consistent. \square

The above results provide a wide swath of u -low degrees. However, our knowledge of u -lowness is still very incomplete. The following question, of course, remains extremely open:

Question 4. Is there an exact characterization of u -lowness in terms of classical computability-theoretic properties?

Finally, while investigating u -lowness, an even stronger notion of weakness with respect to ultrafilters arises. We say a degree \mathbf{a} is u -trivial if $\delta_{\mathcal{U}}(\mathbf{a}) = \delta_{\mathcal{U}}(\mathbf{0})$ for every ultrafilter \mathcal{U} .

Question 5. Is there a nonzero u -trivial degree?

5. COMPARING ULTRAFILTERS

We now turn to what the construction of the maps $\delta_{\mathcal{U}}$ can tell us about the set of ultrafilters. We begin by defining a natural preorder arising from these maps, and then turn to a natural associated algebraic (semigroup) structure; we end by presenting a connection with a classical structure on ultrafilters, the *Rudin-Keisler order*.

Definition 5.1. For \mathcal{U}, \mathcal{V} ultrafilters, let $\mathcal{U} \leq \mathcal{V}$ if for some degree \mathbf{b} , we have $\delta_{\mathcal{U}}(\mathbf{a}) \subseteq \delta_{\mathcal{V}}(\mathbf{a})$ for all $\mathbf{a} \geq_T \mathbf{b}$; that is, $\mathcal{U} \leq \mathcal{V}$ if $\delta_{\mathcal{V}}$ dominates $\delta_{\mathcal{U}}$ on a cone. We write \mathcal{D}_{ult} for the resulting partial order on (equivalence classes of) ultrafilters.

Note that $\mathcal{U} < \mathcal{V}$ does *not* imply that, on a cone, $\delta_{\mathcal{U}}(\mathbf{a}) \subsetneq \delta_{\mathcal{V}}(\mathbf{a})$. Indeed, it is not clear whether such a situation ever occurs.

Question 6. Are there \mathcal{U}, \mathcal{V} such that $\delta_{\mathcal{U}}(\mathbf{a}) \subsetneq \delta_{\mathcal{V}}(\mathbf{a})$ for all \mathbf{a} (on a cone)?

Proposition 5.2. \mathcal{D}_{ult} is 2^{\aleph_0} -directed: given any 2^{\aleph_0} -sized collection \mathcal{C} of ultrafilters, there is a \mathcal{V} with $\mathcal{U} < \mathcal{V}$ for every $\mathcal{U} \in \mathcal{C}$.

Proof. We use Fact 1.13 to construct an ultrafilter \mathcal{V} which dominates each $\mathcal{U} \in \mathcal{C}$ on a cone. Since the set of Turing degrees \mathcal{D} has size 2^{\aleph_0} , we can enumerate $\mathcal{C} = \{\mathcal{U}_{\mathbf{d}} : \mathbf{d} \in \mathcal{D}\}$. Fix for each Turing degree \mathbf{a} , some Turing degree $\mathbf{b}_{\mathbf{a}}$ so that $\mathbf{b}_{\mathbf{a}}$ is an upper bound for $\delta_{\mathcal{U}_{\mathbf{c}}}(\mathbf{a})$ for every $\mathbf{c} \leq \mathbf{a}$. Since there are only countably many Turing degrees $\mathbf{c} \leq \mathbf{a}$ and each $\delta_{\mathcal{U}_{\mathbf{c}}}(\mathbf{a})$ is countable, such a $\mathbf{b}_{\mathbf{a}}$ can be found. Now, it suffices to build an ultrafilter \mathcal{V} so that $\mathbf{b}_{\mathbf{a}} \in \delta_{\mathcal{V}}(\mathbf{a})$ for every $\mathbf{a} \in \mathcal{D}$, since \mathcal{V} will then dominate $\mathcal{U}_{\mathbf{e}}$ on the cone above \mathbf{e} .

Using Fact 1.13, we fix an e so that $\{\Phi_e^X : X \subseteq \omega\}$ is a free family. For each $\mathbf{a} \in \mathcal{D}$, fix some non-empty $X_{\mathbf{a}} \in \mathbf{a}$. Then the family $\{\Phi_e^{0^n \frown X_{\mathbf{a}}} : n \in \omega\}$ is free and uniformly computable from $X_{\mathbf{a}}$. Let $Y_{\mathbf{a}}$ be the set whose n th row is $\Phi_e^{0^n \frown X_{\mathbf{a}}}$. Then since the family of rows of the sets $Y_{\mathbf{a}}$ for all $\mathbf{a} \in \mathcal{D}$ is a free family, Fact 1.12 shows that we can choose an ultrafilter \mathcal{V} to make $\lim_{\mathcal{V}}(Y_{\mathbf{a}})$ be anything we choose. We choose to make $\lim_{\mathcal{V}}(Y_{\mathbf{a}}) \in \mathbf{b}_{\mathbf{a}}$. This ensures that $\mathbf{b}_{\mathbf{a}} \in \delta_{\mathcal{V}}(\mathbf{a})$ for every $\mathbf{a} \in \mathcal{D}$. \square

Note that this argument cannot be easily extended to give upper bounds of larger sets of ultrafilters. Indeed, it is consistent that there are exactly ω_2 -many ultrafilters on ω , in which case 2^{\aleph_0} -directedness (note that here $2^{\aleph_0} = \omega_1$) is the most we could hope for.

Question 7. What can be said about $|\mathcal{D}_{ult}|$? (Note that we have $2^{\aleph_0} < |\mathcal{D}_{ult}| \leq 2^{2^{\aleph_0}}$. Moreover, it is consistent — and follows from GCH — that $|\mathcal{D}_{ult}| = 2^{2^{\aleph_0}}$.)

Additionally, the proof of Proposition 5.2 says nothing about the *optimality* of the upper bound constructed.

Question 8. What sets of ultrafilters have least upper bounds in \mathcal{D}_{ult} ?

Note that it is not even clear whether *finite* sets of ultrafilters have least upper bounds.

We now show that the set of ultrafilters carries a natural semigroup structure which is compatible with the degree structure \mathcal{D}_{ult} :

Definition 5.3. \mathcal{U}, \mathcal{V} ultrafilters, let

$$\mathcal{U} * \mathcal{V} = \{X : \{b : \{a : \langle a, b \rangle \in X\} \in \mathcal{V}\} \in \mathcal{U}\}.$$

It is clear that $\mathcal{U} * \mathcal{V}$ is again an ultrafilter, and that the operation $*$ is associative. The crucial property of $*$ is the following:

Proposition 5.4. For all \mathcal{U}, \mathcal{V} we have $\delta_{\mathcal{U}} \circ \delta_{\mathcal{V}} = \delta_{\mathcal{U} * \mathcal{V}}$.

Proof. For a set X , let

$$X^\sharp = \{\langle \langle i, j \rangle, k \rangle : \langle i, \langle j, k \rangle \rangle \in X\}.$$

We claim that $\lim_{\mathcal{U} * \mathcal{V}}(X^\sharp) = \lim_{\mathcal{U}}(\lim_{\mathcal{V}}(X))$, as follows:

$$\begin{aligned} x \in \lim_{\mathcal{U}}(\lim_{\mathcal{V}}(X)) &\iff \{j : \langle j, x \rangle \in \lim_{\mathcal{V}}(X)\} \in \mathcal{U} \iff \{j : \{k : \langle k, \langle j, x \rangle \rangle \in X\} \in \mathcal{V}\} \in \mathcal{U} \\ &\iff \{j : \{k : \langle \langle k, j \rangle, x \rangle \in X^\sharp\} \in \mathcal{V}\} \in \mathcal{U} \iff \{\langle k, j \rangle : \langle \langle k, j \rangle, x \rangle \in X^\sharp\} \in \mathcal{U} * \mathcal{V} \iff x \in \lim_{\mathcal{U} * \mathcal{V}}(X^\sharp). \end{aligned}$$

Since the operation \sharp is invertible and preserves Turing degree, we have shown that $\delta_{\mathcal{U}} \circ \delta_{\mathcal{V}}(\mathbf{a}) = \delta_{\mathcal{U} * \mathcal{V}}(\mathbf{a})$ for every degree \mathbf{a} . \square

Remark 5.5. Note that Proposition 5.4 only holds on the level of *degrees*: In general, given ultrafilters \mathcal{U} and \mathcal{V} and a set X there need be no ultrafilter \mathcal{W} with $\lim_{\mathcal{W}}(X) = \lim_{\mathcal{U}}(\lim_{\mathcal{V}}(X))$. For example, take $X = (X_i)_{i \in \omega}$ where each $X_i = \{0\}$. Then $\lim_{\mathcal{W}}(X) = \{0\}$ and $\lim_{\mathcal{U}}(\lim_{\mathcal{V}}(X)) = \emptyset$ regardless of the choice of $\mathcal{U}, \mathcal{V}, \mathcal{W}$.

Proposition 5.4 immediately yields:

Corollary 5.6. The operation $*$ is compatible with \mathcal{D}_{ult} : if $\mathcal{U}_0 \leq \mathcal{U}_1$ and $\mathcal{V}_0 \leq \mathcal{V}_1$, then $\mathcal{U}_0 * \mathcal{V}_0 \leq \mathcal{U}_1 * \mathcal{V}_1$. Moreover, $*$ is well-defined on elements of \mathcal{D}_{ult} .

Proposition 5.4 also provides us with a “jump” operator on \mathcal{D}_{ult} :

Definition 5.7. For \mathcal{U} an ultrafilter, let $\mathcal{U}' = \mathcal{U} * \mathcal{U}$.

Proposition 5.8. For every \mathcal{U} we have $\mathcal{U} < \mathcal{U}'$.

Proof. This is a refinement of Corollary 4.2. That $\mathcal{U} \leq \mathcal{U}'$ is immediate. To show that this is strict, fix a sufficiently large degree \mathbf{a} and suppose $\mathcal{U}' \leq \mathcal{U}$. Then we have (using Lemma 2.1(1) for the first equality)

$$\delta_{\mathcal{U}}(\mathbf{a}') \subseteq \delta_{\mathcal{U}'}(\mathbf{a}) \subseteq \delta_{\mathcal{U}}(\mathbf{a}),$$

contradicting the relativized version of Proposition 4.1. \square

This natural algebraic structure, compatible with the preorder, suggests that \mathcal{D}_{ult} may be an interesting degree structure in its own right. We end by providing further evidence for this: a connection between \mathcal{D}_{ult} and a more classical ordering of ultrafilters, the *Rudin-Keisler ordering*:

Definition 5.9. For \mathcal{U}, \mathcal{V} ultrafilters, \mathcal{U} is Rudin-Keisler reducible to \mathcal{V} , written $\mathcal{U} \leq_{RK} \mathcal{V}$, if for some $f: \omega \rightarrow \omega$ we have

$$\mathcal{U} = f^{-1}(\mathcal{V}), \text{ that is, } X \in \mathcal{V} \iff f^{-1}(X) \in \mathcal{U}.$$

We write $\mathcal{U} \leq_{RK}^f \mathcal{V}$ if f witnesses $\mathcal{U} \leq_{RK} \mathcal{V}$.

The connection between Rudin-Keisler reducibility and our \mathcal{D}_{ult} is provided by the following:

Theorem 5.10. Suppose $\mathcal{U} \leq_{RK}^f \mathcal{V}$. Then if $f \leq_T \mathbf{a}$, we have $\delta_{\mathcal{U}}(\mathbf{a}) \subseteq \delta_{\mathcal{V}}(\mathbf{a})$.

Proof. Given $X = (X_i)_{i \in \omega} \in \mathbf{a}$, define $Y = (Y_i)_{i \in \omega}$ where $Y_i = X_{f(i)}$. Now by our assumption on f we have

$$n \in \lim_{\mathcal{V}}(Y) \iff \{i : n \in X_{f(i)}\} \in \mathcal{V} \iff \{i : n \in X_i\} \in \mathcal{U} \iff n \in \lim_{\mathcal{U}}(X).$$

But this means $\delta_{\mathcal{U}}(\mathbf{a}) \subseteq \delta_{\mathcal{V}}(\mathbf{a})$. □

Corollary 5.11. If $\mathcal{U} \leq_{RK} \mathcal{V}$, then $\mathcal{U} \leq \mathcal{V}$.

Given this connection between \mathcal{D}_{ult} and the Rudin-Keisler ordering, it is natural to ask:

Question 9. Is there a characterization of \leq in terms of combinatorial properties of ultrafilters? In particular, does \leq coincide with \leq_{RK} ?

Almost certainly the answer is “no,” but it is not clear how to construct a counterexample.

6. FURTHER DIRECTIONS

We end by presenting two directions for further research.

6.1. Filter jumps. We have investigated maps $\delta_{\mathcal{U}}$ for \mathcal{U} an ultrafilter. However, this construction applies equally well to filters:

Definition 6.1. A *filter* is a collection of sets $\mathcal{F} \subseteq \mathcal{P}(\omega)$ satisfying conditions (1)-(3) of Definition 1.2. For \mathcal{F} a filter and $A = (A_i)_{i \in \omega}$ a sequence of sets, set $\lim_{\mathcal{F}}(A) = \{i : A^i \in \mathcal{F}\}$; then for \mathbf{a} a Turing ideal, define

$$\delta_{\mathcal{F}}(\mathbf{a}) = \{\lim_{\mathcal{F}}(A) : A \leq_T \mathbf{a}\}.$$

In order to preserve some of the analogy with classical limit computability, we demand that the filters we consider contain the Fréchet filter \mathcal{F}_{cof} of cofinite sets; call such filters *nice*.

Intuitively, this is a more “biased” notion of limit computability, since it is in general easier to have $X \notin \mathcal{F}$ than to have $X \in \mathcal{F}$. This is reflected in the fact that, in general, the resulting “filter jumps” $\delta_{\mathcal{F}}$ — while they may correspond to natural computability-theoretic operations — do not always yield Turing ideals. For example, $\delta_{\mathcal{F}_{cof}}(\mathbf{a}) = \Sigma_2^0(\mathbf{a})$, which is not closed under Turing reduction. On the positive side, note that $\delta_{\mathcal{F}}(\mathbf{a})$ is always closed under \oplus , and (as long as \mathcal{F} is nice) the limit lemma immediately implies that $\delta_{\mathcal{F}}(\mathbf{a}) \supseteq \Delta_2^0(\mathbf{a})$. Beyond this, however, it seems difficult to establish how these more general operations behave, and so the question of characterizing the possible images of filter jumps, in analogy with Theorem 3.1, is open:

Question 10. Fix a Turing degree \mathbf{a} . For what classes $\mathcal{I} \subseteq \mathcal{P}(\omega)$ is there some nice filter \mathcal{F} with $\delta_{\mathcal{F}}(\mathbf{a}) = \mathcal{I}$?

In particular, ensuring closure under Turing reducibility appears difficult.

Question 11. What filters \mathcal{F} have the property that $\delta_{\mathcal{F}}(\mathbf{a})$ is a Turing ideal for all \mathbf{a} ?

Moving on to trying to control the action of $\delta_{\mathcal{F}}$, we can (as in section 4) define a degree \mathbf{a} to be *f-low* if there is a nice filter \mathcal{F} such that $\delta_{\mathcal{F}}(\mathbf{a}) = \delta_{\mathcal{F}}(\mathbf{0})$. Clearly *f-lowness* is implied by *u-lowness*, and DNR_2 degrees are not *f-low*.

Question 12. Which degrees are *f-low*?

6.2. Ultrafilter jumps of Turing ideals. Our definition of $\delta_{\mathcal{U}}$ makes sense, not just for degrees, but for Turing ideals:

Definition 6.2. For \mathcal{I} a countable Turing ideal, let $\delta_{\mathcal{U}}(\mathcal{I}) = \{\lim_{\mathcal{U}}(A) : A \in \mathcal{I}\}$.

Then we can develop the theory of ultrafilter jumps in this broader context. By and large, the resulting picture is the same. Most importantly, by essentially the same proof as Theorem 3.1, we obtain:

Corollary 6.3. *Suppose \mathcal{I} is a countable Turing ideal and $\mathfrak{K} \supseteq \mathcal{I}$. Then the following are equivalent:*

- \mathfrak{K} is a countable Scott set containing \mathbf{a}' for every $\mathbf{a} \in \mathcal{I}$.
- There is an ultrafilter \mathcal{U} with $\delta_{\mathcal{U}}(\mathcal{I}) = \mathfrak{K}$.

There are, however, slight differences. For example, note that when generalized to ideals, Question 6 has a simple negative answer:

Proposition 6.4. *Let \mathcal{U}, \mathcal{V} be ultrafilters, and fix a Turing ideal \mathcal{I} . Then there is an ideal $\mathfrak{K} \supseteq \mathcal{I}$ such that $\delta_{\mathcal{U}}(\mathfrak{K}) = \mathfrak{K} = \delta_{\mathcal{V}}(\mathfrak{K})$.*

Proof. We alternately apply $\delta_{\mathcal{U}}$ and $\delta_{\mathcal{V}}$ infinitely many times to \mathcal{I} . Let $\mathcal{I}_0 = \mathcal{I}$, $\mathcal{I}_{n+1} = \delta_{\mathcal{U} * \mathcal{V}}(\mathcal{I}_n)$, and let

$$\mathfrak{K} = \bigcup_{i \in \omega} \mathcal{I}_i.$$

It is easy to check that \mathfrak{K} satisfies the desired properties. □

Having already generalized to countable Turing ideals, we can further consider the question of characterizing $\delta_{\mathcal{U}}(\mathcal{I})$ for *uncountable* Turing ideals \mathcal{I} . To a large extent, the possible behavior of ultrafilter jumps on uncountable ideals is already determined by their possible behavior on countable ideals, and even on individual degrees. However, the proof of Theorem 3.1 relied on enumerating the Turing ideals in question, and so breaks down as soon as we pass to uncountable ideals. This raises the question of whether our characterization still holds for uncountable ideals, and furthermore, to what extent the answer to this question depends on the axioms of set theory.

In general, this is unknown. However, at least a certain amount of set-theoretic independence does occur. Write “ $\mathcal{I} \prec 2^\omega$ ” if the structure $(\omega, \mathcal{I}; 0, 1, +, \times, \in)$ is an elementary substructure of $(\omega, 2^\omega; 0, 1, +, \times, \in)$. For example, if $\mathcal{I} \prec 2^\omega$ then \mathcal{I} is arithmetically closed, since “ $X = Y$ ” is definable in the language of second-order arithmetic. Then we have:

Theorem 6.5 (Schweber). *It is independent of ZFC whether sufficiently nice Turing ideals are always fixed by some $\delta_{\mathcal{U}}$:*

- Consistently with ZFC (in fact, provably from $V=L$), for any $\mathcal{I} \prec 2^\omega$ there is some ultrafilter with $\delta_{\mathcal{U}}(\mathcal{I}) = \mathcal{I}$.
- If V satisfies projective determinacy PD, then there is a forcing extension W of V with the same reals as V and a $\mathcal{I} \in W$ such that $\mathcal{I} \prec 2^\omega$ but no ultrafilter \mathcal{U} satisfies $\delta_{\mathcal{U}}(\mathcal{I}) = \mathcal{I}$.

The first part follows immediately from the fact that $V=L$ implies the existence of definable ultrafilters; the second uses forcing to build the desired Turing ideal, with projective determinacy needed to guarantee that appropriate extensions of conditions always exist. The full proof of Theorem 6.5 appeared in the fourth author’s Ph.D. thesis [Sch16, Theorem 7.6.5]. However, note that this result still falls far short of understanding the extension of Theorem 3.1 to uncountable Turing ideals.

Question 13. Is it consistent with ZFC that every *arithmetically closed* Turing ideal is fixed by some $\delta_{\mathcal{U}}$?

Question 14. Does ZFC+PD *prove* that there is an $\mathfrak{J} < 2^\omega$ not fixed by any $\delta_{\mathcal{U}}$?

REFERENCES

- [CN74] W. W. Comfort and S. Negrepointis. *The theory of ultrafilters*. Springer-Verlag, New York-Heidelberg, 1974. Die Grundlehren der mathematischen Wissenschaften, Band 211.
- [DH10] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic randomness and complexity*. Theory and Applications of Computability. Springer, New York, 2010.
- [Jec03] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded.
- [Nie06] André Nies. Reals which compute little. In *Logic Colloquium '02*, volume 27 of *Lect. Notes Log.*, pages 261–275. Assoc. Symbol. Logic, La Jolla, CA, 2006.
- [Rob66] Robert William Robinson. *The Inclusion Lattice and Degrees of Unsolvability of the Recursively Enumerable Sets*. ProQuest LLC, Ann Arbor, MI, 1966. Thesis (Ph.D.)—Cornell University.
- [Sch16] Noah D. Schweber. *Interactions between computability theory and set theory*. ProQuest LLC, Ann Arbor, MI, 2016. Thesis (Ph.D.)—University of California, Berkeley.
- [Sim99] Stephen G. Simpson. *Subsystems of second order arithmetic*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1999.
- [Soa87] Robert I. Soare. *Recursively enumerable sets and degrees*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1987. A study of computable functions and computably generated sets.